




Cuttack District Block Wise Agro-met Advisory Service Bulletin

*Tuesday, 04th November, 2025
(For the period 04th to 08th November,
2025)*



*Issued by
District Agro-met Unit,
KrishiVigyan Kendra Cuttack*






GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-602
Date- 04/11/2025

STANDARD PRECIPITATION INDEX AND NDVI MAP



LEGEND (Medium range block level rainfall forecast)		
	No Rainfall	00 mm
	Very light rainfall	01-05 mm
	Light rainfall	06-10 mm
	Light to moderate rainfall	11-30 mm
	Moderate to heavy rainfall	31-65 mm
	Heavy to very heavy rainfall	66-115 mm
	Extremely heavy rainfall	>116 mm

LEGEND (Past week temperature)		
Color code	Temperature	Departure
	Below Normal	< 2°C from normal
	Normal	-2°C to +2°C from normal
	Above Normal	>2°C from normal



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-602
Date- 04/11/2025

Weather (Past and Forecasted) based Disease and pest in Major Crops and vegetables

Sl No	Crop	Stage and condition of the crop	Anticipated Disease and pest	
1	Rice	Panicle initiation to Milk dough stage		Leaf folder
				BPH
				Gundhi bug
				False smut
				Bacterial leaf streak
2	Sugarcane	Grand growth stage to cane development stag		Ring spot
				Top borer
				Woolly Aphid
				Scale insect
				White fly
				Mealy bug
3	Vegetables(Brinjal and okra)	Vegetative to fruiting		Fruit and shoot borer
	Okra			YMV
	Chilli , Cowpea			Thrips in Chilli and Aphids in Cowpea



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area





GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-602

Date- 04/11/2025

District Past week average weather condition and Block past week average Rainfall condition

The block received average 16 mm and district received average 10 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.28°C (-1.29 from normal) and 22°C (+1.46 from normal), respectively. The mean relative humidity was 87-92%.

Forecasted period: Today(8.30IST) to 09th November (8.30IST)2025

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 31-32°C and night temperature will be 20-21°C for upcoming 5 days. Maximum and minimum relative humidity will be 70-83% and 38-54% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Athagad (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	04/11/2025	05/11/2025	06/11/2025	07/11/2025	08/11/2025
Rainfall (mm)	00	00	00	00	00
T-Max (°C)	32	31	31	31	31
T-Min (°C)	21	21	21	20	20
Cloud Cover (octa)	4	3	3	3	2
RH Max (%)	70	74	75	83	79
RH Min (%)	38	42	46	54	53
Wind Speed (kmph)	5	5	6	8	8
Wind Direction (deg)	344	321	323	329	338

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to milk dough stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Keep watch on the infestation of gundhi bug and BPH in paddy. Complete nursery sowing for winter vegetable crops. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.

Kharif Rice:

Paddy is at PI to flowering stage. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of Leaf Folder. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of BPH & WBPH occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from Gundhi bug apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of false smut spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water. In the event of bacterial leaf streak i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + copper oxychloride 1g per liter water only in the clear weather condition.

PULSES:

21-26°C soil temperature is required for proper germination of pulses. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT:

As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal. Apply irrigation in field if mustard was sown before 15-20 days.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. As past and forecasted temperature is around 20-30°C, so better to cultivate the offseason cauliflowers in protected structure to maintain the temperature. To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @70ml/acre and Thiodicarb@300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry: Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-602
Date- 04/11/2025

AGROMET ADVISORY BULLETIN FOR BADAMBA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 07 mm and district received average 10 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.28°C (-1.29 from normal) and 22°C (+1.46 from normal), respectively. The mean relative humidity was 87-92%.

Forecasted period: Today(8.30IST) to 09th November (8.30IST)2025

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 31-32°C and night temperature will be 20-21°C for upcoming 5 days. Maximum and minimum relative humidity will be 72-82% and 35-53% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Badamba (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	04/11/2025	05/11/2025	06/11/2025	07/11/2025	08/11/2025
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	32	31	31	31	31
T-Min(°C)	21	21	21	20	20
Cloud Cover(octa)	4	3	3	3	2
RH Max (%)	74	72	73	82	79
RH Min (%)	35	42	44	53	51
Wind Speed (Kmph)	5	4	5	6	6
Wind Direction (deg)	330	326	321	332	334

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to milk dough stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Keep watch on the infestation of gundhi bug and BPH in paddy. Complete nursery sowing for winter vegetable crops. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.

Kharif Rice:

Paddy is at PI to flowering stage. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem baesd insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC@ 2ml/liter of water. In the event of **bacterial leaf streak** i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + copper oxychloride 1g per liter water only in the clear weather condition.

PULSES:

21-26°C soil temperature is required for proper germination of pulses. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT:

As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal. Apply irrigation in field if mustard was sown before 15-20 days.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. As past and forecasted temperature is around 20-30°C, so better to cultivate the offseason cauliflowers in protected structure to maintain the temperature. To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @70ml/acre and Thiodicarb@300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry: Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-602
Date- 04/11/2025

AGROMET ADVISORY BULLETIN FOR BANKI

District Past week average weather condition and Block past week average Rainfall condition

The district received average 10 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.28°C (-1.29 from normal) and 22°C (+1.46 from normal), respectively. The mean relative humidity was 87-92%.

Forecasted period: Today(8.30IST) to 09th November (8.30IST)2025

Weatheradvisory:: Dry weather is forecasted for upcoming 5days in the block. The day temperature will be 31-32°C and night temperature will be 20-21°C for upcoming 5 days. Maximum and minimum relative humidity will be 72-82% and 35-54% respectively. **There will be clear to partly cloudy sky for next 5days.**

Block	Banki(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	04/11/2025	05/11/2025	06/11/2025	07/11/2025	08/11/2025
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	32	31	31	31	31
T-Min(°C)	21	21	21	20	20
Cloud Cover(octa)	4	3	3	3	2
RH Max (%)	72	73	75	82	77
RH Min (%)	35	42	45	54	52
Wind Speed (Kmph)	6	6	6	7	8
Wind Direction (deg)	335	328	325	331	336

Agromet Advisory

General Advisory: **Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to milk dough stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Keep watch on the infestation of gundhi bug and BPH in paddy. Complete nursery sowing for winter vegetable crops. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.**

Kharif Rice:

Paddy is at PI to flowering stage. **As dry weather is forecasted for upcoming 5 days , in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem baesd insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC@ 2ml/liter of water. In the event of **bacterial leaf streak** i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + cupper oxychloride 1g per liter water only in the clear weather condition.

PULSES:

21-26°C soil temperature is required for proper germination of pulses. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT:

As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal. **Apply irrigation in field if mustard was sown before 15-20 days.**

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. **In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. As past and forecasted temperature is around 20-30°C, so better to cultivate the offseason cauliflowers in protected structure to maintain the temperature.** To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @600ml/acre at initial stages. Later ongo for alternate spray of Spinosad 45% SC @70ml/acre and Thiodicarb@300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry: Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





**GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK**

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-602

Date- 04/11/2025

AGROMET ADVISORY BULLETIN FOR BANKI-DAMPARA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 16 mm and district received average 10 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.28°C (-1.29 from normal) and 22°C (+1.46 from normal), respectively. The mean relative humidity was 87-92%.

Forecasted period: Today(8.30IST) to 09th November (8.30IST)2025

Weather Advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 31-32°C and night temperature will be 20-21°C for upcoming 5 days. Maximum and minimum relative humidity will be 72-82% and 35-54% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Banki-Dampara (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	04/11/2025	05/11/2025	06/11/2025	07/11/2025	08/11/2025
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	32	31	31	31	31
T-Min(°C)	21	21	21	20	20
Cloud Cover(octa)	4	3	3	3	2
RH Max (%)	72	73	75	82	77
RH Min (%)	35	42	45	54	52
Wind Speed (Kmph)	6	6	6	7	8
Wind Direction (deg)	335	328	325	331	336

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to milk dough stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Keep watch on the infestation of gundhi bug and BPH in paddy. Complete nursery sowing for winter vegetable crops. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.

Kharif Rice:

Paddy is at PI to flowering stage. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water. In the event of **bacterial leaf streak** i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + copper oxychloride 1g per liter water only in the clear weather condition.

PULSES:

21-26°C soil temperature is required for proper germination of pulses. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT:

As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal. Apply irrigation in field if mustard was sown before 15-20 days.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. As past and forecasted temperature is around 20-30°C, so better to cultivate the offseason cauliflowers in protected structure to maintain the temperature. To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @70ml/acre and Thiodicarb@300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry: Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



**DrRanjan Kumar Mohanta
(Nodal officer)**

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-602
Date- 04/11/2025

AGROMET ADVISORY BULLETIN FOR BARANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 07 mm and district received average 10 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.28°C (-1.29 from normal) and 22°C (+1.46 from normal), respectively. The mean relative humidity was 87-92%.

Forecasted period: Today(8.30IST) to 09th November (8.30IST)2025

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 31-32°C and night temperature will be 20-21°C for upcoming 5 days. Maximum and minimum relative humidity will be 73-84% and 39-55% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Baranga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	04/11/2025	05/11/2025	06/11/2025	07/11/2025	08/11/2025
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	32	31	31	31	31
T-Min(°C)	21	21	21	20	20
Cloud Cover	4	3	3	3	2
RH Max (%)	73	75	77	84	78
RH Min (%)	39	43	47	55	55
Wind Speed (Kmph)	6	5	7	8	9
Wind Direction (deg)	350	334	328	338	346

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to milk dough stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Keep watch on the infestation of gundhi bug and BPH in paddy. Complete nursery sowing for winter vegetable crops. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.

Kharif Rice:

Paddy is at PI to flowering stage. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of Leaf Folder. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of BPH & WBPH occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from Gundhi bug apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of false smut spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water. In the event of bacterial leaf streak i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + copper oxychloride 1g per liter water only in the clear weather condition.

PULSES:

21-26°C soil temperature is required for proper germination of pulses. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT:

As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal. Apply irrigation in field if mustard was sown before 15-20 days.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. As past and forecasted temperature is around 20-30°C, so better to cultivate the offseason cauliflowers in protected structure to maintain the temperature. To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @70ml/acre and Thiodicarb @300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry: Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR CUTTACK SADAR



Bulletin No.-602

Date- 04/11/2025

District Past week average weather condition and Block past week average Rainfall condition

The block received average 04 mm and district received average 10 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.28°C (-1.29 from normal) and 22°C (+1.46 from normal), respectively. The mean relative humidity was 87-92%.

Forecasted period: Today(8.30IST) to 09th November (8.30IST)2025

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 31°C and night temperature will be 20-21°C for upcoming 5 days. Maximum and minimum relative humidity will be 68-85% and 40-54% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Cuttack sadar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	04/11/2025	05/11/2025	06/11/2025	07/11/2025	08/11/2025
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	31	31	31	31	31
T-Min(°C)	21	21	21	20	20
Cloud Cover(octa)	4	3	3	3	2
RH Max (%)	68	74	82	85	77
RH Min (%)	40	43	47	54	53
Wind Speed (Kmph)	6	4	6	7	8
Wind Direction (deg)	3	329	338	342	5

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to milk dough stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Keep watch on the infestation of gundhi bug and BPH in paddy. Complete nursery sowing for winter vegetable crops. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.

Kharif Rice: Paddy is at PI to flowering stage. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of Leaf Folder. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of BPH & WBPH occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from Gundhi bug apply 5 ml neem baesd insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of false smut spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water. In the event of bacterial leaf streak i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + cupper oxychloride 1g per liter water only in the clear weather condition.

Sugarcane: Remove the borer affected tillers and late formed tillers, tie the cane shoot with two or three together with partially dried lower leaves. Present weather conditions are favourable for the incidence of ring spot disease in Sugarcane. To control go for spraying of copperoxychloride @ 3g/l (or) Carbendazim @ 1g/l (or) Mancozeb @ 3g/l for three times once in three weeks. To control top shoot borers spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water. For control of Sucking pest in sugarcane like Woolly aphid, Whitefly, Scale insects, mealybug in present weather conditions go for alternate spraying of thiomethoxam 25WG @ 0.25 gm/lit or imidacloprid 0.45 ml/lit of water or Dinotefuron 20% SG 3gm/lit of Water.

PULSES: 21-26°C soil temperature is required for proper germination of pulses. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT: As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard: Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @ 60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal. Apply irrigation in field if mustard was sown before 15-20 days.

VEGETABLES: Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. As past and forecasted temperature is around 20-30°C, so better to cultivate the offseason cauliflowers in protected structure to maintain the temperature. To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry: Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-602
Date- 04/11/2025

AGROMET ADVISORY BULLETIN FOR KANTAPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 11 mm and district received average 10 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.28°C (-1.29 from normal) and 22°C (+1.46 from normal), respectively. The mean relative humidity was 87-92%.

Forecasted period: Today(8.30IST) to 09th November (8.30IST)2025

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 31-32°C and night temperature will be 20-21°C for upcoming 5 days. Maximum and minimum relative humidity will be 71-87% and 41-55% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Kantapada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	04/11/2025	05/11/2025	06/11/2025	07/11/2025	08/11/2025
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	32	31	31	31	31
T-Min(°C)	21	21	21	21	20
Cloud Cover(octa)	4	3	3	3	2
RH Max (%)	71	78	81	87	81
RH Min (%)	41	45	48	55	55
Wind Speed (Kmph)	7	6	7	8	9
Wind Direction (deg)	351	337	338	345	4

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to milk dough stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Keep watch on the infestation of gundhi bug and BPH in paddy. Complete nursery sowing for winter vegetable crops. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.

Kharif Rice: Paddy is at PI to flowering stage. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water. In the event of **bacterial leaf streak** i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + copper oxychloride 1g per liter water only in the clear weather condition.

Sugarcane: Remove the borer affected tillers and late formed tillers, tie the cane shoot with two or three together with partially dried lower leaves. Present weather conditions are favourable for the incidence of **ring spot disease** in Sugarcane. To control go for spraying of copperoxychloride @ 3g/l (or) Carbendazim @ 1g/l (or) Mancozeb @ 3g/l for three times once in three weeks. To control **top shoot borers** spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water. For control of Sucking pest in sugarcane like **Woolly aphid, Whitefly, Scale insects, mealybug** in present weather conditions go for alternate spraying of thiomethoxam 25WG @ 0.25 gm/lit or imidacloprid 0.45 ml/lit of water or Dinotefuron 20% SG 3gm/lit of Water.

PULSES: 21-26°C soil temperature is required for proper germination of pulses. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT: As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard: Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @ 60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal. Apply irrigation in field if mustard was sown before 15-20 days.

VEGETABLES: Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. As past and forecasted temperature is around 20-30°C, so better to cultivate the offseason cauliflowers in protected structure to maintain the temperature. To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry: Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





**GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK**

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR MAHANGA



Bulletin No.-602

Date- 04/11/2025

District Past week average weather condition and Block past week average Rainfall condition

The block received average 10 mm and district received average 10 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.28°C (-1.29 from normal) and 22°C (+1.46 from normal), respectively. The mean relative humidity was 87-92%.

Forecasted period: Today(8.30IST) to 09th November (8.30IST)2025

Weatheradvisory: Dry weather is forecasted for upcoming 5days in the block. The day temperature will be and it will be 31-32°C and night temperature will be 20-21°C for upcoming 5 days. Maximum and minimum relative humidity will be 66-86% and 42-54% respectively. **There will be clear to partly cloudy sky for next 5days.**

Block	Mahanga(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	04/11/2025	05/11/2025	06/11/2025	07/11/2025	08/11/2025
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	32	32	31	31	31
T-Min(°C)	21	21	21	21	20
Cloud Cover(octal)	4	3	3	3	2
RH Max (%)	66	81	86	86	78
RH Min (%)	42	45	48	54	54
Wind Speed (Kmph)	8	5	7	7	11
Wind Direction (deg)	90	334	345	349	90

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to milk dough stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Keep watch on the infestation of gundhi bug and BPH in paddy. Complete nursery sowing for winter vegetable crops. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.

Kharif Rice: Paddy is at PI to flowering stage. **As dry weather is forecasted for upcoming 5 days , in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem baesd insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC@ 2ml/liter of water. In the event of **bacterial leaf streak** i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + cupper oxychloride 1g per liter water only in the clear weather condition.

Sugarcane: Remove the borer affected tillers and late formed tillers, tie the cane shoot with two or three together with partially dried lower leaves. Present weather conditions are favourable for the incidence of **ring spot disease** in Sugarcane. To control go for spraying of copperoxychloride @ 3g/l (or) Carbendazim @ 1g/l (or) Mancozeb @3g/l for three times once in three weeks. To control **top shoot borers** spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water For control of Sucking pest in sugarcane like **Wooly aphid, Whitefly, Scale insects, mealybug** in present weather conditions go for alternate spraying of thiomethoxam 25WG @ 0.25 gm/lit or imidacloprid 0.45 ml/lit of water or Dinotefuron 20% SG 3gm/lit of Water.

PULSES: 21-26°C soil temperature is required for proper germination of pulses. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT: As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard: Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal. **Apply irrigation in field if mustard was sown before 15-20 days.**

VEGETABLES: Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. **In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. As past and forecasted temperature is around 20-30°C, so better to cultivate the offseason cauliflowers in protected structure to maintain the temperature.** To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @70ml/acre and Thiodicarb@300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry: Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



**DrRanjan Kumar Mohanta
(Nodal officer)**

**Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area**





GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-602
Date- 04/11/2025

AGROMET ADVISORY BULLETIN FOR NARASINGHPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 11 mm and district received average 10 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.28°C (-1.29 from normal) and 22°C (+1.46 from normal), respectively. The mean relative humidity was 87-92%.

Forecasted period: Today(8.30IST) to 09th November (8.30IST)2025

Weatheradvisory:: *Dry weather is forecasted for upcoming 5 days in the block.* The day temperature will be 31°C and night temperature will be 19-21°C for upcoming 5 days. Maximum and minimum relative humidity will be 73-87% and 33-51% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Narasinghpur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	04/11/2025	05/11/2025	06/11/2025	07/11/2025	08/11/2025
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	31	31	31	31	31
T-Min(°C)	21	21	21	20	19
Cloud Cover(/octa)	3	2	3	3	2
RH Max (%)	77	73	76	87	82
RH Min (%)	33	41	43	51	50
Wind Speed (Kmph)	5	3	4	5	4
Wind Direction (deg)	328	319	323	334	326

Agromet Advisory

General Advisory: *Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to milk dough stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Keep watch on the infestation of gundhi bug and BPH in paddy. Complete nursery sowing for winter vegetable crops. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.*

Kharif Rice:

Paddy is at PI to flowering stage. **As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water. In the event of **bacterial leaf streak** i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + copper oxychloride 1g per liter water only in the clear weather condition.

PULSES:

21-26°C soil temperature is required for proper germination of pulses. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT:

As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal. **Apply irrigation in field if mustard was sown before 15-20 days.**

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. **In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. As past and forecasted temperature is around 20-30°C, so better to cultivate the offseason cauliflowers in protected structure to maintain the temperature.** To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @70ml/acre and Thiodicarb@300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry: Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area





GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-602
Date- 04/11/2025

AGROMET ADVISORY BULLETIN FOR NIALI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 12 mm and district received average 10 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.28°C (-1.29 from normal) and 22°C (+1.46 from normal), respectively. The mean relative humidity was 87-92%.

Forecasted period: Today(8.30IST) to 09th November (8.30IST)2025

Weather advisory: *Dry weather is forecasted for upcoming 5 days in the block.* The day temperature will be 32°C and night temperature will be 21-22°C for upcoming 5 days. Maximum and minimum relative humidity will be 71-87% and 40-57% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Niali (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	04/11/2025	05/11/2025	06/11/2025	07/11/2025	08/11/2025
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	32	32	32	32	32
T-Min(°C)	22	22	22	21	21
Cloud Cover(octa)	4	3	3	4	3
RH Max (%)	71	82	79	87	81
RH Min (%)	40	46	48	55	57
Wind Speed(Kmph)	8	6	8	9	11
Wind Direction(deg)	338	332	341	346	356

Agromet Advisory

General Advisory: *Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to milk dough stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Keep watch on the infestation of gundhi bug and BPH in paddy. Complete nursery sowing for winter vegetable crops. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.*

Kharif Rice: Paddy is at PI to flowering stage. **As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water. In the event of **bacterial leaf streak** i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + copper oxychloride 1g per liter water only in the clear weather condition.

Sugarcane: Remove the borer affected tillers and late formed tillers, tie the cane shoot with two or three together with partially dried lower leaves. Present weather conditions are favourable for the incidence of **ring spot disease** in Sugarcane. To control go for spraying of copperoxychloride @ 3g/l (or) Carbendazim @ 1g/l (or) Mancozeb @ 3g/l for three times once in three weeks. To control **top shoot borers** spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water. For control of Sucking pest in sugarcane like **Woolly aphid, Whitefly, Scale insects, mealybug** in present weather conditions go for alternate spraying of thiomethoxam 25WG @ 0.25 gm/lit or imidacloprid 0.45 ml/lit of water or Dinotefuron 20% SG 3gm/lit of Water.

PULSES: **21-26°C soil temperature is required for proper germination of pulses.** This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT: **As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done** if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard: Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @ 60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal. **Apply irrigation in field if mustard was sown before 15-20 days.**

VEGETABLES: Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. **In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. As past and forecasted temperature is around 20-30°C, so better to cultivate the offseason cauliflowers in protected structure to maintain the temperature.** To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry: Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area





GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)



Bulletin No.-602
Date- 04/11/2025

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR NISCHINTAKOILI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 11 mm and district received average 10 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.28°C (-1.29 from normal) and 22°C (+1.46 from normal), respectively. The mean relative humidity was 87-92%.

Forecasted period: Today(8.30IST) to 09th November (8.30IST)2025

Weatheradvisory: Dry weather is forecasted for upcoming 5days in the block. The day temperature will be 31-32°C and night temperature will be 20-21°C for upcoming 5 days. Maximum and minimum relative humidity will be 68-86% and 42-54% respectively. **There will be clear to partly cloudy sky for next 5days.**

Block	Nischintakoili(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	04/11/2025	05/11/2025	06/11/2025	07/11/2025	08/11/2025
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	32	32	31	31	31
T-Min(°C)	21	21	21	21	20
Cloud Cover(octa)	4	3	3	3	2
RH Max (%)	68	81	86	86	78
RH Min (%)	42	45	48	54	54
Wind Speed (Kmph)	8	5	7	8	12
Wind Direction (deg)	90	334	348	350	2

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to milk dough stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Keep watch on the infestation of gundhi bug and BPH in paddy. Complete nursery sowing for winter vegetable crops. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.

Kharif Rice: Paddy is at PI to flowering stage. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC@ 2ml/liter of water. In the event of **bacterial leaf streak** i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + cupper oxychloride 1g per liter water only in the clear weather condition.

Sugarcane: Remove the borer affected tillers and late formed tillers, tie the cane shoot with two or three together with partially dried lower leaves. Present weather conditions are favourable for the incidence of **ring spot disease** in Sugarcane. To control go for spraying of copperoxychloride @ 3g/l (or) Carbendazim @ 1g/l (or) Mancozeb @3g/l for three times once in three weeks. To control **top shoot borers** spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water For control of Sucking pest in sugarcane like **Woolly aphid, Whitefly, Scale insects, mealybug** in present weather conditions go for alternate spraying of thiomethoxam 25WG @ 0.25 gm/lit or imidacloprid 0.45 ml/lit of water or Dinotefuron 20% SG 3gm/lit of Water.

PULSES: 21-26°C soil temperature is required for proper germination of pulses. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT: As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard: Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal. **Apply irrigation in field if mustard was sown before 15-20 days.**

VEGETABLES: Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. As past and forecasted temperature is around 20-30°C, so better to cultivate the offseason cauliflowers in protected structure to maintain the temperature. To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @70ml/acre and Thiodicarb@300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry: Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-602
Date- 04/11/2025

AGROMET ADVISORY BULLETIN FOR SALEPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 11 mm and district received average 10 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.28°C (-1.29 from normal) and 22°C (+1.46 from normal), respectively. The mean relative humidity was 87-92%.

Forecasted period: Today(8.30IST) to 09th November (8.30IST)2025

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 31°C and night temperature will be 20-21°C for upcoming 5 days. Maximum and minimum relative humidity will be 68-86% and 42-54% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Salepur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	04/11/2025	05/11/2025	06/11/2025	07/11/2025	08/11/2025
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	31	31	31	31	31
T-Min(°C)	21	21	21	21	20
Cloud Cover(octa)	4	3	3	3	2
RH Max (%)	68	81	86	86	78
RH Min (%)	42	44	48	54	54
Wind Speed (Kmph)	8	5	7	8	11
Wind Direction (deg)	90	334	345	349	90

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to milk dough stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Keep watch on the infestation of gundhi bug and BPH in paddy. Complete nursery sowing for winter vegetable crops. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.

Kharif Rice: Paddy is at PI to flowering stage. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water. In the event of **bacterial leaf streak** i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + copper oxychloride 1g per liter water only in the clear weather condition.

Sugarcane: Remove the borer affected tillers and late formed tillers, tie the cane shoot with two or three together with partially dried lower leaves. Present weather conditions are favourable for the incidence of **ring spot disease** in Sugarcane. To control go for spraying of copperoxychloride @ 3g/l (or) Carbendazim @ 1g/l (or) Mancozeb @ 3g/l for three times once in three weeks. To control **top shoot borers** spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water. For control of Sucking pest in sugarcane like **Woolly aphid, Whitefly, Scale insects, mealybug** in present weather conditions go for alternate spraying of thiomethoxam 25WG @ 0.25 gm/lit or imidacloprid 0.45 ml/lit of water or Dinotefuron 20% SG 3gm/lit of Water.

PULSES: 21-26°C soil temperature is required for proper germination of pulses. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT: As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard: Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @ 60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal. Apply irrigation in field if mustard was sown before 15-20 days.

VEGETABLES: Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. As past and forecasted temperature is around 20-30°C, so better to cultivate the offseason cauliflowers in protected structure to maintain the temperature. To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry: Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area





GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)



Bulletin No.-602
Date- 04/11/2025

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR TANGI-CHOUDWAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 11 mm and district received average 10 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.28°C (-1.29 from normal) and 22°C (+1.46 from normal), respectively. The mean relative humidity was 87-92%.

Forecasted period: Today(8.30IST) to 09th November (8.30IST)2025

Weatheradvisory: Dry weather is forecasted for upcoming 5days in the block. The day temperature will be 31°C and night temperature will be 20-21°C for upcoming 5 days. Maximum and minimum relative humidity will be 67-84% and 45-61% respectively. **There will be clear to partly cloudy sky for next 5days.**

Block	Tangi-Choudwar(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	04/11/2025	05/11/2025	06/11/2025	07/11/2025	08/11/2025
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	31	31	31	31	31
T-Min(°C)	21	21	21	20	20
Cloud Cover(octa)	4	3	3	3	2
RH Max (%)	67	77	83	84	82
RH Min (%)	45	53	55	60	61
Wind Speed (Kmph)	12	8	9	11	13
Wind Direction (deg)	355	350	340	350	3

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to milk dough stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Keep watch on the infestation of gundhi bug and BPH in paddy. Complete nursery sowing for winter vegetable crops. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.

Kharif Rice:

Paddy is at PI to flowering stage. **As dry weather is forecasted for upcoming 5 days , in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem baesd insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC@ 2ml/liter of water. In the event of **bacterial leaf streak** i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + cupper oxychloride 1g per liter water only in the clear weather condition.

PULSES:

21-26°C soil temperature is required for proper germination of pulses. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT:

As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal. **Apply irrigation in field if mustard was sown before 15-20 days.**

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. **In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. As past and forecasted temperature is around 20-30°C, so better to cultivate the offseason cauliflowers in protected structure to maintain the temperature.** To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @600ml/acre at initial stages. Later ongo for alternate spray of Spinosad 45% SC @70ml/acre and Thiodicarb@300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry: Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-602
Date- 04/11/2025

AGROMET ADVISORY BULLETIN FOR TIGIRIA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 08 mm and district received average 10 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.28°C (-1.29 from normal) and 22°C (+1.46 from normal), respectively. The mean relative humidity was 87-92%.

Forecasted period: Today(8.30IST) to 09th November (8.30IST)2025

Weatheradvisory:: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 31-32°C and night temperature will be 20-21°C for upcoming 5 days. Maximum and minimum relative humidity will be 71-82% and 38-55% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Tigiria(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	04/11/2025	05/11/2025	06/11/2025	07/11/2025	08/11/2025
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	32	31	31	31	31
T-Min(°C)	21	21	21	20	20
Cloud Cover(octa)	4	3	3	3	2
RH Max (%)	71	71	73	82	79
RH Min (%)	38	42	45	55	53
Wind Speed (Kmph)	6	5	5	7	8
Wind Direction (deg)	327	324	326	331	332

Agro met Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to milk dough stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Keep watch on the infestation of gundhi bug and BPH in paddy. Complete nursery sowing for winter vegetable crops. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.

Kharif Rice:

Paddy is at PI to flowering stage. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC@ 2ml/liter of water. In the event of **bacterial leaf streak** i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + copper oxychloride 1g per liter water only in the clear weather condition.

PULSES:

21-26°C soil temperature is required for proper germination of pulses. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT:

As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal. Apply irrigation in field if mustard was sown before 15-20 days.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. As past and forecasted temperature is around 20-30°C, so better to cultivate the offseason cauliflowers in protected structure to maintain the temperature. To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @70ml/acre and Thiodicarb@300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry: Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area








GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-602
Date- 04/11/2025

				
MEGHDOOT APP	DAMINI APP	FACEBOOK (Damu Cuttack)	BLOCK SPECIFIC WHAT'S APP GROUPS	@DAMUCUTTACK

- Kindly install and use **“Meghdoot mobile App”** for Cuttack district weather forecast, warnings and weather based agromet advisory services.

<https://play.google.com/store/apps/details?id=com.aas.meghdoot>

- Kindly install and use **“Damini mobile App”** for location specific for lightening forecast and warnings.

<https://play.google.com/store/apps/details?id=com.lightening.live.damini>

- Follow the **block specific whats app groups** and join in your block wise groups for getting weather based agromet advisory services twice in a week (Tuesday and Friday) and other weather warning based farm management advisories.
- Follow the **Official Website links** to get the weather based agromet advisory service bulletins (**English & Odia**) twice in a week (Tuesday and Friday).

1	ICAR-NRRI, Cuttack, Odisha	https://icar-nrri.in/agromate/	
2	Agrimet division, IMD, Pune	https://imdagrmet.gov.in/imdproject/AGIndex.php	



DrRanjan Kumar Mohanta
 (Nodal officer)

**Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area**

