




Cuttack District Block Wise Agro-met Advisory Service Bulletin

*Tuesday, 11th November, 2025
(For the period 11th to 15th November,
2025)*



*Issued by
District Agro-met Unit,
KrishiVigyan Kendra Cuttack*






GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-604
Date- 11/11/2025

STANDARD PRECIPITATION INDEX AND NDVI MAP



LEGEND (Medium range block level rainfall forecast)		
	No Rainfall	00 mm
	Very light rainfall	01-05 mm
	Light rainfall	06-10 mm
	Light to moderate rainfall	11-30 mm
	Moderate to heavy rainfall	31-65 mm
	Heavy to very heavy rainfall	66-115 mm
	Extremely heavy rainfall	>116 mm

LEGEND (Past week temperature)		
Color code	Temperature	Departure
	Below Normal	< 2°C from normal
	Normal	-2°C to +2°C from normal
	Above Normal	>2°C from normal



DrRanjan Kumar Mohanta
 (Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK




(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-604

Date- 11/11/2025

Weather (Past and Forecasted) based Disease and pest in Major Crops and vegetables

Sl No	Crop	Stage and condition of the crop	Anticipated Disease and pest	
1	Rice	Milky grain stage to harvesting stage		Rice ear head bug
				Neck blast
				BPH
				Gundhi bug
2	Sugarcane	Grand growth stage to cane development stag		Wooly Aphid
				Scale insect
				White fly
				Mealy bug
3	Cole crops	Vegetative to fruiting		DBM
	Vegetables	Vegetative to fruiting		sucking pest(Aphid, Mites, Thrips)



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area





GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-604
Date- 11/11/2025

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30°C (-1.25 from normal) and 18°C (-1.77 from normal), respectively. The mean relative humidity was 72-79%.

Forecasted period: Today(8.30IST) to 16th November (8.30IST)2025

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 30-31°C and night temperature will be 15-16°C for upcoming 5 days. Maximum and minimum relative humidity will be 67-76% and 31-37% respectively. There will be clear sky for next 5 days.

Block	Athagad (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2025	12/11/2025	13/11/2025	14/11/2025	15/11/2025
Rainfall (mm)	00	00	00	00	00
T-Max (°C)	30	30	31	30	30
T-Min (°C)	15	15	16	16	15
Cloud Cover (octa)	0	0	0	0	1
RH Max (%)	76	74	70	70	67
RH Min (%)	37	33	32	31	32
Wind Speed (kmph)	8	7	7	7	6
Wind Direction (deg)	317	317	328	326	327

Agromet Advisory

General Advisory: Harvest matured paddy crops and store them in safer place after sun drying. Complete nursery sowing or transplanting of winter vegetable crops. Irrigate the nursery during morning and evening time. Use low cost polythene cover above the vegetables nursery to avoid high evapotranspiration. Go for sowing of pea, gram and green gram crop by utilizing residual soil moisture

Kharif Rice:

Paddy is grain formation to harvesting stage. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. To control **rice ear head bug**, go for spraying of Dichlorovos @ 1ml/l during evening hours. Spray the chemical from borders to middle of the field. Present weather conditions is favourable for the incidence of **Neck blast** disease. To control, go for spraying of tricyclazole @ 0.6 g/l or Carbendazim @ 1 g / L. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis.

PULSES:

21-26°C soil temperature is required for proper germination of pulses. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT:

As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @ 60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal. Apply irrigation in field if mustard was sown before 15-20 days.

VEGETABLES:

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied.
- Forecasted max and min temp is congenial for infestation of diamond back moth in cole crops. To control this apply Spinosad 45SC @ 0.5ml/l.
- There is a chance of occurrence of thrip, mites and Aphids in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.

Animal husbandry: In winter during cold hours farmer should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide sufficient clean drinking water to the animals.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)



Bulletin No.-604
Date- 11/11/2025

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR BADAMBA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 30°C (-1.25 from normal) and 18°C (-1.77 from normal), respectively.** The mean relative humidity was 72-79%.

Forecasted period: Today(8.30IST) to 16th November (8.30IST)2025

Weatheradvisory:: Dry weather is forecasted for upcoming 5days in the block. The daytemperature will be 30-31°C and night temperature will be 15-16°C for upcoming 5 days. Maximum and minimum relative humidity will be 75-81% and 37-39% respectively. **There will be clear sky for next 5days.**

Block	Badamba(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2025	12/11/2025	13/11/2025	14/11/2025	15/11/2025
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	30	30	31	30	30
T-Min(°C)	15	15	16	16	15
Cloud Cover(octa)	1	0	0	0	1
RH Max (%)	75	76	79	81	79
RH Min (%)	39	37	38	38	38
Wind Speed (Kmph)	6	7	7	6	5
Wind Direction (deg)	325	319	331	325	324

Agromet Advisory

General Advisory: **Harvest matured paddy crops and store them in safer place after sun drying. Complete nursery sowing or transplanting of winter vegetable crops. Irrigate the nursery during morning and evening time. Use low cost polythene cover above the vegetables nursery to avoid high evapotranspiration. Go for sowing of pea, gram and green gram crop by utilizing residual soil moisture**

Kharif Rice:

Paddy is grain formation to harvesting stage. **As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** To prevent crop from **Gundhi bug** apply 5 ml neem baesd insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. To control **rice ear head bug**, go for spraying of Dichlorovos @ 1ml/l during evening hours. Spray the chemical from borders to middle of the field. Present weather conditions is favourable for the incidence of **Neck blast** disease. To control, go for spraying of tricyclazole @ 0.6 g/l or Carbendazim @ 1 g / L. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis.

PULSES:

21-26°C soil temperature is required for proper germination of pulses. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT:

As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal. **Apply irrigation in field if mustard was sown before 15-20 days.**

VEGETABLES:

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. **In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied.**
- **Forecasted max and min temp is congenial for infestation of diamond back moth in cole crops.** To control this apply Spinosad 45SC @ 0.5ml/l.
- There is a chance of occurrence of **thrip, mites and Aphids** in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- **As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.**

Animal husbandry: In winter during cold hours farmer should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide sufficient clean drinking water to the animals.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-604
Date- 11/11/2025

AGROMET ADVISORY BULLETIN FOR BANKI

District Past week average weather condition and Block past week average Rainfall condition

The district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30°C (-1.25 from normal) and 18°C (-1.77 from normal), respectively. The mean relative humidity was 72-79%.

Forecasted period: Today(8.30IST) to 16th November (8.30IST)2025

Weather advisory:: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 30-31°C and night temperature will be 15-16°C for upcoming 5 days. Maximum and minimum relative humidity will be 67-73% and 30-34% respectively. **There will be clear sky for next 5 days.**

Block	Banki(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2025	12/11/2025	13/11/2025	14/11/2025	15/11/2025
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	30	30	31	30	30
T-Min(°C)	16	16	16	16	15
Cloud Cover(octa)	0	0	0	0	1
RH Max (%)	72	73	71	72	67
RH Min (%)	34	32	32	31	30
Wind Speed (Kmph)	7	7	7	7	6
Wind Direction (deg)	321	319	331	326	325

Agromet Advisory

General Advisory: Harvest matured paddy crops and store them in safer place after sun drying. Complete nursery sowing or transplanting of winter vegetable crops. Irrigate the nursery during morning and evening time. Use low cost polythene cover above the vegetables nursery to avoid high evapotranspiration. Go for sowing of pea, gram and green gram crop by utilizing residual soil moisture

Kharif Rice:

Paddy is grain formation to harvesting stage. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. To prevent crop from **Gundhi bug** apply 5 ml neem baesd insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. To control **rice ear head bug**, go for spraying of Dichlorovos @ 1ml/l during evening hours. Spray the chemical from borders to middle of the field. Present weather conditions is favourable for the incidence of **Neck blast** disease. To control, go for spraying of tricyclazole @ 0.6 g/l or Carbendazim @ 1 g / L. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis.

PULSES:

21-26°C soil temperature is required for proper germination of pulses. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT:

As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal. Apply irrigation in field if mustard was sown before 15-20 days.

VEGETABLES:

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied.
- Forecasted max and min temp is congenial for infestation of diamond back moth in cole crops. To control this apply Spinosad 45SC @ 0.5ml/l.
- There is a chance of occurrence of thrip, mites and Aphids in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.

Animal husbandry: In winter during cold hours farmer should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide sufficient clean drinking water to the animals.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area





GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-604

Date- 11/11/2025

AGROMET ADVISORY BULLETIN FOR BANKI-DAMPARA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30°C (-1.25 from normal) and 18°C (-1.77 from normal), respectively. The mean relative humidity was 72-79%.

Forecasted period: Today(8.30IST) to 16th November (8.30IST)2025

Weather Advisory:: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 30-31°C and night temperature will be 16°C for upcoming 5 days. Maximum and minimum relative humidity will be 67-73% and 30-34% respectively. There will be clear sky for next 5 days.

Block	Banki-Dampara (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2025	12/11/2025	13/11/2025	14/11/2025	15/11/2025
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	30	30	31	30	30
T-Min(°C)	16	16	16	16	16
Cloud Cover(octa)	0	0	0	0	1
RH Max (%)	72	73	71	72	67
RH Min (%)	34	32	32	31	30
Wind Speed (Kmph)	7	7	7	7	6
Wind Direction (deg)	321	319	331	326	325

Agromet Advisory

General Advisory: Harvest matured paddy crops and store them in safer place after sun drying. Complete nursery sowing or transplanting of winter vegetable crops. Irrigate the nursery during morning and evening time. Use low cost polythene cover above the vegetables nursery to avoid high evapotranspiration. Go for sowing of pea, gram and green gram crop by utilizing residual soil moisture

Kharif Rice:

Paddy is grain formation to harvesting stage. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. To prevent crop from Gundhi bug apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. To control rice ear head bug, go for spraying of Dichlorovos @ 1ml/l during evening hours. Spray the chemical from borders to middle of the field. Present weather conditions is favourable for the incidence of Neck blast disease. To control, go for spraying of tricyclazole @ 0.6 g/l or Carbendazim @ 1 g / L. If infestation of BPH & WBPH occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis.

PULSES:

21-26°C soil temperature is required for proper germination of pulses. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT:

As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal. Apply irrigation in field if mustard was sown before 15-20 days.

VEGETABLES:

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied.
- Forecasted max and min temp is congenial for infestation of diamond back moth in cole crops. To control this apply Spinosad 45SC @ 0.5ml/l.
- There is a chance of occurrence of thrip, mites and Aphids in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.

Animal husbandry: In winter during cold hours farmer should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide sufficient clean drinking water to the animals.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-604
Date- 11/11/2025

AGROMET ADVISORY BULLETIN FOR BARANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 30°C (-1.25 from normal) and 18°C (-1.77 from normal), respectively.** The mean relative humidity was 72-79%.

Forecasted period: Today(8.30IST) to 16th November (8.30IST)2025

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 30-31°C and night temperature will be 16°C for upcoming 5 days. Maximum and minimum relative humidity will be 74-77% and 37-40% respectively. **There will be clear sky for next 5 days.**

Block	Baranga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2025	12/11/2025	13/11/2025	14/11/2025	15/11/2025
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	30	30	31	30	30
T-Min(°C)	16	16	16	16	16
Cloud Cover	1	1	0	0	1
RH Max (%)	74	74	77	77	74
RH Min (%)	40	39	38	39	37
Wind Speed (Kmph)	7	8	8	6	6
Wind Direction (deg)	319	321	335	338	335

Agromet Advisory

General Advisory: **Harvest matured paddy crops and store them in safer place after sun drying. Complete nursery sowing or transplanting of winter vegetable crops. Irrigate the nursery during morning and evening time. Use low cost polythene cover above the vegetables nursery to avoid high evapotranspiration. Go for sowing of pea, gram and green gram crop by utilizing residual soil moisture**

Kharif Rice:

Paddy is grain formation to harvesting stage. **As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. To control **rice ear head bug**, go for spraying of Dichlorovos @ 1ml/l during evening hours. Spray the chemical from borders to middle of the field. Present weather conditions is favourable for the incidence of **Neck blast** disease. To control, go for spraying of triacyclazole @ 0.6 g/l or Carbendazim @ 1 g / L. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis.

PULSES:

21-26°C soil temperature is required for proper germination of pulses. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT:

As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @ 60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal. **Apply irrigation in field if mustard was sown before 15-20 days.**

VEGETABLES:

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. **In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied.**
- **Forecasted max and min temp is congenial for infestation of diamond back moth in cole crops.** To control this apply Spinosad 45SC @ 0.5ml/l.
- There is a chance of occurrence of **thrip, mites and Aphids** in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- **As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.**

Animal husbandry: In winter during cold hours farmer should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide sufficient clean drinking water to the animals.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area





GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)



Bulletin No.-604
Date- 11/11/2025

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR CUTTACK SADAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 30°C (-1.25 from normal) and 18°C (-1.77 from normal), respectively.** The mean relative humidity was 72-79%.

Forecasted period: Today(8.30IST) to 16th November (8.30IST)2025

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 29-30°C and night temperature will be 16-17°C for upcoming 5 days. Maximum and minimum relative humidity will be 63-74% and 31-35% respectively. **There will be clear sky for next 5 days.**

Block	Cuttack sadar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2025	12/11/2025	13/11/2025	14/11/2025	15/11/2025
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	30	30	30	30	29
T-Min(°C)	16	16	17	17	16
Cloud Cover(octa)	1	0	0	0	1
RH Max (%)	74	70	68	67	63
RH Min (%)	35	32	31	32	31
Wind Speed (Kmph)	7	7	8	7	7
Wind Direction (deg)	323	331	338	342	336

Agromet Advisory

General Advisory: Harvest matured paddy crops and store them in safer place after sun drying. Complete nursery sowing or transplanting of winter vegetable crops. Irrigate the nursery during morning and evening time. Use low cost polythene cover above the vegetables nursery to avoid high evapotranspiration. Go for sowing of pea, gram and green gram crop by utilizing residual soil moisture

Kharif Rice:

Paddy is grain formation to harvesting stage. **As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. To control **rice ear head bug**, go for spraying of Dichlorovos @ 1ml/l during evening hours. Spray the chemical from borders to middle of the field. Present weather conditions is favourable for the incidence of **Neck blast** disease. To control, go for spraying of tricyclazole @ 0.6 g/l or Carbendazim @ 1 g / L. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis.

PULSES: **21-26°C soil temperature is required for proper germination of pulses.** This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

Sugarcane:

The crop is at maturity (40 - 41 week) stage. Postpone irrigation to crops and spraying insecticides and pesticides. For control of Sucking pest in sugarcane like **Woolly aphid, Whitefly, Scale insects, mealybug** in present weather conditions go for alternate spraying of thiomethoxam 25WG @ 0.25 gm/lit or imidacloprid 0.45 ml/lit of water or Dinotefuron 20% SG 3gm/lit of Water.

GROUNDNUT: **As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done** if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @ 60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal. **Apply irrigation in field if mustard was sown before 15-20 days.**

VEGETABLES:

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. **In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied.**
- **Forecasted max and min temp is congenial for infestation of diamond back moth in cole crops.** To control this apply Spinosad 45SC @ 0.5ml/l.
- There is a chance of occurrence of **thrip, mites and Aphids** in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- **As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.**

Animal husbandry: In winter during cold hours farmer should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide sufficient clean drinking water to the animals.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-604
Date- 11/11/2025

AGROMET ADVISORY BULLETIN FOR KANTAPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30°C (-1.25 from normal) and 18°C (-1.77 from normal), respectively. The mean relative humidity was 72-79%.

Forecasted period: Today(8.30IST) to 16th November (8.30IST)2025

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 30-31°C and night temperature will be 17-18°C for upcoming 5 days. Maximum and minimum relative humidity will be 65-76% and 29-31% respectively. There will be clear sky for next 5 days.

Block	Kantapada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2025	12/11/2025	13/11/2025	14/11/2025	15/11/2025
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	30	30	31	31	30
T-Min(°C)	17	17	18	18	17
Cloud Cover(octa)	1	0	0	0	1
RH Max (%)	76	73	70	69	65
RH Min (%)	31	30	29	29	29
Wind Speed (Kmph)	9	9	9	8	8
Wind Direction (deg)	328	337	342	342	338

Agromet Advisory

General Advisory: Harvest matured paddy crops and store them in safer place after sun drying. Complete nursery sowing or transplanting of winter vegetable crops. Irrigate the nursery during morning and evening time. Use low cost polythene cover above the vegetables nursery to avoid high evapotranspiration. Go for sowing of pea, gram and green gram crop by utilizing residual soil moisture

Kharif Rice:

Paddy is grain formation to harvesting stage. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. To control **rice ear head bug**, go for spraying of Dichlorovos @ 1ml/l during evening hours. Spray the chemical from borders to middle of the field. Present weather conditions is favourable for the incidence of **Neck blast** disease. To control, go for spraying of tricyclazole @ 0.6 g/l or Carbendazim @ 1 g / L. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis.

PULSES: 21-26°C soil temperature is required for proper germination of pulses. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

Sugarcane:

The crop is at maturity (40 - 41 week) stage. Postpone irrigation to crops and spraying insecticides and pesticides. For control of Sucking pest in sugarcane like **Woolly aphid, Whitefly, Scale insects, mealybug** in present weather conditions go for alternate spraying of thiomethoxam 25WG @ 0.25 gm/lit or imidacloprid 0.45 ml/lit of water or Dinotefuron 20% SG 3gm/lit of Water.

GROUNDNUT: As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @ 60:30:30 N-P₂O₅-K₂O Kg/ha Mustard @ 80:40:40 N-P₂O₅-K₂O Kg/ha. Apply 50%N and all P₂O₅ and K₂O fertilizers should be applied as basal. Apply irrigation in field if mustard was sown before 15-20 days.

VEGETABLES:

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied.
- Forecasted max and min temp is congenial for infestation of diamond back moth in cole crops. To control this apply Spinosad 45SC @ 0.5ml/l.
- There is a chance of occurrence of **thrip, mites and Aphids** in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.

Animal husbandry: In winter during cold hours farmer should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide sufficient clean drinking water to the animals.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)



Bulletin No.-604
Date- 11/11/2025

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR MAHANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30°C (-1.25 from normal) and 18°C (-1.77 from normal), respectively. The mean relative humidity was 72-79%.

Forecasted period: Today(8.30IST) to 16th November (8.30IST)2025

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be and it will be 30-31°C and night temperature will be 18°C for upcoming 5 days. Maximum and minimum relative humidity will be 63-74% and 29-32% respectively. There will be clear sky for next 5 days.

Block	Mahanga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2025	12/11/2025	13/11/2025	14/11/2025	15/11/2025
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	30	30	31	30	30
T-Min(°C)	18	18	18	18	18
Cloud Cover(octal)	1	0	0	0	1
RH Max (%)	74	70	67	67	63
RH Min (%)	32	30	29	30	30
Wind Speed (Kmph)	7	8	8	7	7
Wind Direction (deg)	326	338	355	354	348

Agromet Advisory

General Advisory: Harvest matured paddy crops and store them in safer place after sun drying. Complete nursery sowing or transplanting of winter vegetable crops. Irrigate the nursery during morning and evening time. Use low cost polythene cover above the vegetables nursery to avoid high evapotranspiration. Go for sowing of pea, gram and green gram crop by utilizing residual soil moisture

Kharif Rice:

Paddy is grain formation to harvesting stage. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. To prevent crop from **Gundhi bug** apply 5 ml neem baesd insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. To control **rice ear head bug**, go for spraying of Dichlorovos @ 1ml/l during evening hours. Spray the chemical from borders to middle of the field. Present weather conditions is favourable for the incidence of **Neck blast** disease. To control, go for spraying of tricyclazole @ 0.6 g/l or Carbendazim @ 1 g / L. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis.

PULSES: 21-26°C soil temperature is required for proper germination of pulses. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

Sugarcane:

The crop is at maturity (40 - 41 week) stage. Postpone irrigation to crops and spraying insecticides and pesticides. For control of Sucking pest in sugarcane like **Woolly aphid, Whitefly, Scale insects, mealybug** in present weather conditions go for alternate spraying of thiomethoxam 25WG @ 0.25 gm/lit or imidacloprid 0.45 ml/lit of water or Dinotefuron 20% SG 3gm/lit of Water.

GROUNDNUT: As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @ 60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal. Apply irrigation in field if mustard was sown before 15-20 days.

VEGETABLES:

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied.
- Forecasted max and min temp is congenial for infestation of diamond back moth in cole crops. To control this apply Spinosad 45SC @ 0.5ml/l.
- There is a chance of occurrence of **thrip, mites and Aphids** in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.

Animal husbandry: In winter during cold hours farmer should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide sufficient clean drinking water to the animals.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area





GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-604
Date- 11/11/2025

AGROMET ADVISORY BULLETIN FOR NARASINGHPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 30°C (-1.25 from normal) and 18°C (-1.77 from normal), respectively.** The mean relative humidity was 72-79%.

Forecasted period: Today(8.30IST) to 16th November (8.30IST)2025

Weatheradvisory:: Dry weather is forecasted for upcoming 5days in the block. The day temperature will be 29-30°C and night temperature will be 15°C for upcoming 5 days. Maximum and minimum relative humidity will be 71-73% and 30-33% respectively. **There will be clear sky for next 5days.**

Block	Narasinghpur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2025	12/11/2025	13/11/2025	14/11/2025	15/11/2025
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	30	30	30	30	29
T-Min(°C)	15	15	15	15	15
Cloud Cover(/octa)	0	0	0	0	1
RH Max (%)	73	71	72	73	72
RH Min (%)	33	32	31	31	30
Wind Speed (Kmph)	6	6	6	6	5
Wind Direction (deg)	330	317	330	330	326

Agromet Advisory

General Advisory: Harvest matured paddy crops and store them in safer place after sun drying. Complete nursery sowing or transplanting of winter vegetable crops. Irrigate the nursery during morning and evening time. Use low cost polythene cover above the vegetables nursery to avoid high evapotranspiration. Go for sowing of pea, gram and green gram crop by utilizing residual soil moisture

Kharif Rice:

Paddy is grain formation to harvesting stage. **As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** To prevent crop from **Gundhi bug** apply 5 ml neem baesd insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. To control **rice ear head bug**, go for spraying of Dichlorovos @ 1ml/l during evening hours. Spray the chemical from borders to middle of the field. Present weather conditions is favourable for the incidence of **Neck blast** disease. To control, go for spraying of tricyclazole @ 0.6 g/l or Carbendazim @ 1 g / L. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis.

PULSES:

21-26°C soil temperature is required for proper germination of pulses. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT:

As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal. **Apply irrigation in field if mustard was sown before 15-20 days.**

VEGETABLES:

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. **In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied.**
- **Forecasted max and min temp is congenial for infestation of diamond back moth in cole crops.** To control this apply Spinosad 45SC @ 0.5ml/l.
- There is a chance of occurrence of **thrip, mites and Aphids** in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- **As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.**

Animal husbandry: In winter during cold hours farmer should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide sufficient clean drinking water to the animals.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area





GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-604
Date- 11/11/2025

AGROMET ADVISORY BULLETIN FOR NIALI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 30°C (-1.25 from normal) and 18°C (-1.77 from normal), respectively.** The mean relative humidity was 72-79%.

Forecasted period: Today(8.30IST) to 16th November (8.30IST)2025

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 30-31°C and night temperature will be 18-19°C for upcoming 5 days. Maximum and minimum relative humidity will be 78-83% and 33-35% respectively. **There will be clear sky for next 5 days.**

Block	Niali (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2025	12/11/2025	13/11/2025	14/11/2025	15/11/2025
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	30	30	31	31	30
T-Min(°C)	18	18	19	19	18
Cloud Cover(octa)	1	1	0	0	1
RH Max (%)	83	80	79	79	78
RH Min (%)	34	33	34	35	34
Wind Speed(Kmph)	7	9	9	8	8
Wind Direction(deg)	328	328	346	352	347

Agromet Advisory

General Advisory: **Harvest matured paddy crops and store them in safer place after sun drying. Complete nursery sowing or transplanting of winter vegetable crops. Irrigate the nursery during morning and evening time. Use low cost polythene cover above the vegetables nursery to avoid high evapotranspiration. Go for sowing of pea, gram and green gram crop by utilizing residual soil moisture**

Kharif Rice:

Paddy is grain formation to harvesting stage. **As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. To control **rice ear head bug**, go for spraying of Dichlorovos @ 1ml/l during evening hours. Spray the chemical from borders to middle of the field. Present weather conditions is favourable for the incidence of **Neck blast** disease. To control, go for spraying of tricyclazole @ 0.6 g/l or Carbendazim @ 1 g / L. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis.

PULSES: **21-26°C soil temperature is required for proper germination of pulses.** This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

Sugarcane:

The crop is at maturity (40 - 41 week) stage. Postpone irrigation to crops and spraying insecticides and pesticides. For control of Sucking pest in sugarcane like **Wooly aphid, Whitefly, Scale insects, mealybug** in present weather conditions go for alternate spraying of thiomethoxam 25WG @ 0.25 gm/lit or imidacloprid 0.45 ml/lit of water or Dinotefuron 20% SG 3gm/lit of Water.

GROUNDNUT: **As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done** if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @ 60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal. **Apply irrigation in field if mustard was sown before 15-20 days.**

VEGETABLES:

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. **In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied.**
- **Forecasted max and min temp is congenial for infestation of diamond back moth in cole crops.** To control this apply Spinosad 45SC @ 0.5ml/l.
- There is a chance of occurrence of **thrip, mites and Aphids** in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- **As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.**

Animal husbandry: In winter during cold hours farmer should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide sufficient clean drinking water to the animals.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area





GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)



Bulletin No.-604
Date- 11/11/2025

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR NISCHINTAKOILI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 30°C (-1.25 from normal) and 18°C (-1.77 from normal), respectively.** The mean relative humidity was 72-79%.

Forecasted period: Today(8.30IST) to 16th November (8.30IST)2025

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 30-31°C and night temperature will be 18°C for upcoming 5 days. Maximum and minimum relative humidity will be 63-75% and 30-32% respectively. **There will be clear sky for next 5 days.**

Block	Nischintakoili (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2025	12/11/2025	13/11/2025	14/11/2025	15/11/2025
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	30	30	31	30	30
T-Min(°C)	18	18	18	18	18
Cloud Cover(octa)	1	0	0	0	1
RH Max (%)	75	70	67	67	63
RH Min (%)	32	30	30	30	31
Wind Speed (Kmph)	7	8	8	7	7
Wind Direction (deg)	328	339	355	354	352

Agromet Advisory

General Advisory: **Harvest matured paddy crops and store them in safer place after sun drying. Complete nursery sowing or transplanting of winter vegetable crops. Irrigate the nursery during morning and evening time. Use low cost polythene cover above the vegetables nursery to avoid high evapotranspiration. Go for sowing of pea, gram and green gram crop by utilizing residual soil moisture**

Kharif Rice:

Paddy is grain formation to harvesting stage. **As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. To control **rice ear head bug**, go for spraying of Dichlorovos @ 1ml/l during evening hours. Spray the chemical from borders to middle of the field. Present weather conditions is favourable for the incidence of **Neck blast** disease. To control, go for spraying of tricyclazole @ 0.6 g/l or Carbendazim @ 1 g / L. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis.

PULSES: **21-26°C soil temperature is required for proper germination of pulses.** This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

Sugarcane:

The crop is at maturity (40 - 41 week) stage. Postpone irrigation to crops and spraying insecticides and pesticides. For control of Sucking pest in sugarcane like **Woolly aphid, Whitefly, Scale insects, mealybug** in present weather conditions go for alternate spraying of thiomethoxam 25WG @ 0.25 gm/lit or imidacloprid 0.45 ml/lit of water or Dinotefuron 20% SG 3gm/lit of Water.

GROUNDNUT: **As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done** if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @ 60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal. **Apply irrigation in field if mustard was sown before 15-20 days.**

VEGETABLES:

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. **In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied.**
- **Forecasted max and min temp is congenial for infestation of diamond back moth in cole crops.** To control this apply Spinosad 45SC @ 0.5ml/l.
- There is a chance of occurrence of **thrip, mites and Aphids** in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- **As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.**

Animal husbandry: In winter during cold hours farmer should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide sufficient clean drinking water to the animals.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-604
Date- 11/11/2025

AGROMET ADVISORY BULLETIN FOR SALEPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 30°C (-1.25 from normal) and 18°C (-1.77 from normal), respectively.** The mean relative humidity was 72-79%.

Forecasted period: Today(8.30IST) to 16th November (8.30IST)2025

Weather advisory: *Dry weather is forecasted for upcoming 5days in the block.* The day temperature will be 29-30°C and night temperature will be 16-17°C for upcoming 5 days. Maximum and minimum relative humidity will be 63-75% and 30-32% respectively. **There will be clear sky for next 5days.**

Block	Salepur(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2025	12/11/2025	13/11/2025	14/11/2025	15/11/2025
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	30	30	30	30	29
T-Min(°C)	17	17	17	17	16
Cloud Cover(octa)	1	0	0	0	1
RH Max (%)	75	69	67	67	63
RH Min (%)	32	31	30	30	31
Wind Speed (Kmph)	7	8	8	7	7
Wind Direction (deg)	328	338	355	354	348

Agromet Advisory

General Advisory: *Harvest matured paddy crops and store them in safer place after sun drying. Complete nursery sowing or transplanting of winter vegetable crops. Irrigate the nursery during morning and evening time. Use low cost polythene cover above the vegetables nursery to avoid high evapotranspiration. Go for sowing of pea, gram and green gram crop by utilizing residual soil moisture*

Kharif Rice:

Paddy is grain formation to harvesting stage. **As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. To control **rice ear head bug**, go for spraying of Dichlorovos @ 1ml/l during evening hours. Spray the chemical from borders to middle of the field. Present weather conditions is favourable for the incidence of **Neck blast** disease. To control, go for spraying of tricyclazole @ 0.6 g/l or Carbendazim @ 1 g / L. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis.

PULSES: **21-26°C soil temperature is required for proper germination of pulses.** This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

Sugarcane:

The crop is at maturity (40 - 41 week) stage. Postpone irrigation to crops and spraying insecticides and pesticides. For control of Sucking pest in sugarcane like **Wooly aphid, Whitefly, Scale insects, mealybug** in present weather conditions go for alternate spraying of thiomethoxam 25WG @ 0.25 gm/lit or imidacloprid 0.45 ml/lit of water or Dinotefuron 20% SG 3gm/lit of Water.

GROUNDNUT: **As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done** if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal. **Apply irrigation in field if mustard was sown before 15-20 days.**

VEGETABLES:

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. **In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied.**
- **Forecasted max and min temp is congenial for infestation of diamond back moth in cole crops.** To control this apply Spinosad 45SC @ 0.5ml/l.
- There is a chance of occurrence of **thrip, mites and Aphids** in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- **As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.**

Animal husbandry: In winter during cold hours farmer should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide sufficient clean drinking water to the animals.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR TANGI-CHOUDWAR



Bulletin No.-604

Date- 11/11/2025

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30°C (-1.25 from normal) and 18°C (-1.77 from normal), respectively. The mean relative humidity was 72-79%.

Forecasted period: Today(8.30IST) to 16th November (8.30IST)2025

Weatheradvisory: Dry weather is forecasted for upcoming 5days in the block. The day temperature will be 29-30°C and night temperature will be 16-17°C for upcoming 5 days. Maximum and minimum relative humidity will be 62-75% and 37-39% respectively. There will be clear sky for next 5days.

Block	Tangi-Choudwar(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2025	12/11/2025	13/11/2025	14/11/2025	15/11/2025
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	30	30	30	30	29
T-Min(°C)	16	16	17	17	16
Cloud Cover(octa)	1	1	0	0	1
RH Max (%)	75	70	69	66	62
RH Min (%)	38	37	38	39	39
Wind Speed (Kmph)	11	11	12	13	13
Wind Direction (deg)	334	334	350	351	351

Agromet Advisory

General Advisory: Harvest matured paddy crops and store them in safer place after sun drying. Complete nursery sowing or transplanting of winter vegetable crops. Irrigate the nursery during morning and evening time. Use low cost polythene cover above the vegetables nursery to avoid high evapotranspiration. Go for sowing of pea, gram and green gram crop by utilizing residual soil moisture

Kharif Rice:

Paddy is grain formation to harvesting stage. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. To prevent crop from Gundhi bug apply 5 ml neem baesd insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. To control rice ear head bug, go for spraying of Dichlorovos @ 1ml/l during evening hours. Spray the chemical from borders to middle of the field. Present weather conditions is favourable for the incidence of Neck blast disease. To control, go for spraying of tricyclazole @ 0.6 g/l or Carbendazim @ 1 g / L. If infestation of BPH & WBPH occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis.

PULSES:

21-26°C soil temperature is required for proper germination of pulses. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT:

As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal. Apply irrigation in field if mustard was sown before 15-20 days.

VEGETABLES:

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied.
- Forecasted max and min temp is congenial for infestation of diamond back moth in cole crops. To control this apply Spinosad 45SC @ 0.5ml/l.
- There is a chance of occurrence of thrip, mites and Aphids in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.

Animal husbandry: In winter during cold hours farmer should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide sufficient clean drinking water to the animals.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-604
Date- 11/11/2025

AGROMET ADVISORY BULLETIN FOR TIGIRIA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30°C (-1.25 from normal) and 18°C (-1.77 from normal), respectively. The mean relative humidity was 72-79%.

Forecasted period: Today(8.30IST) to 16th November (8.30IST)2025

Weatheradvisory:: Dry weather is forecasted for upcoming 5days in the block. The day temperature will be 30-31°C and night temperature will be 15°C for upcoming 5 days. Maximum and minimum relative humidity will be 67-73% and 31-37% respectively. **There will be clear sky for next 5days.**

Block	Tigiria(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2025	12/11/2025	13/11/2025	14/11/2025	15/11/2025
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	30	30	31	30	30
T-Min(°C)	15	15	15	15	15
Cloud Cover(octa)	0	0	0	0	1
RH Max (%)	73	71	71	72	67
RH Min (%)	37	33	33	32	31
Wind Speed (Kmph)	6	7	7	6	5
Wind Direction (deg)	325	321	331	327	326

Agro met Advisory

General Advisory: Harvest matured paddy crops and store them in safer place after sun drying. Complete nursery sowing or transplanting of winter vegetable crops. Irrigate the nursery during morning and evening time. Use low cost polythene cover above the vegetables nursery to avoid high evapotranspiration. Go for sowing of pea, gram and green gram crop by utilizing residual soil moisture

Kharif Rice:

Paddy is grain formation to harvesting stage. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. To prevent crop from Gundhi bug apply 5 ml neem baesd insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. To control rice ear head bug, go for spraying of Dichlorovos @ 1ml/l during evening hours. Spray the chemical from borders to middle of the field. Present weather conditions is favourable for the incidence of Neck blast disease. To control, go for spraying of tricyclazole @ 0.6 g/l or Carbendazim @ 1 g / L. If infestation of BPH & WBPH occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis.

PULSES:

21-26°C soil temperature is required for proper germination of pulses. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT:

As dry weather is forecasted for upcoming 5 days, harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal. Apply irrigation in field if mustard was sown before 15-20 days.

VEGETABLES:

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied.
- Forecasted max and min temp is congenial for infestation of diamond back moth in cole crops. To control this apply Spinosad 45SC @ 0.5ml/l.
- There is a chance of occurrence of thrip, mites and Aphids in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.

Animal husbandry: In winter during cold hours farmer should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide sufficient clean drinking water to the animals.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area








GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-604
Date- 11/11/2025

				
MEGHDOOT APP	DAMINI APP	FACEBOOK (Damu Cuttack)	BLOCK SPECIFIC WHAT'S APP GROUPS	@DAMUCUTTACK

- Kindly install and use **“Meghdoot mobile App”** for Cuttack district weather forecast, warnings and weather based agromet advisory services.

<https://play.google.com/store/apps/details?id=com.aas.meghdoot>

- Kindly install and use **“Damini mobile App”** for location specific for lightening forecast and warnings.

<https://play.google.com/store/apps/details?id=com.lightening.live.damini>

- Follow the **block specific whats app groups** and join in your block wise groups for getting weather based agromet advisory services twice in a week (Tuesday and Friday) and other weather warning based farm management advisories.
- Follow the **Official Website links** to get the weather based agromet advisory service bulletins (**English & Odia**) twice in a week (Tuesday and Friday).

1	ICAR-NRRI, Cuttack, Odisha	https://icar-nrri.in/agromate/	
2	Agrimet division, IMD, Pune	https://imdagrmet.gov.in/imdproject/AGIndex.php	



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

