




Cuttack District Block Wise Agro-met Advisory Service Bulletin



Tuesday, 06th January, 2026
(For the period 06th to 10th January, 2026)



Issued by
District Agromet Unit,
KrishiVigyan Kendra Cuttack



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kykcuttack@gmail.com/damukykcuttack@gmail.com



Bulletin No.-611
Date- 06/01/2026

STANDARD PRECIPITATION INDEX AND NDVI MAP

| Standard Precipitation Index (SPI) | Normalized difference vegetation Index (NDVI) |
|---|---|
| <p>MINISTRY OF EARTH SCIENCES INDIA METEOROLOGICAL DEPARTMENT HYDROMET SECTION, CAUI, CRS PUNE</p> <p>STANDARDISED PRECIPITATION INDEX (SPI) RECENT 4 WEEK 04 DECEMBER 2025 to 31 DECEMBER 2025</p> | <p>NOAA/VIIRS/BLENDED NDVI Composite ending on Week no 49 (03.12.2025 to 10.12.2025) over Agricultural region</p> <p>Odisha</p> <p>Agriculture vigour is good at isolated places of the state.</p> |
| <p><i>As per SPI value (cumulative 4 weeks -04 – 31 December 2025), mildly dry condition was there in Cuttack district.</i></p> | <p><i>According to NDVI map (03.12.2025 to 10.12.2025) the agriculture vigor was light to moderate over most blocks of Cuttack district.</i></p> |

| LEGEND (Medium range block level rainfall forecast) | | |
|--|------------------------------|-----------|
| | No Rainfall | 00 mm |
| | Very light rainfall | 01-05 mm |
| | Light rainfall | 06-10 mm |
| | Light to moderate rainfall | 11-30 mm |
| | Moderate to heavy rainfall | 31-65 mm |
| | Heavy to very heavy rainfall | 66-115 mm |
| | Extremely heavy rainfall | >116 mm |



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kykcuttack@gmail.com/damukykcuttack@gmail.com



Bulletin No.-611
Date- 06/01/2026

Weather (Past and Forecasted) based Disease and pest in Major Crops and vegetables

| Sl No | Crop | Stage and condition of the crop | Anticipated Disease and pest | |
|-------|-------------------------------------|---------------------------------|------------------------------|--------------------------|
| 1 | Pulses (Green gram and black gram) | Vegetative stage | | Aphid |
| | | | | Black headed caterpillar |
| | | | | Powdery mildew |
| | | | | YMV |
| 2 | Mustard | Vegetative stage | | Aphid |
| | | | | Saw fly |
| | | | | Leaf webber |
| | | | | Stink Bug |
| | | | | White rust |
| 3 | Rabi Groundnut | Vegetative stage | | Leaf Miner |
| 4 | Vegetables (Tomato,Chilli, Brinjal) | Vegetative to fruiting | | Thrips |
| | | | | Mites |
| | | | | Chili Aphid |
| 5 | Brinjal | Fruiting | | Fruit and shoot borer |
| 6 | Tomato | Vegetative | | White fly |



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kykcuttack@gmail.com/damukykcuttack@gmail.com



Bulletin No.-611
Date- 06/01/2026

| | | | | |
|---|--------------------------------------|----------------------------|--|-------------------------|
| | | | | Leaf Curl virus |
| | | Fruiting | | Late blight |
| 7 | Potato | Sprouting/Vegetative stage | | Leaf Blight |
| 8 | Cole crops (Cauliflower and cabbage) | Vegetative to fruiting | | Leaf eating caterpillar |
| | | | | Aphid |
| | | | | Fruit borer |
| | | | | Diamond back moth |
| | | | | Tobacco Caterpillar |



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)
Email: kykcuttack@gmail.com/damukykcuttack@gmail.com



Bulletin No.-611
Date- 06/01/2026

AGROMET ADVISORY BULLETIN FOR ATHAGAD

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 23.85°C (-4.06 from normal) and 12°C (-1.45 from normal), respectively. The mean relative humidity was 80-95%.

Forecasted period: Today(8.30IST) to 11thJanuary (8.30IST)2026

Weather advisory: *Dry weather is forecasted for upcoming 5 days in the block.* The day temperature will be 25-28°C and night temperature will be 12-13°C for upcoming 5 days. Maximum and minimum relative humidity will be 58-79% and 20-30% respectively. There will be clear to partly cloudy sky for next 5 days.

| Block | Athagad (VALID TILL 08:30 IST OF THE NEXT 5 DAYS) | | | | |
|----------------------|---|------------|------------|------------|------------|
| Date | 06/01/2026 | 07/01/2026 | 08/01/2026 | 09/01/2026 | 10/01/2026 |
| Rainfall (mm) | 00 | 00 | 00 | 00 | 00 |
| T-Max (°C) | 25 | 26 | 26 | 27 | 28 |
| T-Min (°C) | 12 | 12 | 12 | 12 | 13 |
| Cloud Cover (octa) | 0 | 2 | 1 | 2 | 2 |
| RH Max (%) | 79 | 68 | 71 | 63 | 58 |
| RH Min (%) | 30 | 24 | 23 | 23 | 20 |
| Wind Speed (kmph) | 8 | 5 | 5 | 5 | 6 |
| Wind Direction (deg) | 50 | 348 | 331 | 344 | 329 |

Agromet Advisory

General Advisory: Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Due to slight decrease in minimum temperature in upcoming 3-4 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress.

Pulses: In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Foggy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew. Present weather condition was favourable for the incidence of yellow mosaic virus which was caused by the vector white fly. To control go for spraying of triazophos @ 2ml/lt or Acetamiprid@0.2gm/lt.

Rabi-Groundnut:

To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

Mustard:

In view of dry weather condition during next five days; farmers are advised to irrigate the mustard crop at 30-40 DAS (4-6 leaf stage). Due to foggy weather in past , there are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids and stink bug at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf webber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/liter of water.

Vegetables:

- Use low cost polythene cover above the vegetables nursery to avoid poor germination due to low temperature.
- Transplanting of mature seedlings of tomato, cauliflower, cabbage, knolkhola and broccoli crop may be done in the raised bed.

Potato:

After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyriphos @ 2.5 ml/lit of water against the attack of cut worm and leaf eating insect. Due to past foggy weather there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

- Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water
- To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.
- Current weather is favourable for attack of Cabbage webber and tobacco caterpillar in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. Leaf eating caterpillar and fruit borer may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb14.5% SC mixing in 200lit of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l.

Animal husbandry: In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-611
Date- 06/01/2026

AGROMET ADVISORY BULLETIN FOR BADAMBA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 23.85°C (-4.06 from normal) and 12°C (-1.45 from normal), respectively. The mean relative humidity was 80-95%.

Forecasted period: Today(8.30IST) to 11thJanuary (8.30IST)2026

Weather advisory: *Dry weather is forecasted for upcoming 5 days in the block.* The day temperature will be 25-27°C and night temperature will be 11-12°C for upcoming 5 days. Maximum and minimum relative humidity will be 69-90% and 22-37% respectively. There will be clear to partly cloudy sky for next 5 days.

| Block | Badamba(VALID TILL 08:30 IST OF THE NEXT 5 DAYS) | | | | |
|----------------------|--|------------|------------|------------|------------|
| Date | 06/01/2026 | 07/01/2026 | 08/01/2026 | 09/01/2026 | 10/01/2026 |
| Rainfall(mm) | 00 | 00 | 00 | 00 | 00 |
| T-Max(°C) | 25 | 26 | 26 | 26 | 27 |
| T-Min(°C) | 12 | 12 | 12 | 11 | 11 |
| Cloud Cover(octa) | 1 | 1 | 1 | 1 | 3 |
| RH Max (%) | 90 | 72 | 72 | 69 | 70 |
| RH Min (%) | 37 | 28 | 26 | 22 | 24 |
| Wind Speed (Kmph) | 11 | 6 | 4 | 5 | 7 |
| Wind Direction (deg) | 53 | 337 | 340 | 333 | 326 |

Agromet Advisory

General Advisory: Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Due to slight decrease in minimum temperature in upcoming 3-4 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress.

Pulses: In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Foggy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew. Present weather condition was favourable for the incidence of yellow mosaic virus which was caused by the vector white fly. To control go for spraying of triazophos @ 2ml/lt or Acetamipride@0.2gm/lt.

Rabi-Groundnut:

To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

Mustard:

In view of dry weather condition during next five days; farmers are advised to irrigate the mustard crop at 30-40 DAS (4-6 leaf stage). Due to foggy weather in past , there are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids and stink bug at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf webberspray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/liter of water.

Vegetables:

- Use low cost polythene cover above the vegetables nursery to avoid poor germination due to low temperature.
- Transplanting of mature seedlings of tomato, cauliflower, cabbage, knolkhol and broccoli crop may be done in the raised bed.

Potato:

After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyriphos @ 2.5 ml/lt of water against the attack of cut worm and leaf eating insect. Due to past foggy weather there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

- Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water
- To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.
- Current weather is favourable for attack of Cabbage webber and tobacco caterpillar in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. Leaf eating caterpillar and fruit borer may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb14.5% SC mixing in 200lit of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l.

Animal husbandry: In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kykcuttack@gmail.com/damukyvkcuttack@gmail.com



Bulletin No.-611
Date- 06/01/2026

AGROMET ADVISORY BULLETIN FOR BANKI

District Past week average weather condition

The district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 23.85°C (-4.06 from normal) and 12°C (-1.45 from normal), respectively. The mean relative humidity was 80-95%.

Forecasted period: Today(8.30IST) to 11thJanuary (8.30IST)2026

Weather advisory: *Dry weather is forecasted for upcoming 5 days in the block.* The day temperature will be 25-28°C and night temperature will be 12-13°C for upcoming 5 days. Maximum and minimum relative humidity will be 61-83% and 20-31% respectively. There will be clear to partly cloudy sky for next 5 days.

| Block | Banki(VALID TILL 08:30 IST OF THE NEXT 5 DAYS) | | | | |
|----------------------|--|------------|------------|------------|------------|
| | 06/01/2026 | 07/01/2026 | 08/01/2026 | 09/01/2026 | 10/01/2026 |
| Rainfall(mm) | 00 | 00 | 00 | 00 | 00 |
| T-Max(°C) | 25 | 26 | 27 | 27 | 28 |
| T-Min(°C) | 12 | 12 | 12 | 12 | 13 |
| Cloud Cover(octa) | 1 | 2 | 1 | 2 | 2 |
| RH Max (%) | 83 | 67 | 72 | 64 | 61 |
| RH Min (%) | 31 | 23 | 22 | 22 | 20 |
| Wind Speed (Kmph) | 8 | 5 | 4 | 6 | 7 |
| Wind Direction (deg) | 49 | 344 | 336 | 345 | 328 |

Agromet Advisory

General Advisory: Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Due to slight decrease in minimum temperature in upcoming 3-4 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress.

Pulses: In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Foggy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew. Present weather condition was favourable for the incidence of yellow mosaic virus which was caused by the vector white fly. To control go for spraying of triazophos @ 2ml/ltr or Acetamiprid@0.2gm/ltr.

Rabi-Groundnut:

To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

Mustard:

In view of dry weather condition during next five days; farmers are advised to irrigate the mustard crop at 30-40 DAS (4-6 leaf stage). Due to foggy weather in past , there are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids and stink bug at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf webberspray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/liter of water.

Vegetables:

- Use low cost polythene cover above the vegetables nursery to avoid poor germination due to low temperature.
- Transplanting of mature seedlings of tomato, cauliflower, cabbage, knolkhola and broccoli crop may be done in the raised bed.

Potato:

After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyriphos @ 2.5 ml/ltr of water against the attack of cut worm and leaf eating insect. Due to past foggy weather there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

- **Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water
- To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.
- Current weather is favourable for attack of Cabbage webber and tobacco caterpillar in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. Leaf eating caterpillar and fruit borer may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb14.5% SC mixing in 200lit of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l.
- **Animal husbandry:** In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)
Email: kykcuttack@gmail.com/damukykcuttack@gmail.com



Bulletin No.-611
Date- 06/01/2026

AGROMET ADVISORY BULLETIN FOR BANKI-DAMPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 23.85°C (-4.06 from normal) and 12°C (-1.45 from normal), respectively. The mean relative humidity was 80-95%.

Forecasted period: Today(8.30IST) to 11thJanuary (8.30IST)2026

Weather Advisory: *Dry weather is forecasted for upcoming 5 days in the block.* The day temperature will be 25-28°C and night temperature will be 12-13°C for upcoming 5 days. Maximum and minimum relative humidity will be 61-83% and 20-31% respectively. There will be clear to partly cloudy sky for next 5 days.

| Block | Banki-Dampada(VALID TILL 08:30 IST OF THE NEXT 5 DAYS) | | | | |
|----------------------|--|------------|------------|------------|------------|
| | 06/01/2026 | 07/01/2026 | 08/01/2026 | 09/01/2026 | 10/01/2026 |
| Rainfall(mm) | 00 | 00 | 00 | 00 | 00 |
| T-Max(°C) | 25 | 26 | 27 | 27 | 28 |
| T-Min(°C) | 12 | 12 | 12 | 12 | 13 |
| Cloud Cover(octa) | 1 | 2 | 1 | 2 | 2 |
| RH Max (%) | 83 | 67 | 72 | 64 | 61 |
| RH Min (%) | 31 | 23 | 22 | 22 | 20 |
| Wind Speed (Kmph) | 8 | 5 | 4 | 6 | 7 |
| Wind Direction (deg) | 49 | 344 | 336 | 345 | 328 |

Agromet Advisory

General Advisory: Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Due to slight decrease in minimum temperature in upcoming 3-4 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress.

Pulses: In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Foggy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew. Present weather condition was favourable for the incidence of yellow mosaic virus which was caused by the vector white fly. To control go for spraying of triazophos @ 2ml/lt or Acetamipride@0.2gm/lt.

Rabi-Groundnut:

To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

Mustard:

In view of dry weather condition during next five days; farmers are advised to irrigate the mustard crop at 30-40 DAS (4-6 leaf stage). Due to foggy weather in past , there are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids and stink bug at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf webberspray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/liter of water.

Vegetables:

- Use low cost polythene cover above the vegetables nursery to avoid poor germination due to low temperature.
- Transplanting of mature seedlings of tomato, cauliflower, cabbage, knolkhola and broccoli crop may be done in the raised bed.

Potato:

After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyriphos @ 2.5 ml/lt of water against the attack of cut worm and leaf eating insect. Due to past foggy weather there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

- Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water
- To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.
- Current weather is favourable for attack of Cabbage webber and tobacco caterpillar in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. Leaf eating caterpillar and fruit borer may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb14.5% SC mixing in 200lit of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l.

Animal husbandry: In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kykcuttack@gmail.com/damukyvkcuttack@gmail.com



Bulletin No.-611
Date- 06/01/2026

AGROMET ADVISORY BULLETIN FOR BARANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 23.85°C (-4.06 from normal) and 12°C (-1.45 from normal), respectively. The mean relative humidity was 80-95%.

Forecasted period: Today(8.30IST) to 11thJanuary (8.30IST)2026

Weather advisory: *Dry weather is forecasted for upcoming 5 days in the block.* The day temperature will be 24-26°C and night temperature will be 13-14°C for upcoming 5 days. Maximum and minimum relative humidity will be 65-81% and 26-37% respectively. There will be clear to partly cloudy sky for next 5 days.

| Block | Baranga(VALID TILL 08:30 IST OF THE NEXT 5 DAYS) | | | | |
|----------------------|--|------------|------------|------------|------------|
| Date | 06/01/2026 | 07/01/2026 | 08/01/2026 | 09/01/2026 | 10/01/2026 |
| Rainfall(mm) | 00 | 00 | 00 | 00 | 00 |
| T-Max(°C) | 24 | 25 | 26 | 26 | 26 |
| T-Min(°C) | 14 | 13 | 13 | 14 | 14 |
| Cloud Cover | 1 | 1 | 1 | 1 | 3 |
| RH Max (%) | 81 | 72 | 69 | 68 | 65 |
| RH Min (%) | 37 | 31 | 28 | 27 | 26 |
| Wind Speed (Kmph) | 15 | 8 | 7 | 7 | 8 |
| Wind Direction (deg) | 51 | 347 | 342 | 348 | 331 |

Agromet Advisory

General Advisory: Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Due to slight decrease in minimum temperature in upcoming 3-4 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress.

Pulses: In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Foggy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew. Present weather condition was favourable for the incidence of yellow mosaic virus which was caused by the vector white fly. To control go for spraying of triazophos @ 2ml/lt or Acetamiprid@0.2gm/lt.

Rabi-Groundnut:

To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

Mustard:

In view of dry weather condition during next five days; farmers are advised to irrigate the mustard crop at 30-40 DAS (4-6 leaf stage). Due to foggy weather in past , there are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids and stink bug at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf webber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/liter of water.

Vegetables:

- Use low cost polythene cover above the vegetables nursery to avoid poor germination due to low temperature.
- Transplanting of mature seedlings of tomato, cauliflower, cabbage, knolkhola and broccoli crop may be done in the raised bed.

Potato:

After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyriphos @ 2.5 ml/lt of water against the attack of cut worm and leaf eating insect. Due to past foggy weather there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

- Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water
- To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.
- Current weather is favourable for attack of Cabbage webber and tobacco caterpillar in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. Leaf eating caterpillar and fruit borer may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb14.5% SC mixing in 200lit of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l.
- **Animal husbandry:** In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-611
Date- 06/01/2026

AGROMET ADVISORY BULLETIN FOR CUTTACK SADAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 23.85°C (-4.06 from normal) and 12°C (-1.45 from normal), respectively. The mean relative humidity was 80-95%.

Forecasted period: Today(8.30IST) to 11thJanuary (8.30IST)2026

Weather advisory; Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 25-28°C and night temperature will be 13-14°C for upcoming 5 days. Maximum and minimum relative humidity will be 54-69% and 20-30% respectively. There will be clear to partly cloudy sky for next 5 days.

| Block | Cuttack sadar(VALID TILL 08:30 IST OF THE NEXT 5 DAYS) | | | | |
|----------------------|--|------------|------------|------------|------------|
| | 06/01/2026 | 07/01/2026 | 08/01/2026 | 09/01/2026 | 10/01/2026 |
| Rainfall(mm) | 00 | 00 | 00 | 00 | 00 |
| T-Max(°C) | 25 | 26 | 26 | 27 | 28 |
| T-Min(°C) | 13 | 13 | 13 | 13 | 14 |
| Cloud Cover(octa) | 0 | 2 | 1 | 2 | 2 |
| RH Max (%) | 69 | 65 | 66 | 63 | 54 |
| RH Min (%) | 30 | 24 | 23 | 22 | 20 |
| Wind Speed (Kmph) | 12 | 7 | 7 | 7 | 8 |
| Wind Direction (deg) | 49 | 360 | 354 | 360 | 355 |

Agromet Advisory

General Advisory: Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Due to slight decrease in minimum temperature in upcoming 3-4 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress.

Pulses: In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Foggy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew. Present weather condition was favourable for the incidence of yellow mosaic virus which was caused by the vector white fly. To control go for spraying of triazophos @ 2ml/lt or Acetamiprid@0.2gm/lt.

Rabi-Groundnut:

To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

Mustard:

In view of dry weather condition during next five days; farmers are advised to irrigate the mustard crop at 30-40 DAS (4-6 leaf stage). Due to foggy weather in past , there are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids and stink bug at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf webber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/liter of water.

Vegetables:

- Use low cost polythene cover above the vegetables nursery to avoid poor germination due to low temperature.
- Transplanting of mature seedlings of tomato, cauliflower, cabbage, knolkhola and broccoli crop may be done in the raised bed.

Potato:

After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyriphos @ 2.5 ml/lt of water against the attack of cut worm and leaf eating insect. Due to past foggy weather there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

- Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water
- To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.
- Current weather is favourable for attack of Cabbage webber and tobacco caterpillar in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. Leaf eating caterpillar and fruit borer may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb14.5% SC mixing in 200lit of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l.

Animal husbandry: In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightning alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-611
Date- 06/01/2026

AGROMET ADVISORY BULLETIN FOR KANTAPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 23.85°C (-4.06 from normal) and 12°C (-1.45 from normal), respectively. The mean relative humidity was 80-95%.

Forecasted period: Today(8.30IST) to 11thJanuary (8.30IST)2026

Weather advisory: *Dry weather is forecasted for upcoming 5 days in the block.* The day temperature will be 25-28°C and night temperature will be 12-14°C for upcoming 5 days. Maximum and minimum relative humidity will be 60-72% and 20-29% respectively. There will be clear to partly cloudy sky for next 5 days.

| Block | Kantapada(VALID TILL 08:30 IST OF THE NEXT 5 DAYS) | | | | | |
|----------------------|--|------------|------------|------------|------------|------------|
| | Date | 06/01/2026 | 07/01/2026 | 08/01/2026 | 09/01/2026 | 10/01/2026 |
| Rainfall(mm) | 00 | 00 | 00 | 00 | 00 | 00 |
| T-Max(°C) | 25 | 26 | 27 | 27 | 28 | |
| T-Min(°C) | 13 | 12 | 13 | 13 | 14 | |
| Cloud Cover(octa) | 0 | 2 | 2 | 2 | 3 | |
| RH Max (%) | 71 | 68 | 68 | 72 | 60 | |
| RH Min (%) | 29 | 23 | 22 | 23 | 20 | |
| Wind Speed (Kmph) | 14 | 9 | 8 | 9 | 9 | |
| Wind Direction (deg) | 46 | 5 | 355 | 5 | 356 | |

Agromet Advisory

General Advisory: Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Due to slight decrease in minimum temperature in upcoming 3-4 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress.

Pulses: In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Foggy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew. Present weather condition was favourable for the incidence of yellow mosaic virus which was caused by the vector white fly. To control go for spraying of triazophos @ 2ml/lt or Acetamipride@0.2gm/lt.

Rabi-Groundnut:

To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

Mustard:

In view of dry weather condition during next five days; farmers are advised to irrigate the mustard crop at 30-40 DAS (4-6 leaf stage). Due to foggy weather in past , there are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids and stink bug at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf webber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/liter of water.

Vegetables:

- Use low cost polythene cover above the vegetables nursery to avoid poor germination due to low temperature.
- Transplanting of mature seedlings of tomato, cauliflower, cabbage, knolkhol and broccoli crop may be done in the raised bed.

Potato:

After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyriphos @ 2.5 ml/lt of water against the attack of cut worm and leaf eating insect. Due to past foggy weather there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

- **Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water
- To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.
- Current weather is favourable for attack of Cabbage webber and tobacco caterpillar in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. Leaf eating caterpillar and fruit borer may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb14.5% SC mixing in 200lit of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l.

Animal husbandry: In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)
Email: kykcuttack@gmail.com/damukykcuttack@gmail.com



Bulletin No.-611
Date- 06/01/2026

AGROMET ADVISORY BULLETIN FOR MAHANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 23.85°C (-4.06 from normal) and 12°C (-1.45 from normal), respectively. The mean relative humidity was 80-95%.

Forecasted period: Today(8.30IST) to 11thJanuary (8.30IST)2026

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be and it will be 25-27°C and night temperature will be 12-14°C for upcoming 5 days. Maximum and minimum relative humidity will be 54-68% and 21-30% respectively. There will be clear to partly cloudy sky for next 5 days.

| Block | Mahanga(VALID TILL 08:30 IST OF THE NEXT 5 DAYS) | | | | | |
|----------------------|--|------------|------------|------------|------------|------------|
| | Date | 06/01/2026 | 07/01/2026 | 08/01/2026 | 09/01/2026 | 10/01/2026 |
| Rainfall(mm) | 00 | 00 | 00 | 00 | 00 | 00 |
| T-Max(°C) | 25 | 26 | 26 | 26 | 27 | |
| T-Min(°C) | 13 | 12 | 13 | 12 | 14 | |
| Cloud Cover(octal) | 0 | 2 | 2 | 2 | 3 | |
| RH Max (%) | 68 | 66 | 64 | 66 | 54 | |
| RH Min (%) | 30 | 24 | 23 | 23 | 21 | |
| Wind Speed (Kmph) | 13 | 8 | 8 | 7 | 9 | |
| Wind Direction (deg) | 39 | 13 | 360 | 6 | 360 | |

Agromet Advisory

General Advisory: Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Due to slight decrease in minimum temperature in upcoming 3-4 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress.

Pulses: In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Foggy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew. Present weather condition was favourable for the incidence of yellow mosaic virus which was caused by the vector white fly. To control go for spraying of triazophos @ 2ml/lt or Acetamipride@0.2gm/lt.

Rabi-Groundnut:

To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

Mustard:

In view of dry weather condition during next five days; farmers are advised to irrigate the mustard crop at 30-40 DAS (4-6 leaf stage). Due to foggy weather in past , there are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids and stink bug at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf webber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/liter of water.

Vegetables:

- Use low cost polythene cover above the vegetables nursery to avoid poor germination due to low temperature.
- Transplanting of mature seedlings of tomato, cauliflower, cabbage, knolkhol and broccoli crop may be done in the raised bed.

Potato:

After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyriphos @ 2.5 ml/lt of water against the attack of cut worm and leaf eating insect. Due to past foggy weather there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

- Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water
- To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.
- Current weather is favourable for attack of Cabbage webber and tobacco caterpillar in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. Leaf eating caterpillar and fruit borer may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb14.5% SC mixing in 200lit of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l.

Animal husbandry: In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kykcuttack@gmail.com/damukykcuttack@gmail.com



Bulletin No.-611
Date- 06/01/2026

AGROMET ADVISORY BULLETIN FOR NARASINGHPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 23.85°C (-4.06 from normal) and 12°C (-1.45 from normal), respectively. The mean relative humidity was 80-95%.

Forecasted period: Today(8.30IST) to 11thJanuary (8.30IST)2026

Weather advisory: *Dry weather is forecasted for upcoming 5 days in the block.* The day temperature will be 26-28°C and night temperature will be 10-12°C for upcoming 5 days. Maximum and minimum relative humidity will be 58-84% and 17-26% respectively. There will be clear to partly cloudy sky for next 5 days.

| Block | Narasinghpur(VALID TILL 08:30 IST OF THE NEXT 5 DAYS) | | | | |
|----------------------|---|------------|------------|------------|------------|
| Date | 06/01/2026 | 07/01/2026 | 08/01/2026 | 09/01/2026 | 10/01/2026 |
| Rainfall(mm) | 00 | 00 | 00 | 00 | 00 |
| T-Max(°C) | 26 | 27 | 27 | 27 | 28 |
| T-Min(°C) | 10 | 10 | 10 | 11 | 12 |
| Cloud Cover(/octa) | 1 | 1 | 1 | 3 | 3 |
| RH Max (%) | 84 | 66 | 65 | 58 | 58 |
| RH Min (%) | 26 | 20 | 19 | 17 | 19 |
| Wind Speed (Kmph) | 5 | 4 | 3 | 5 | 6 |
| Wind Direction (deg) | 34 | 325 | 333 | 331 | 327 |

Agromet Advisory

General Advisory: Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Due to slight decrease in minimum temperature in upcoming 3-4 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress.

Pulses: In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Foggy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew. Present weather condition was favourable for the incidence of yellow mosaic virus which was caused by the vector white fly. To control go for spraying of triazophos @ 2ml/lt or Acetamipride@0.2gm/lt.

Rabi-Groundnut:

To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

Mustard:

In view of dry weather condition during next five days; farmers are advised to irrigate the mustard crop at 30-40 DAS (4-6 leaf stage). Due to foggy weather in past , there are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids and stink bug at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf webberspray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/liter of water.

Vegetables:

- Use low cost polythene cover above the vegetables nursery to avoid poor germination due to low temperature.
- Transplanting of mature seedlings of tomato, cauliflower, cabbage, knolkhol and broccoli crop may be done in the raised bed.

Potato:

After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyriphos @ 2.5 ml/lt of water against the attack of cut worm and leaf eating insect. Due to past foggy weather there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

- Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water
- To control the sucking pests like aphid & white fly, spray Imidaclorpid @ 5ml/10 lit of water.
- Current weather is favourable for attack of Cabbage webber and tobacco caterpillar in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. Leaf eating caterpillar and fruit borer may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb14.5% SC mixing in 200lit of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l.

• **Animal husbandry:** In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-611
Date- 06/01/2026

AGROMET ADVISORY BULLETIN FOR NIALI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 23.85°C (-4.06 from normal) and 12°C (-1.45 from normal), respectively. The mean relative humidity was 80-95%.

Forecasted period: Today(8.30IST) to 11thJanuary (8.30IST)2026

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 25-26°C and night temperature will be 13-14°C for upcoming 5 days. Maximum and minimum relative humidity will be 69-80% and 26-35% respectively. There will be clear to partly cloudy sky for next 5 days.

| Block | Niali(VALID TILL 08:30 IST OF THE NEXT 5 DAYS) | | | | |
|---------------------|--|------------|------------|------------|------------|
| Date | 06/01/2026 | 07/01/2026 | 08/01/2026 | 09/01/2026 | 10/01/2026 |
| Rainfall(mm) | 00 | 00 | 00 | 00 | 00 |
| T-Max(°C) | 25 | 26 | 26 | 26 | 26 |
| T-Min(°C) | 14 | 13 | 13 | 13 | 13 |
| Cloud Cover(octa) | 0 | 1 | 1 | 1 | 3 |
| RH Max (%) | 80 | 72 | 70 | 77 | 69 |
| RH Min (%) | 35 | 29 | 26 | 27 | 26 |
| Wind Speed(Kmph) | 16 | 10 | 10 | 9 | 9 |
| Wind Direction(deg) | 41 | 352 | 349 | 349 | 344 |

Agromet Advisory

General Advisory: Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Due to slight decrease in minimum temperature in upcoming 3-4 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress.

Pulses: In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Foggy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew. Present weather condition was favourable for the incidence of yellow mosaic virus which was caused by the vector white fly. To control go for spraying of triazophos @ 2ml/lt or Acetamipride@0.2gm/lt.

Rabi-Groundnut:

To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

Mustard:

In view of dry weather condition during next five days; farmers are advised to irrigate the mustard crop at 30-40 DAS (4-6 leaf stage). Due to foggy weather in past, there are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids and stink bug at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf webber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/liter of water.

Vegetables:

- Use low cost polythene cover above the vegetables nursery to avoid poor germination due to low temperature.

- Transplanting of mature seedlings of tomato, cauliflower, cabbage, knolkhola and broccoli crop may be done in the raised bed.

Potato:

After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyriphos @ 2.5 ml/lt of water against the attack of cut worm and leaf eating insect. Due to past foggy weather there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

- Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water
- To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.
- Current weather is favourable for attack of Cabbage webber and tobacco caterpillar in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. Leaf eating caterpillar and fruit borer may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb14.5% SC mixing in 200lit of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l.

- **Animal husbandry:** In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kykcuttack@gmail.com/damukyvkcuttack@gmail.com



Bulletin No.-611
Date- 06/01/2026

AGROMET ADVISORY BULLETIN FOR NISCHINTAKOILI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 23.85°C (-4.06 from normal) and 12°C (-1.45 from normal), respectively. The mean relative humidity was 80-95%.

Forecasted period: Today(8.30IST) to 11thJanuary (8.30IST)2026

Weather advisory: *Dry weather is forecasted for upcoming 5 days in the block.* The day temperature will be 25-28°C and night temperature will be 12-14°C for upcoming 5 days. Maximum and minimum relative humidity will be 54-68% and 21-30% respectively. There will be clear to partly cloudy sky for next 5 days.

| Block | Nischintakoili(VALID TILL 08:30 IST OF THE NEXT 5 DAYS) | | | | |
|----------------------|---|------------|------------|------------|------------|
| Date | 06/01/2026 | 07/01/2026 | 08/01/2026 | 09/01/2026 | 10/01/2026 |
| Rainfall(mm) | 00 | 00 | 00 | 00 | 00 |
| T-Max(°C) | 25 | 26 | 26 | 26 | 28 |
| T-Min(°C) | 13 | 12 | 13 | 12 | 14 |
| Cloud Cover(octa) | 0 | 2 | 2 | 2 | 3 |
| RH Max (%) | 68 | 66 | 64 | 67 | 54 |
| RH Min (%) | 30 | 24 | 23 | 23 | 21 |
| Wind Speed (Kmph) | 13 | 8 | 8 | 8 | 9 |
| Wind Direction (deg) | 39 | 13 | 3 | 5 | 360 |

Agromet Advisory

General Advisory: Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Due to slight decrease in minimum temperature in upcoming 3-4 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress.

Pulses: In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Foggy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew. Present weather condition was favourable for the incidence of yellow mosaic virus which was caused by the vector white fly. To control go for spraying of triazophos @ 2ml/lt or Acetamiprid@0.2gm/lt.

Rabi-Groundnut:

To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

Mustard:

In view of dry weather condition during next five days; farmers are advised to irrigate the mustard crop at 30-40 DAS (4-6 leaf stage). Due to foggy weather in past , there are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids and stink bug at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf webberspray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/liter of water.

Vegetables:

- Use low cost polythene cover above the vegetables nursery to avoid poor germination due to low temperature.
- Transplanting of mature seedlings of tomato, cauliflower, cabbage, knolkhola and broccoli crop may be done in the raised bed.

Potato:

After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyriphos @ 2.5 ml/lt of water against the attack of cut worm and leaf eating insect. Due to past foggy weather there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

- **Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water
- To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.
- Current weather is favourable for attack of Cabbage webber and tobacco caterpillar in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. Leaf eating caterpillar and fruit borer may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb14.5% SC mixing in 200lit of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l.
- **Animal husbandry:** In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kykcuttack@gmail.com/damukykcuttack@gmail.com



Bulletin No.-611
Date- 06/01/2026

AGROMET ADVISORY BULLETIN FOR SALEPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 23.85°C (-4.06 from normal) and 12°C (-1.45 from normal), respectively. The mean relative humidity was 80-95%.

Forecasted period: Today(8.30IST) to 11thJanuary (8.30IST)2026

Weather advisory: *Dry weather is forecasted for upcoming 5 days in the block.* The day temperature will be 25-28°C and night temperature will be 13-14°C for upcoming 5 days. Maximum and minimum relative humidity will be 54-68% and 21-30% respectively. There will be clear to partly cloudy sky for next 5 days.

| Block | Salepur(VALID TILL 08:30 IST OF THE NEXT 5 DAYS) | | | | |
|----------------------|--|------------|------------|------------|------------|
| Date | 06/01/2026 | 07/01/2026 | 08/01/2026 | 09/01/2026 | 10/01/2026 |
| Rainfall(mm) | 00 | 00 | 00 | 00 | 00 |
| T-Max(°C) | 25 | 26 | 26 | 26 | 28 |
| T-Min(°C) | 13 | 13 | 13 | 13 | 14 |
| Cloud Cover(octa) | 0 | 2 | 2 | 2 | 3 |
| RH Max (%) | 68 | 66 | 64 | 67 | 54 |
| RH Min (%) | 30 | 24 | 23 | 23 | 21 |
| Wind Speed (Kmph) | 13 | 8 | 8 | 8 | 9 |
| Wind Direction (deg) | 41 | 13 | 360 | 5 | 360 |

Agromet Advisory

General Advisory: Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Due to slight decrease in minimum temperature in upcoming 3-4 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress.

Pulses: In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Foggy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew. Present weather condition was favourable for the incidence of yellow mosaic virus which was caused by the vector white fly. To control go for spraying of triazophos @ 2ml/lt or Acetamipride@0.2gm/lt.

Rabi-Groundnut:

To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

Mustard:

In view of dry weather condition during next five days; farmers are advised to irrigate the mustard crop at 30-40 DAS (4-6 leaf stage). Due to foggy weather in past , there are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids and stink bug at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf webber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/liter of water.

Vegetables:

- Use low cost polythene cover above the vegetables nursery to avoid poor germination due to low temperature.
- Transplanting of mature seedlings of tomato, cauliflower, cabbage, knolkhol and broccoli crop may be done in the raised bed.

Potato:

After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyriphos @ 2.5 ml/lt of water against the attack of cut worm and leaf eating insect. Due to past foggy weather there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

- Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water
- To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.
- Current weather is favourable for attack of Cabbage webber and tobacco caterpillar in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. Leaf eating caterpillar and fruit borer may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb14.5% SC mixing in 200lit of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l.

Animal husbandry: In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)
Email: kykcuttack@gmail.com/damukykcuttack@gmail.com



Bulletin No.-611
Date- 06/01/2026

AGROMET ADVISORY BULLETIN FOR TANGI-CHOWDWAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 23.85°C (-4.06 from normal) and 12°C (-1.45 from normal), respectively. The mean relative humidity was 80-95%.

Forecasted period: Today(8.30IST) to 11thJanuary (8.30IST)2026

Weather advisory: *Dry weather is forecasted for upcoming 5 days in the block.* The day temperature will be 24-26°C and night temperature will be 13-14°C for upcoming 5 days. Maximum and minimum relative humidity will be 63-68% and 29-38% respectively. There will be clear to partly cloudy sky for next 5 days.

| Block | Tangi-Chowdwar(VALID TILL 08:30 IST OF THE NEXT 5 DAYS) | | | | |
|----------------------|---|------------|------------|------------|------------|
| | 06/01/2026 | 07/01/2026 | 08/01/2026 | 09/01/2026 | 10/01/2026 |
| Rainfall(mm) | 00 | 00 | 00 | 00 | 00 |
| T-Max(°C) | 24 | 24 | 25 | 25 | 26 |
| T-Min(°C) | 13 | 13 | 13 | 13 | 14 |
| Cloud Cover(octa) | 0 | 1 | 2 | 2 | 3 |
| RH Max (%) | 68 | 65 | 64 | 64 | 63 |
| RH Min (%) | 38 | 34 | 33 | 29 | 29 |
| Wind Speed (Kmph) | 18 | 15 | 14 | 14 | 12 |
| Wind Direction (deg) | 21 | 4 | 5 | 1 | 356 |

Agromet Advisory

General Advisory: Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Due to slight decrease in minimum temperature in upcoming 3-4 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress.

Pulses: In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Foggy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew. Present weather condition was favourable for the incidence of yellow mosaic virus which was caused by the vector white fly. To control go for spraying of triazophos @ 2ml/lt or Acetamiprid@0.2gm/lt.

Rabi-Groundnut:

To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

Mustard:

In view of dry weather condition during next five days; farmers are advised to irrigate the mustard crop at 30-40 DAS (4-6 leaf stage). Due to foggy weather in past, there are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids and stink bug at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf webber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/liter of water.

Vegetables:

- Use low cost polythene cover above the vegetables nursery to avoid poor germination due to low temperature.
- Transplanting of mature seedlings of tomato, cauliflower, cabbage, knolkhol and broccoli crop may be done in the raised bed.

Potato:

After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyriphos @ 2.5 ml/lt of water against the attack of cut worm and leaf eating insect. Due to past foggy weather there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

- Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water
- To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.
- Current weather is favourable for attack of Cabbage webber and tobacco caterpillar in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. Leaf eating caterpillar and fruit borer may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb14.5% SC mixing in 200lit of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l.
- **Animal husbandry:** In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-611
Date- 06/01/2026

AGROMET ADVISORY BULLETIN FOR TIGIRIA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 23.85°C (-4.06 from normal) and 12°C (-1.45 from normal), respectively. The mean relative humidity was 80-95%.

Forecasted period: Today(8.30IST) to 11thJanuary (8.30IST)2026

Weather advisory: *Dry weather is forecasted for upcoming 5 days in the block.* The day temperature will be 25-28°C and night temperature will be 11-13°C for upcoming 5 days. Maximum and minimum relative humidity will be 61-84% and 20-30% respectively. There will be clear to partly cloudy sky for next 5 days.

| Block | Tigiria(VALID TILL 08:30 IST OF THE NEXT 5 DAYS) | | | | |
|----------------------|--|------------|------------|------------|------------|
| Date | 06/01/2026 | 07/01/2026 | 08/01/2026 | 09/01/2026 | 10/01/2026 |
| Rainfall(mm) | 00 | 00 | 00 | 00 | 00 |
| T-Max(°C) | 25 | 26 | 26 | 27 | 28 |
| T-Min(°C) | 12 | 11 | 11 | 12 | 13 |
| Cloud Cover(octa) | 1 | 2 | 1 | 2 | 3 |
| RH Max (%) | 84 | 67 | 72 | 64 | 61 |
| RH Min (%) | 30 | 23 | 22 | 22 | 20 |
| Wind Speed (Kmph) | 7 | 5 | 4 | 6 | 7 |
| Wind Direction (deg) | 47 | 337 | 326 | 337 | 326 |

Agro met Advisory

General Advisory: Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Due to slight decrease in minimum temperature in upcoming 3-4 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress.

Pulses: In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Foggy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew. Present weather condition was favourable for the incidence of yellow mosaic virus which was caused by the vector white fly. To control go for spraying of triazophos @ 2ml/ltr or Acetamiprid@0.2gm/ltr.

Rabi-Groundnut:

To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

Mustard:

In view of dry weather condition during next five days; farmers are advised to irrigate the mustard crop at 30-40 DAS (4-6 leaf stage). Due to foggy weather in past , there are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids and stink bug at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf webberspray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/liter of water.

Vegetables:

- Use low cost polythene cover above the vegetables nursery to avoid poor germination due to low temperature.
- Transplanting of mature seedlings of tomato, cauliflower, cabbage, knolkhola and broccoli crop may be done in the raised bed.

Potato:

After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyriphos @ 2.5 ml/ltr of water against the attack of cut worm and leaf eating insect. Due to past foggy weather there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

- **Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water
- To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.
- Current weather is favourable for attack of Cabbage webber and tobacco caterpillar in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. Leaf eating caterpillar and fruit borer may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb14.5% SC mixing in 200lit of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l.
- **Animal husbandry:** In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kykcuttack@gmail.com/damukykcuttack@gmail.com



Bulletin No.-611
Date- 06/01/2026

| | | | | |
|---------------------|-------------------|--------------------------------|---|---------------------|
| | | | | |
| MEGHDOOT APP | DAMINI APP | FACEBOOK (Damu Cuttack) | BLOCK SPECIFIC WHAT'S APP GROUPS | @DAMUCUTTACK |

- Kindly install and use “**Meghdoot mobile App**” for Cuttack district weather forecast, warnings and weather based agromet advisory services.

<https://play.google.com/store/apps/details?id=com.aas.meghdoott>

- Kindly install and use “**Damini mobile App**” for location specific for lightening forecast and warnings.

<https://play.google.com/store/apps/details?id=com.lightening.live.damini>

- Follow the **block specific whats app groups** and join in your block wise groups for getting weather based agromet advisory services twice in a week (Tuesday and Friday) and other weather warning based farm management advisories.

- Follow the **Official Website links** to get the weather based agromet advisory service bulletins (**English & Odia**) twice in a week (Tuesday and Friday).

| | | | |
|---|---------------------------------------|---|--|
| 1 | ICAR-NRRI, Cuttack, Odisha | https://icar-nrri.in/agromate/ | |
| 2 | Agrimet division, IMD, Pune | https://imdagrimet.gov.in/imdproject/AGIndex.php | |



*Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area*

**DrRanjan Kumar Mohanta
(Nodal officer)**

