




# Cuttack District Block Wise Agro-met Advisory Service Bulletin



**Tuesday, 20<sup>th</sup> January, 2026**  
**(For the period 20<sup>th</sup> to 24<sup>th</sup> January, 2026)**



**Issued by**  
**District Agromet Unit,**  
**KrishiVigyan Kendra Cuttack**



Follow Meghdoot Mobile APP for Agromet  
advisories in local language and DAMINI  
APP for Lightening alert in your area

DrRanjan Kumar Mohanta  
(Nodal officer)





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
**(A Unit of ICAR-National Rice Research Institute)**  
**Email: kykcuttack@gmail.com/damukykcuttack@gmail.com**



Bulletin No.-613  
Date- 20/01/2026

**STANDARD PRECIPITATION INDEX AND NDVI MAP**

Standard Precipitation Index (SPI)	Normalized difference vegetation Index (NDVI)
<p>MINISTRY OF EARTH SCIENCES  INDIA METEOROLOGICAL DEPARTMENT  HYDROMET SECTION, CAUI, CRS PUNE</p> <p>STANDARDISED PRECIPITATION INDEX (SPI)  RECENT 4 WEEK  04 DECEMBER 2025 to 31 DECEMBER 2025</p> <p><b>SPI Values</b></p> <ul style="list-style-type: none"> <li>No Data</li> <li>&lt; -2 (Extremely Dry)</li> <li>-1.5 to -1.99 (Severely Dry)</li> <li>-1 to -1.49 (Moderately Dry)</li> <li>0 to -0.99 (Mildly Dry)</li> <li>0 to 0.99 (Mildly Wet)</li> <li>1 to 1.49 (Moderately Wet)</li> <li>1.5 to 1.99 (Severely Wet)</li> <li>&gt; 2 (Extremely Wet)</li> </ul>	<p>NOAA/VIIRS/BLENDED NDVI Composite ending on Week no 01  (01.01.2026 to 07.01.2026) over Agricultural region</p> <p style="text-align: center;"><b>Odisha</b></p> <p><b>NDVI Legend</b></p> <ul style="list-style-type: none"> <li>-9999 - 0</li> <li>0 - 0.05</li> <li>0.05 - 0.1</li> <li>0.1 - 0.15</li> <li>0.15 - 0.2</li> <li>0.2 - 0.25</li> <li>0.25 - 0.3</li> <li>0.3 - 0.35</li> <li>0.35 - 0.4</li> <li>0.4 - 0.45</li> <li>0.45 - 0.5</li> <li>0.5 - 0.55</li> <li>0.55 - 0.6</li> <li>&gt; 0.6</li> </ul> <p>Agriculture vigour is good at isolated places of the state.</p>
<p><i>As per SPI value (cumulative 4weeks -04 – 31 December 2025), mildly dry condition was there in Cuttack district.</i></p>	<p><i>According to NDVI map (01.01.2026 to 07.01.2026) the agriculture vigor was very light to light over most blocks of Cuttack district.</i></p>

<b>LEGEND (Medium range block level rainfall forecast)</b>		
	No Rainfall	00 mm
	Very light rainfall	01-05 mm
	Light rainfall	06-10 mm
	Light to moderate rainfall	11-30 mm
	Moderate to heavy rainfall	31-65 mm
	Heavy to very heavy rainfall	66-115 mm
	Extremely heavy rainfall	>116 mm



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta  
(Nodal officer)





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
**(A Unit of ICAR-National Rice Research Institute)**  
**Email: kykcuttack@gmail.com/damukykcuttack@gmail.com**



Bulletin No.-613  
Date- 20/01/2026

**Weather (Past and Forecasted) based Disease and pest in Major Crops and vegetables**

Sl No	Crop	Stage and condition of the crop	Anticipated Disease and pest	
1	Pulses (Green gram and black gram)	Vegetative stage to flowering		Aphid
				White fly
				Powdery mildew
2	Mustard	Vegetative stage to flowering		Aphid
				Saw fly
				Leaf webber
				Stink Bug
3	Rabi Groundnut	Vegetative stage to flowering		Tikka disease
				Aphid and Mites
4	Vegetables (Tomato,Chilli, Brinjal)	Vegetative to fruiting		Thrips
				Mites
				Chili Aphid
6	Tomato	Flowering to Fruting		Leaf Curl virus
7	Potato	Sprouting/Vegetative stage		Leaf Blight
				Aphid



Follow Meghdoot Mobile APP for Agromet  
advisories in local language and DAMINI  
APP for Lightening alert in your area

DrRanjan Kumar Mohanta  
(Nodal officer)





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
**(A Unit of ICAR-National Rice Research Institute)**  
Email: [kvkcuttack@gmail.com](mailto:kvkcuttack@gmail.com)/[damukvkcuttack@gmail.com](mailto:damukvkcuttack@gmail.com)



Bulletin No.-613  
Date- 20/01/2026

8	<p><b>Cole crops (Cauliflower and cabbage)</b></p> <p><b>Vegetative to fruiting</b></p>		<b>Leaf eating caterpillar</b>
			<b>Aphid</b>
			<b>Fruit borer</b>
			<b>Diamond back moth</b>
			<b>Tobacco Caterpillar</b>



Follow Meghdoot Mobile APP for Agromet  
advisories in local language and DAMINI  
APP for Lightening alert in your area

**DrRanjan Kumar Mohanta**  
(Nodal officer)





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
**(A Unit of ICAR-National Rice Research Institute)**  
**Email: kykcuttack@gmail.com/damukykcuttack@gmail.com**



Bulletin No.-613  
Date- 20/01/2026

**AGROMET ADVISORY BULLETIN FOR ATHAGAD**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 26.14°C (-2.57 from normal) and 10.57°C (-4.4 from normal), respectively. The mean relative humidity was 65-77%.

**Forecasted period: Today(8.30IST) to 25<sup>th</sup>January (8.30IST)2026**

**Weather advisory:** *Dry weather is forecasted for upcoming 5 days in the block.* The day temperature will be 30-31°C and night temperature will be 13-14°C for upcoming 5 days. Maximum and minimum relative humidity will be 88-94% and 23-27% respectively. There will be clear sky for next 5 days.

Block	Athagad (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	20/01/2026	21/01/2026	22/01/2026	23/01/2026	24/01/2026
Rainfall (mm)	00	00	00	00	00
T-Max (°C)	31	31	30	30	31
T-Min (°C)	13	13	14	14	14
Cloud Cover (octa)	0	0	0	0	0
RH Max (%)	88	91	91	94	91
RH Min (%)	23	24	27	27	24
Wind Speed (kmph)	3	5	5	2	2
Wind Direction (deg)	297	333	315	333	321

**Agromet Advisory**

**GeneralAdvisory:** *As shallow to moderate fog has occurred over isolated places in past 4-5 days and also likely to occur in next 2-3 days, so farmers are advised to monitor potato, tomato and onion crops for incidence of early/late blight disease and mustard for white rust, green and black gram crop for powdery mildew. if symptoms are noticed, apply recommended plant protection measures.*

**Pulses:**

**Farmers should start sowing of pulses by utilization of irrigation water.**

Cloudy and foggy weather may aggravate the infestation of **white fly, aphid and powdery mildew** in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and **Whitefly** and spray 3gm Sulfex or 1ml Propiconazole 10% EC or 1 ml Tebuconazole 25.9%EC per litre water for controlling **Powdery mildew**.

**Rabi-Groundnut:**

**As early sown groundnut crop is at post flowering stage (After 30-40 days of sowing), so farmers are advised to irrigate their field after this stage due to mildly dry condition in Cuttack district in last one month and also in upcoming days.**

Farmers are advised to do intercultivation and apply **Gypsum @200 kg/ acre** in **Groundnut** which is at **flowering stage**. To manage **aphids** and **thrips** spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. To manage **tikka** disease in groundnut farmers should spray with 500g Blitox-50 powder in 200 litre water.

**Mustard: As dry weather is needed after sowing of mustard, so farmers should complete sowing of Mustard in between upcoming 5 days.**

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of **Aphid** and **stink bug** infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage **Mustard sawfly** and **leafWebber** spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

**Vegetables:**

**Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water. To control the **sucking pests like aphid & white fly**, spray Imidacloprid @ 5ml/10 lit of water.

**Cabbage and Cauliflower:** Current weather is favorable for attack of **Cabbage Webber** and **tobacco caterpillar** in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. **Leaf eating caterpillar and fruit borer** may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectin benzoate 5% SG 80gm or indoxacarb 14.5% SC mixing in 200lit. of water per acre crop. At three-week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by **diamond back moth**. To control this, apply Spinosad 45SC @ 0.5ml/l.

**Tomato:** Transplanted seedlings are at 6-8-week stage in main field. Complete hoeing, weeding and apply 27 kg urea at six-week stage. Irrigate the crops after fertilizer application. To control **tomato leaf curl virus** in field, Spray Imidachloprid 17.8% SL @ 0.5 ml or Dimethoate 30% EC @ 1 ml/litre of 10 water at 15, 25, 45 days after transplanting.

**Potato:** There are chances of Leaf Blight disease incidence in potato crop. To manage **early leaf blight** in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400-gram/acre at 10 days' interval.

**Onion:** The transplanted seedlings are at seedling establishment to four-week stage in main field. **Irrigate the crop at 10 days' interval.** Go for hoeing, weeding and 1<sup>st</sup> top dressing with 45kg urea per acre crop area at 4-week stage of crop followed by irrigation.

**Animal husbandry:** In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

**DrRanjan Kumar Mohanta**  
**(Nodal officer)**





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**

**(A Unit of ICAR-National Rice Research Institute)**  
**Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com**



Bulletin No.-613  
Date- 20/01/2026

**AGROMET ADVISORY BULLETIN FOR BADAMBA**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 26.14°C (-2.57 from normal) and 10.57°C (-4.4 from normal), respectively. The mean relative humidity was 65-77%.

**Forecasted period: Today(8.30IST) to 25<sup>th</sup>January (8.30IST)2026**

**Weather advisory:** *Dry weather is forecasted for upcoming 5 days in the block.* The day temperature will be 29-30°C and night temperature will be 14-15°C for upcoming 5 days. Maximum and minimum relative humidity will be 87-91% and 25-33% respectively. There will be clear sky for next 5 days.

Block	Badamba(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)					
	Date	20/01/2026	21/01/2026	22/01/2026	23/01/2026	24/01/2026
Rainfall(mm)	00	00	00	00	00	00
T-Max(°C)	30	30	30	29	30	30
T-Min(°C)	14	14	15	15	15	15
Cloud Cover(octa)	0	0	0	0	0	0
RH Max (%)	87	88	90	91	91	91
RH Min (%)	26	25	28	33	29	29
Wind Speed (Kmph)	2	5	5	2	2	2
Wind Direction (deg)	259	330	331	248	288	

**Agromet Advisory**

**GeneralAdvisory:** *As shallow to moderate fog has occurred over isolated places in past 4-5 days and also likely to occur in next 2-3 days, so farmers are advised to monitor potato, tomato and onion crops for incidence of early/late blight disease and mustard for white rust, green and black gram crop for powdery mildew. if symptoms are noticed, apply recommended plant protection measures.*

**Pulses:**

**Farmers should start sowing of pulses by utilization of irrigation water.**

Cloudy and foggy weather may aggravate the infestation of **white fly, aphid and powdery mildew** in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and **Whitefly** and spray 3gm Sulfex or 1ml Propiconazole 10% EC or 1 ml Tebuconazole 25.9%EC per litre water for controlling **Powdery mildew**.

**Rabi-Groundnut:**

**As early sown groundnut crop are at post flowering stage(After 30-40 days of sowing), so farmers are advised to irrigate their field after this stage due to mildly dry condition in Cuttack district in last one month and also in upcoming days.**

Farmers are advised to do intercultivation and apply **Gypsum @200 kg/ acre** in **Groundnut** which is at flowering stage. To manage **aphids** and **thrips** spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. To manage **tikka** disease in groundnut farmers should spray with 500g Blitox-50 powder in 200 litre water.

**Mustard:** *As dry weather is needed after sowing of mustard, so farmers should complete sowing of Mustard in between upcoming 5 days.*

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of **Aphid** and **stink bug** infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage **Mustard sawfly** and **leafWebber** spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

**Vegetables:**

**Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water. To control the **sucking pests like aphid & white fly**, spray Imidacloprid @ 5ml/10 lit of water.

**Cabbage and Cauliflower:** Current weather is favorable for attack of **Cabbage Webber** and **tobacco caterpillar** in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. **Leaf eating caterpillar and fruit borer** may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectin benzoate 5% SG 80gm or indoxacarb 14.5% SC mixing in 200lit. of water per acre crop. At three-week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by **diamond back moth**. To control this, apply Spinosad 45SC @ 0.5ml/l.

**Tomato:** Transplanted seedlings are at 6-8-week stage in main field. Complete hoeing, weeding and apply 27 kg urea at six-week stage. Irrigate the crops after fertilizer application. To control **tomato leaf curl virus** in field, Spray Imidachloprid 17.8% SL @ 0.5 ml or Dimethoate 30% EC @ 1 ml/litre of 10 water at 15, 25, 45 days after transplanting.

**Potato:** There are chances of Leaf Blight disease incidence in potato crop. To manage **early leaf blight** in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days' interval.

**Onion:** The transplanted seedlings are at seedling establishment to four-week stage in main field. **Irrigate the crop at 10 days' interval.** Go for hoeing, weeding and 1<sup>st</sup> top dressing with 45kg urea per acre crop area at 4-week stage of crop followed by irrigation.

**Animal husbandry:** In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta  
(Nodal officer)





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
**(A Unit of ICAR-National Rice Research Institute)**  
**Email: kykcuttack@gmail.com/damukykcuttack@gmail.com**



Bulletin No.-613  
Date- 20/01/2026

**AGROMET ADVISORY BULLETIN FOR BANKI**

**District Past week average weather condition**

The district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 26.14°C (-2.57 from normal) and 10.57°C (-4.4 from normal), respectively. The mean relative humidity was 65-77%.

**Forecasted period: Today(8.30IST) to 25<sup>th</sup>January (8.30IST)2026**

**Weather advisory:** *Dry weather is forecasted for upcoming 5 days in the block.* The day temperature will be 30-31°C and night temperature will be 13-15°C for upcoming 5 days. Maximum and minimum relative humidity will be 89-93% and 23-28% respectively. There will be clear sky for next 5days.

Block	Banki(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	20/01/2026	21/01/2026	22/01/2026	23/01/2026	24/01/2026
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	30	30	30	30	31
T-Min(°C)	13	13	14	15	14
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	89	91	92	93	92
RH Min (%)	23	24	27	28	24
Wind Speed (Kmph)	3	5	6	1	2
Wind Direction (deg)	297	326	320	270	315

**Agromet Advisory**

**GeneralAdvisory:** *As shallow to moderate fog has occurred over isolated places in past4-5 days and also likely to occur in next 2-3 days, so farmers are advised to monitor potato, tomato and onion crops for incidence of early/late blight disease and mustard for white rust, green and black gram crop for powdery mildew. if symptoms are noticed, apply recommended plant protection measures.*

**Pulses:**

**Farmers should start sowing of pulses by utilization of irrigation water.**

Cloudy and foggy weather may aggravate the infestation of white fly, aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and Whitefly and spray 3gm Sulfex or 1ml Propiconazole 10% EC or 1 ml Tebuconazole 25.9%EC per litre water for controlling Powdery mildew.

**Rabi-Groundnut:**

**As early sown groundnut crop are at post flowering stage(After 30-40 days of sowing), so farmers are advised to irrigate their field after this stage due to mildly dry condition in Cuttack district in last one month and also in upcoming days.**

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. To manage aphids and thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. To manage tikka disease in groundnut farmers should spray with 500g Blitox-50 powder in 200 litre water.

**Mustard:** *As dry weather is needed after sowing of mustard, so farmers should complete sowing of Mustard in between upcoming 5 days.*

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leafWebber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

**Vegetables:**

**Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water. To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.

**Cabbage and Cauliflower:** Current weather is favorable for attack of **Cabbage Webber** and **tobacco caterpillar** in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. **Leaf eating caterpillar and fruit borer** may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectin benzoate 5% SG 80gm or indoxacarb14.5% SC mixing in 200lit. of water per acre crop. At three-week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by **diamond back moth**. To control this, apply Spinosad 45SC @ 0.5ml/l.

**Tomato:** Transplanted seedlings are at 6-8-week stage in main field. Complete hoeing, weeding and apply 27 kg urea at six-week stage. Irrigate the crops after fertilizer application. To control **tomato leaf curl virus** in field, Spray Imidachloprid 17.8% SL @ 0.5 ml or Dimethoate 30% EC @ 1 ml/litre of 10 water at 15, 25, 45 days after transplanting.

**Potato:** There are chances of Leaf Blight disease incidence in potato crop. To manage **early leaf blight** in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days' interval.

**Onion:** The transplanted seedlings are at seedling establishment to four-week stage in main field. **Irrigate the crop at 10 days' interval.** Go for hoeing, weeding and 1<sup>st</sup> top dressing with 45kg urea per acre crop area at 4-week stage of crop followed by irrigation.

**Animal husbandry:** In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta  
(Nodal officer)





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**

**(A Unit of ICAR-National Rice Research Institute)**  
**Email: kykcuttack@gmail.com/damukykcuttack@gmail.com**



Bulletin No.-613  
Date- 20/01/2026

**AGROMET ADVISORY BULLETIN FOR BANKI-DAMPADA**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 26.14°C (-2.57 from normal) and 10.57°C (-4.4 from normal), respectively. The mean relative humidity was 65-77%.

**Forecasted period: Today(8.30IST) to 25<sup>th</sup>January (8.30IST)2026**

**Weather Advisory:** *Dry weather is forecasted for upcoming 5 days in the block.* The day temperature will be 30-31°C and night temperature will be 13-15°C for upcoming 5 days. Maximum and minimum relative humidity will be 89-93% and 23-28% respectively. There will be clear sky for next 5 days.

Block	Banki-Dampada(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	20/01/2026	21/01/2026	22/01/2026	23/01/2026	24/01/2026
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	30	30	30	30	31
T-Min(°C)	13	13	14	15	14
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	89	91	92	93	92
RH Min (%)	23	24	27	28	24
Wind Speed (Kmph)	3	5	6	1	2
Wind Direction (deg)	297	326	320	270	315

**Agromet Advisory**

**General Advisory:** *As shallow to moderate fog has occurred over isolated places in past 4-5 days and also likely to occur in next 2-3 days, so farmers are advised to monitor potato, tomato and onion crops for incidence of early/late blight disease and mustard for white rust, green and black gram crop for powdery mildew. If symptoms are noticed, apply recommended plant protection measures.*

**Pulses:**

**Farmers should start sowing of pulses by utilization of irrigation water.**

Cloudy and foggy weather may aggravate the infestation of **white fly, aphid and powdery mildew** in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and **Whitefly** and spray 3gm Sulfex or 1ml Propiconazole 10% EC or 1 ml Tebuconazole 25.9%EC per litre water for controlling **Powdery mildew**.

**Rabi-Groundnut:**

**As early sown groundnut crop are at post flowering stage(After 30-40 days of sowing), so farmers are advised to irrigate their field after this stage due to mildly dry condition in Cuttack district in last one month and also in upcoming days.**

Farmers are advised to do intercultivation and apply **Gypsum @200 kg/ acre** in Groundnut which is at flowering stage. To manage **aphids and thrips** spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. To manage **tikka** disease in groundnut farmers should spray with 500g Blitox-50 powder in 200 litre water.

**Mustard:** *As dry weather is needed after sowing of mustard, so farmers should complete sowing of Mustard in between upcoming 5 days.*

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of **Aphid and stink bug** infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage **Mustard sawfly and leafWebber** spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

**Vegetables:**

**Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water. To control the **sucking pests like aphid & white fly**, spray Imidacloprid @ 5ml/10 lit of water.

**Cabbage and Cauliflower:** Current weather is favorable for attack of **Cabbage Webber** and **tobacco caterpillar** in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. **Leaf eating caterpillar and fruit borer** may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectin benzoate 5% SG 80gm or indoxacarb 14.5% SC mixing in 200lit. of water per acre crop. At three-week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by **diamond back moth**. To control this, apply Spinosad 45SC @ 0.5ml/l.

**Tomato:** Transplanted seedlings are at 6-8-week stage in main field. Complete hoeing, weeding and apply 27 kg urea at six-week stage. Irrigate the crops after fertilizer application. To control **tomato leaf curl virus** in field, Spray Imidachloprid 17.8% SL @ 0.5 ml or Dimethoate 30% EC @ 1 ml/litre of 10 water at 15, 25, 45 days after transplanting.

**Potato:** There are chances of Leaf Blight disease incidence in potato crop. To manage **early leaf blight** in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days' interval.

**Onion:** The transplanted seedlings are at seedling establishment to four-week stage in main field. **Irrigate the crop at 10 days' interval.** Go for hoeing, weeding and 1<sup>st</sup> top dressing with 45kg urea per acre crop area at 4-week stage of crop followed by irrigation.

**Animal husbandry:** In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightning alert in your area

DrRanjan Kumar Mohanta  
(Nodal officer)





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
**(A Unit of ICAR-National Rice Research Institute)**  
**Email: kykcuttack@gmail.com/damukyvkcuttack@gmail.com**



Bulletin No.-613  
Date- 20/01/2026

**AGROMET ADVISORY BULLETIN FOR BARANGA**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 26.14°C (-2.57 from normal) and 10.57°C (-4.4 from normal), respectively. The mean relative humidity was 65-77%.

**Forecasted period: Today(8.30IST) to 25<sup>th</sup>January (8.30IST)2026**

**Weather advisory:** *Dry weather is forecasted for upcoming 5 days in the block.* The day temperature will be 29-30°C and night temperature will be 16-17°C for upcoming 5 days. Maximum and minimum relative humidity will be 85-93% and 29-35% respectively. There will be clear sky for next 5 days.

Block	Baranga(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	20/01/2026	21/01/2026	22/01/2026	23/01/2026	24/01/2026
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	30	30	29	29	30
T-Min(°C)	16	16	17	17	17
Cloud Cover	0	0	0	0	0
RH Max (%)	85	88	90	93	92
RH Min (%)	30	29	33	35	32
Wind Speed (Kmph)	3	7	7	3	4
Wind Direction (deg)	276	335	317	306	336

**Agromet Advisory**

**GeneralAdvisory:** *As shallow to moderate fog has occurred over isolated places in past 4-5 days and also likely to occur in next 2-3 days, so farmers are advised to monitor potato, tomato and onion crops for incidence of early/late blight disease and mustard for white rust, green and black gram crop for powdery mildew. If symptoms are noticed, apply recommended plant protection measures.*

**Pulses:**

**Farmers should start sowing of pulses by utilization of irrigation water.**

Cloudy and foggy weather may aggravate the infestation of white fly, aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and Whitefly and spray 3gm Sulfex or 1ml Propiconazole 10% EC or 1 ml Tebuconazole 25.9%EC per litre water for controlling Powdery mildew.

**Rabi-Groundnut:**

**As early sown groundnut crop are at post flowering stage(After 30-40 days of sowing), so farmers are advised to irrigate their field after this stage due to mildly dry condition in Cuttack district in last one month and also in upcoming days.**

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. To manage aphids and thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. To manage tikka disease in groundnut farmers should spray with 500g Blitox-50 powder in 200 litre water.

**Mustard:** *As dry weather is needed after sowing of mustard, so farmers should complete sowing of Mustard in between upcoming 5 days.*

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leafWebber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

**Vegetables:**

**Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water. To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.

**Cabbage and Cauliflower:** Current weather is favorable for attack of **Cabbage Webber** and **tobacco caterpillar** in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. **Leaf eating caterpillar and fruit borer** may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectin benzoate 5% SG 80gm or indoxacarb 14.5% SC mixing in 200lit. of water per acre crop. At three-week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by **diamond back moth**. To control this, apply Spinosad 45SC @ 0.5ml/l.

**Tomato:** Transplanted seedlings are at 6-8-week stage in main field. Complete hoeing, weeding and apply 27 kg urea at six-week stage. Irrigate the crops after fertilizer application. To control **tomato leaf curl virus** in field, Spray Imidachloprid 17.8% SL @ 0.5 ml or Dimethoate 30% EC @ 1 ml/litre of 10 water at 15, 25, 45 days after transplanting.

**Potato:** There are chances of Leaf Blight disease incidence in potato crop. To manage **early leaf blight** in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days' interval.

**Onion:** The transplanted seedlings are at seedling establishment to four-week stage in main field. **Irrigate the crop at 10 days' interval.** Go for hoeing, weeding and 1<sup>st</sup> top dressing with 45kg urea per acre crop area at 4-week stage of crop followed by irrigation.

• **Animal husbandry:** In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta  
(Nodal officer)





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
**(A Unit of ICAR-National Rice Research Institute)**  
**Email: kykcuttack@gmail.com/damukyvkcuttack@gmail.com**



Bulletin No.-613  
Date- 20/01/2026

**AGROMET ADVISORY BULLETIN FOR CUTTACK SADAR**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 26.14°C (-2.57 from normal) and 10.57°C (-4.4 from normal), respectively. The mean relative humidity was 65-77%.

**Forecasted period: Today(8.30IST) to 25<sup>th</sup>January (8.30IST)2026**

**Weatheradvisory:** *Dry weather is forecasted for upcoming 5 days in the block.* The day temperature will be 30-31°C and night temperature will be 14-15°C for upcoming 5 days. Maximum and minimum relative humidity will be 85-93% and 25-28% respectively. There will be clear sky for next 5 days.

Block	Cuttack sadar(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	20/01/2026	21/01/2026	22/01/2026	23/01/2026	24/01/2026
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	30	30	30	30	31
T-Min(°C)	14	15	15	15	15
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	85	89	91	93	87
RH Min (%)	25	25	28	28	25
Wind Speed (Kmph)	5	6	7	3	4
Wind Direction (deg)	303	335	313	328	319

**Agromet Advisory**

**GeneralAdvisory:** *As shallow to moderate fog has occurred over isolated places in past 4-5 days and also likely to occur in next 2-3 days, so farmers are advised to monitor potato, tomato and onion crops for incidence of early/late blight disease and mustard for white rust, green and black gram crop for powdery mildew. if symptoms are noticed, apply recommended plant protection measures.*

**Pulses:**

**Farmers should start sowing of pulses by utilization of irrigation water.**

Cloudy and foggy weather may aggravate the infestation of **white fly, aphid and powdery mildew** in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and **Whitefly** and spray 3gm Sulfex or 1ml Propiconazole 10% EC or 1 ml Tebuconazole 25.9%EC per litre water for controlling **Powdery mildew**.

**Rabi-Groundnut:**

**As early sown groundnut crop are at post flowering stage(After 30-40 days of sowing), so farmers are advised to irrigate their field after this stage due to mildly dry condition in Cuttack district in last one month and also in upcoming days.**

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. To manage **aphids** and **thrips** spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. To manage **tikka** disease in groundnut farmers should spray with 500g Blitox-50 powder in 200 litre water.

**Mustard:** *As dry weather is needed after sowing of mustard, so farmers should complete sowing of Mustard in between upcoming 5 days.*

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of **Aphid** and **stink bug** infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage **Mustard sawfly** and **leafWebber** spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

**Vegetables:**

**Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water. To control the **sucking pests like aphid & white fly**, spray Imidacloprid @ 5ml/10 lit of water.

**Cabbage and Cauliflower:** Current weather is favorable for attack of **Cabbage Webber** and **tobacco caterpillar** in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. **Leaf eating caterpillar and fruit borer** may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectin benzoate 5% SG 80gm or indoxacarb 14.5% SC mixing in 200lit. of water per acre crop. At three-week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by **diamond back moth**. To control this, apply Spinosad 45SC @ 0.5ml/l.

**Tomato:** Transplanted seedlings are at 6-8-week stage in main field. Complete hoeing, weeding and apply 27 kg urea at six-week stage. Irrigate the crops after fertilizer application. To control **tomato leaf curl virus** in field, Spray Imidachloprid 17.8% SL @ 0.5 ml or Dimethoate 30% EC @ 1 ml/litre of 10 water at 15, 25, 45 days after transplanting.

**Potato:** There are chances of Leaf Blight disease incidence in potato crop. To manage **early leaf blight** in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days' interval.

**Onion:** The transplanted seedlings are at seedling establishment to four-week stage in main field. **Irrigate the crop at 10 days' interval.** Go for hoeing, weeding and 1<sup>st</sup> top dressing with 45kg urea per acre crop area at 4-week stage of crop followed by irrigation.

**Animal husbandry:** In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



**Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area**

**DrRanjan Kumar Mohanta**  
**(Nodal officer)**





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
**(A Unit of ICAR-National Rice Research Institute)**  
**Email: [kvkcuttack@gmail.com](mailto:kvkcuttack@gmail.com)/[damukvkcuttack@gmail.com](mailto:damukvkcuttack@gmail.com)**



Bulletin No.-613  
Date- 20/01/2026

**AGROMET ADVISORY BULLETIN FOR KANTAPADA**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 26.14°C (-2.57 from normal) and 10.57°C (-4.4 from normal), respectively. The mean relative humidity was 65-77%.

**Forecasted period: Today(8.30IST) to 25<sup>th</sup>January (8.30IST)2026**

**Weather advisory:** *Dry weather is forecasted for upcoming 5 days in the block.* The day temperature will be 30-31°C and night temperature will be 14-15°C for upcoming 5 days. Maximum and minimum relative humidity will be 88-93% and 24-29% respectively. There will be clear sky for next 5 days.

Block	<b>Kantapada(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)</b>				
	<b>20/01/2026</b>	<b>21/01/2026</b>	<b>22/01/2026</b>	<b>23/01/2026</b>	<b>24/01/2026</b>
Rainfall(mm)	<b>00</b>	<b>00</b>	<b>00</b>	<b>00</b>	<b>00</b>
T-Max(°C)	31	30	30	30	31
T-Min(°C)	<b>14</b>	<b>15</b>	<b>15</b>	<b>15</b>	<b>15</b>
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	88	92	92	93	91
RH Min (%)	25	26	29	29	24
Wind Speed (Kmph)	5	6	7	3	4
Wind Direction (deg)	315	338	301	311	338

**Agromet Advisory**

**GeneralAdvisory:** *As shallow to moderate fog has occurred over isolated places in past 4-5 days and also likely to occur in next 2-3 days, so farmers are advised to monitor potato, tomato and onion crops for incidence of early/late blight disease and mustard for white rust, green and black gram crop for powdery mildew. if symptoms are noticed, apply recommended plant protection measures.*

**Pulses:**

**Farmers should start sowing of pulses by utilization of irrigation water.**

Cloudy and foggy weather may aggravate the infestation of **white fly, aphid and powdery mildew** in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and **Whitefly** and spray 3gm Sulfex or 1ml Propiconazole 10% EC or 1 ml Tebuconazole 25.9%EC per litre water for controlling **Powdery mildew**.

**Rabi-Groundnut:**

**As early sown groundnut crop are at post flowering stage(After 30-40 days of sowing), so farmers are advised to irrigate their field after this stage due to mildly dry condition in Cuttack district in last one month and also in upcoming days.**

Farmers are advised to do intercultivation and apply **Gypsum @200 kg/ acre** in Groundnut which is at flowering stage. To manage **aphids** and **thrips** spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. To manage **tikka** disease in groundnut farmers should spray with 500g Blitox-50 powder in 200 litre water.

**Mustard:** *As dry weather is needed after sowing of mustard, so farmers should complete sowing of Mustard in between upcoming 5 days.*

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of **Aphid** and **stink bug** infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage **Mustard sawfly** and **leafWebber** spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

**Vegetables:**

**Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water. To control the **sucking pests like aphid & white fly**, spray Imidacloprid @ 5ml/10 lit of water.

**Cabbage and Cauliflower:** Current weather is favorable for attack of **Cabbage Webber** and **tobacco caterpillar** in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. **Leaf eating caterpillar and fruit borer** may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectin benzoate 5% SG 80gm or indoxacarb 14.5% SC mixing in 200lit. of water per acre crop. At three-week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by **diamond back moth**. To control this, apply Spinosad 45SC @ 0.5ml/l.

**Tomato:** Transplanted seedlings are at 6-8-week stage in main field. Complete hoeing, weeding and apply 27 kg urea at six-week stage. Irrigate the crops after fertilizer application. To control **tomato leaf curl virus** in field, Spray Imidachloprid 17.8% SL @ 0.5 ml or Dimethoate 30% EC @ 1 ml/litre of 10 water at 15, 25, 45 days after transplanting.

**Potato:** There are chances of Leaf Blight disease incidence in potato crop. To manage **early leaf blight** in potato, spray Mancozeb 75% WP @ 600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days' interval.

**Onion:** The transplanted seedlings are at seedling establishment to four-week stage in main field. **Irrigate the crop at 10 days' interval.** Go for hoeing, weeding and 1<sup>st</sup> top dressing with 45kg urea per acre crop area at 4-week stage of crop followed by irrigation.

**Animal husbandry:** In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

**DrRanjan Kumar Mohanta**  
**(Nodal officer)**





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**

**(A Unit of ICAR-National Rice Research Institute)**  
**Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com**



Bulletin No.-613  
Date- 20/01/2026

**AGROMET ADVISORY BULLETIN FOR MAHANGA**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 26.14°C (-2.57 from normal) and 10.57°C (-4.4 from normal), respectively. The mean relative humidity was 65-77%.

**Forecasted period: Today(8.30IST) to 25<sup>th</sup>January (8.30IST)2026**

**Weather advisory:** Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be and it will be 30-31°C and night temperature will be 13-15°C for upcoming 5 days. Maximum and minimum relative humidity will be 84-91% and 23-28% respectively. There will be clear sky for next 5 days.

Block	Mahanga(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	20/01/2026	21/01/2026	22/01/2026	23/01/2026	24/01/2026
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	31	30	30	30	31
T-Min(°C)	13	14	14	15	14
Cloud Cover(octal)	0	0	0	0	0
RH Max (%)	84	88	90	91	89
RH Min (%)	24	25	28	27	23
Wind Speed (Kmph)	5	5	7	5	5
Wind Direction (deg)	300	326	311	324	299

**Agromet Advisory**

**GeneralAdvisory:** As shallow to moderate fog has occurred over isolated places in past 4-5 days and also likely to occur in next 2-3 days, so farmers are advised to monitor potato, tomato and onion crops for incidence of early/late blight disease and mustard for white rust, green and black gram crop for powdery mildew. If symptoms are noticed, apply recommended plant protection measures.

**Pulses:**

Farmers should start sowing of pulses by utilization of irrigation water.

Cloudy and foggy weather may aggravate the infestation of white fly, aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and Whitefly and spray 3gm Sulfex or 1ml Propiconazole 10% EC or 1 ml Tebuconazole 25.9%EC per litre water for controlling Powdery mildew.

**Rabi-Groundnut:**

As early sown groundnut crop are at post flowering stage(After 30-40 days of sowing), so farmers are advised to irrigate their field after this stage due to mildly dry condition in Cuttack district in last one month and also in upcoming days.

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. To manage aphids and thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. To manage tikka disease in groundnut farmers should spray with 500g Blitox-50 powder in 200 litre water.

**Mustard:** As dry weather is needed after sowing of mustard, so farmers should complete sowing of Mustard in between upcoming 5 days.

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leafWebber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

**Vegetables:**

Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water. To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.

**Cabbage and Cauliflower:** Current weather is favorable for attack of Cabbage Webber and tobacco caterpillar in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. Leaf eating caterpillar and fruit borer may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectin benzoate 5% SG 80gm or indoxacarb 14.5% SC mixing in 200lit. of water per acre crop. At three-week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this, apply Spinosad 45SC @ 0.5ml/l.

**Tomato:** Transplanted seedlings are at 6-8-week stage in main field. Complete hoeing, weeding and apply 27 kg urea at six-week stage. Irrigate the crops after fertilizer application. To control tomato leaf curl virus in field, Spray Imidachloprid 17.8% SL @ 0.5 ml or Dimethoate 30% EC @ 1 ml/litre of 10 water at 15, 25, 45 days after transplanting.

**Potato:** There are chances of Leaf Blight disease incidence in potato crop. To manage early leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage late blight in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days' interval.

**Onion:** The transplanted seedlings are at seedling establishment to four-week stage in main field. Irrigate the crop at 10 days' interval. Go for hoeing, weeding and 1<sup>st</sup> top dressing with 45kg urea per acre crop area at 4-week stage of crop followed by irrigation.

**Animal husbandry:** In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightning alert in your area

DrRanjan Kumar Mohanta  
(Nodal officer)





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**

**(A Unit of ICAR-National Rice Research Institute)**  
**Email: kykcuttack@gmail.com/damukykcuttack@gmail.com**



Bulletin No.-613  
Date- 20/01/2026

**AGROMET ADVISORY BULLETIN FOR NARASINGHPUR**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 26.14°C (-2.57 from normal) and 10.57°C (-4.4 from normal), respectively. The mean relative humidity was 65-77%.

**Forecasted period: Today(8.30IST) to 25<sup>th</sup>January (8.30IST)2026**

**Weather advisory:** *Dry weather is forecasted for upcoming 5 days in the block.* The day temperature will be 30-31°C and night temperature will be 12-13°C for upcoming 5 days. Maximum and minimum relative humidity will be 82-89% and 21-27% respectively. There will be clear sky for next 5 days.

Block	Narasinghpur(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	20/01/2026	21/01/2026	22/01/2026	23/01/2026	24/01/2026
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	30	31	30	30	31
T-Min(°C)	12	13	13	13	13
Cloud Cover(/octa)	0	0	0	0	0
RH Max (%)	84	82	87	89	83
RH Min (%)	21	23	24	27	23
Wind Speed (Kmph)	2	4	4	1	1
Wind Direction (deg)	307	311	331	236	214

**Agromet Advisory**

**GeneralAdvisory:** *As shallow to moderate fog has occurred over isolated places in past 4-5 days and also likely to occur in next 2-3 days, so farmers are advised to monitor potato, tomato and onion crops for incidence of early/late blight disease and mustard for white rust, green and black gram crop for powdery mildew. If symptoms are noticed, apply recommended plant protection measures.*

**Pulses:**

**Farmers should start sowing of pulses by utilization of irrigation water.**

Cloudy and foggy weather may aggravate the infestation of **white fly, aphid and powdery mildew** in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and **Whitefly** and spray 3gm Sulfex or 1ml Propiconazole 10% EC or 1 ml Tebuconazole 25.9%EC per litre water for controlling **Powdery mildew**.

**Rabi-Groundnut:**

**As early sown groundnut crop are at post flowering stage(After 30-40 days of sowing), so farmers are advised to irrigate their field after this stage due to mildly dry condition in Cuttack district in last one month and also in upcoming days.**

Farmers are advised to do intercultivation and apply **Gypsum @200 kg/ acre** in Groundnut which is at flowering stage. To manage **aphids** and **thrips** spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. To manage **tikka** disease in groundnut farmers should spray with 500g Blitox-50 powder in 200 litre water.

**Mustard:** *As dry weather is needed after sowing of mustard, so farmers should complete sowing of Mustard in between upcoming 5 days.*

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of **Aphid** and **stink bug** infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage **Mustard sawfly** and **leafWebber** spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

**Vegetables:**

**Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water. To control the **sucking pests like aphid & white fly**, spray Imidacloprid @ 5ml/10 lit of water.

**Cabbage and Cauliflower:** Current weather is favorable for attack of **Cabbage Webber** and **tobacco caterpillar** in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. **Leaf eating caterpillar and fruit borer** may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectin benzoate 5% SG 80gm or indoxacarb 14.5% SC mixing in 200lit. of water per acre crop. At three-week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by **diamond back moth**. To control this, apply Spinosad 45SC @ 0.5ml/l.

**Tomato:** Transplanted seedlings are at 6-8-week stage in main field. Complete hoeing, weeding and apply 27 kg urea at six-week stage. Irrigate the crops after fertilizer application. To control **tomato leaf curl virus** in field, Spray Imidachloprid 17.8% SL @ 0.5 ml or Dimethoate 30% EC @ 1 ml/litre of 10 water at 15, 25, 45 days after transplanting.

**Potato:** There are chances of Leaf Blight disease incidence in potato crop. To manage **early leaf blight** in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days' interval.

**Onion:** The transplanted seedlings are at seedling establishment to four-week stage in main field. **Irrigate the crop at 10 days' interval.** Go for hoeing, weeding and 1<sup>st</sup> top dressing with 45kg urea per acre crop area at 4-week stage of crop followed by irrigation.

- Animal husbandry:** In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta  
(Nodal officer)





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
**(A Unit of ICAR-National Rice Research Institute)**  
**Email: kykcuttack@gmail.com/damukykcuttack@gmail.com**



Bulletin No.-613  
Date- 20/01/2026

**AGROMET ADVISORY BULLETIN FOR NIALI**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 26.14°C (-2.57 from normal) and 10.57°C (-4.4 from normal), respectively. The mean relative humidity was 65-77%.

**Forecasted period: Today(8.30IST) to 25<sup>th</sup>January (8.30IST)2026**

**Weather advisory:** Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 29-30°C and night temperature will be 15-17°C for upcoming 5 days. Maximum and minimum relative humidity will be 87-94% and 30-37% respectively. There will be clear sky for next 5 days.

Block	Niali(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	20/01/2026	21/01/2026	22/01/2026	23/01/2026	24/01/2026
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	30	30	29	29	30
T-Min(°C)	15	16	16	17	16
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	87	91	92	92	94
RH Min (%)	30	30	33	37	30
Wind Speed(Kmph)	4	8	9	4	6
Wind Direction(deg)	294	344	318	312	343

**Agromet Advisory**

**GeneralAdvisory:** As shallow to moderate fog has occurred over isolated places in past 4-5 days and also likely to occur in next 2-3 days, so farmers are advised to monitor potato, tomato and onion crops for incidence of early/late blight disease and mustard for white rust, green and black gram crop for powdery mildew. If symptoms are noticed, apply recommended plant protection measures.

**Pulses:**

Farmers should start sowing of pulses by utilization of irrigation water.

Cloudy and foggy weather may aggravate the infestation of white fly, aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and Whitefly and spray 3gm Sulfex or 1ml Propiconazole 10% EC or 1 ml Tebuconazole 25.9%EC per litre water for controlling Powdery mildew.

**Rabi-Groundnut:**

As early sown groundnut crop are at post flowering stage (After 30-40 days of sowing), so farmers are advised to irrigate their field after this stage due to mildly dry condition in Cuttack district in last one month and also in upcoming days.

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. To manage aphids and thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. To manage tikka disease in groundnut farmers should spray with 500g Blitox-50 powder in 200 litre water.

**Mustard:** As dry weather is needed after sowing of mustard, so farmers should complete sowing of Mustard in between upcoming 5 days.

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leafWebber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

**Vegetables:** Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water. To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.

**Cabbage and Cauliflower:** Current weather is favorable for attack of Cabbage Webber and tobacco caterpillar in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. Leaf eating caterpillar and fruit borer may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectin benzoate 5% SG 80gm or indoxacarb 14.5% SC mixing in 200lit. of water per acre crop. At three-week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this, apply Spinosad 45SC @ 0.5ml/l.

**Tomato:** Transplanted seedlings are at 6-8-week stage in main field. Complete hoeing, weeding and apply 27 kg urea at six-week stage. Irrigate the crops after fertilizer application. To control tomato leaf curl virus in field, Spray Imidachloprid 17.8% SL @ 0.5 ml or Dimethoate 30% EC @ 1 ml/litre of 10 water at 15, 25, 45 days after transplanting.

**Potato:** There are chances of Leaf Blight disease incidence in potato crop. To manage early leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage late blight in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days' interval.

**Onion:** The transplanted seedlings are at seedling establishment to four-week stage in main field. Irrigate the crop at 10 days' interval. Go for hoeing, weeding and 1<sup>st</sup> top dressing with 45kg urea per acre crop area at 4-week stage of crop followed by irrigation.

- Animal husbandry:** In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals.
- In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightning alert in your area

DrRanjan Kumar Mohanta  
(Nodal officer)





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
**(A Unit of ICAR-National Rice Research Institute)**  
**Email: kykcuttack@gmail.com/damukyvkcuttack@gmail.com**



Bulletin No.-613  
Date- 20/01/2026

**AGROMET ADVISORY BULLETIN FOR NISCHINTAKOILI**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 26.14°C (-2.57 from normal) and 10.57°C (-4.4 from normal), respectively. The mean relative humidity was 65-77%.

**Forecasted period: Today(8.30IST) to 25<sup>th</sup>January (8.30IST)2026**

**Weather advisory:** Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 30-31°C and night temperature will be 13-15°C for upcoming 5 days. Maximum and minimum relative humidity will be 84-92% and 23-28% respectively. There will be clear sky for next 5 days.

Block	Nischtakoli(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	20/01/2026	21/01/2026	22/01/2026	23/01/2026	24/01/2026
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	31	31	30	30	31
T-Min(°C)	13	14	15	15	14
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	84	89	90	92	90
RH Min (%)	23	24	28	28	23
Wind Speed (Kmph)	5	6	7	5	5
Wind Direction (deg)	306	330	315	321	309

**Agromet Advisory**

**General Advisory:** As shallow to moderate fog has occurred over isolated places in past 4-5 days and also likely to occur in next 2-3 days, so farmers are advised to monitor potato, tomato and onion crops for incidence of early/late blight disease and mustard for white rust, green and black gram crop for powdery mildew. If symptoms are noticed, apply recommended plant protection measures.

**Pulses:**

Farmers should start sowing of pulses by utilization of irrigation water.

Cloudy and foggy weather may aggravate the infestation of white fly, aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and Whitefly and spray 3gm Sulfex or 1ml Propiconazole 10% EC or 1 ml Tebuconazole 25.9% EC per litre water for controlling Powdery mildew.

**Rabi-Groundnut:**

As early sown groundnut crop are at post flowering stage (After 30-40 days of sowing), so farmers are advised to irrigate their field after this stage due to mildly dry condition in Cuttack district in last one month and also in upcoming days.

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. To manage aphids and thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. To manage tikka disease in groundnut farmers should spray with 500g Blitox-50 powder in 200 litre water.

**Mustard:** As dry weather is needed after sowing of mustard, so farmers should complete sowing of Mustard in between upcoming 5 days.

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf Webber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

**Vegetables:**

Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin @ 1g/litre or Profenophos @ 2ml/litre of water. To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.

**Cabbage and Cauliflower:** Current weather is favorable for attack of Cabbage Webber and tobacco caterpillar in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm @ 600ml/acre and Emamectin benzoate @ 70g/acre in 200 lit water. Leaf eating caterpillar and fruit borer may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectin benzoate 5% SG 80gm or indoxacarb 14.5% SC mixing in 200lit. of water per acre crop. At three-week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this, apply Spinosad 45SC @ 0.5ml/l.

**Tomato:** Transplanted seedlings are at 6-8-week stage in main field. Complete hoeing, weeding and apply 27 kg urea at six-week stage. Irrigate the crops after fertilizer application. To control tomato leaf curl virus in field, Spray Imidachloprid 17.8% SL @ 0.5 ml or Dimethoate 30% EC @ 1 ml/litre of 10 water at 15, 25, 45 days after transplanting.

**Potato:** There are chances of Leaf Blight disease incidence in potato crop. To manage early leaf blight in potato, spray Mancozeb 75% WP @ 600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage late blight in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400-gram/acre at 10 days' interval.

**Onion:** The transplanted seedlings are at seedling establishment to four-week stage in main field. Irrigate the crop at 10 days' interval. Go for hoeing, weeding and 1<sup>st</sup> top dressing with 45kg urea per acre crop area at 4-week stage of crop followed by irrigation.

• **Animal husbandry:** In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightning alert in your area

**Dr Ranjan Kumar Mohanta**  
**(Nodal officer)**





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
**(A Unit of ICAR-National Rice Research Institute)**  
**Email: kykcuttack@gmail.com/damukykcuttack@gmail.com**



Bulletin No.-613  
Date- 20/01/2026

**AGROMET ADVISORY BULLETIN FOR SALEPUR**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 26.14°C (-2.57 from normal) and 10.57°C (-4.4 from normal), respectively. The mean relative humidity was 65-77%.

**Forecasted period: Today(8.30IST) to 25<sup>th</sup>January (8.30IST)2026**

**Weather advisory:** Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 30-31°C and night temperature will be 13-15°C for upcoming 5 days. Maximum and minimum relative humidity will be 84-92% and 24-28% respectively. There will be clear sky for next 5 days.

Block	Salepur(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	20/01/2026	21/01/2026	22/01/2026	23/01/2026	24/01/2026
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	31	31	30	30	31
T-Min(°C)	13	14	15	15	15
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	84	89	91	92	90
RH Min (%)	24	24	28	28	24
Wind Speed (Kmph)	5	6	7	5	5
Wind Direction (deg)	306	330	315	321	309

**Agromet Advisory**

**GeneralAdvisory:** As shallow to moderate fog has occurred over isolated places in past 4-5 days and also likely to occur in next 2-3 days, so farmers are advised to monitor potato, tomato and onion crops for incidence of early/late blight disease and mustard for white rust, green and black gram crop for powdery mildew. If symptoms are noticed, apply recommended plant protection measures.

**Pulses:**

Farmers should start sowing of pulses by utilization of irrigation water.

Cloudy and foggy weather may aggravate the infestation of white fly, aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and Whitefly and spray 3gm Sulfex or 1ml Propiconazole 10% EC or 1 ml Tebuconazole 25.9%EC per litre water for controlling Powdery mildew.

**Rabi-Groundnut:**

As early sown groundnut crop are at post flowering stage(After 30-40 days of sowing), so farmers are advised to irrigate their field after this stage due to mildly dry condition in Cuttack district in last one month and also in upcoming days.

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. To manage aphids and thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. To manage tikka disease in groundnut farmers should spray with 500g Blitox-50 powder in 200 litre water.

**Mustard:** As dry weather is needed after sowing of mustard, so farmers should complete sowing of Mustard in between upcoming 5 days.

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leafWebber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

**Vegetables:**

Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water. To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.

**Cabbage and Cauliflower:** Current weather is favorable for attack of Cabbage Webber and tobacco caterpillar in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. Leaf eating caterpillar and fruit borer may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectin benzoate 5% SG 80gm or indoxacarb 14.5% SC mixing in 200lit. of water per acre crop. At three-week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this, apply Spinosad 45SC @ 0.5ml/l.

**Tomato:** Transplanted seedlings are at 6-8-week stage in main field. Complete hoeing, weeding and apply 27 kg urea at six-week stage. Irrigate the crops after fertilizer application. To control tomato leaf curl virus in field, Spray Imidachloprid 17.8% SL @ 0.5 ml or Dimethoate 30% EC @ 1 ml/litre of 10 water at 15, 25, 45 days after transplanting.

**Potato:** There are chances of Leaf Blight disease incidence in potato crop. To manage early leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage late blight in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days' interval.

**Onion:** The transplanted seedlings are at seedling establishment to four-week stage in main field. Irrigate the crop at 10 days' interval. Go for hoeing, weeding and 1<sup>st</sup> top dressing with 45kg urea per acre crop area at 4-week stage of crop followed by irrigation.

**Animal husbandry:** In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightning alert in your area

DrRanjan Kumar Mohanta  
(Nodal officer)





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**

**(A Unit of ICAR-National Rice Research Institute)**  
**Email: [kvkcuttack@gmail.com](mailto:kvkcuttack@gmail.com)/[damukvkcuttack@gmail.com](mailto:damukvkcuttack@gmail.com)**



Bulletin No.-613  
Date- 20/01/2026

**AGROMET ADVISORY BULLETIN FOR TANGI-CHOWDWAR**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 26.14°C (-2.57 from normal) and 10.57°C (-4.4 from normal), respectively. The mean relative humidity was 65-77%.

**Forecasted period: Today(8.30IST) to 25<sup>th</sup>January (8.30IST)2026**

**Weather advisory:** *Dry weather is forecasted for upcoming 5 days in the block.* The day temperature will be 30-31°C and night temperature will be 13-15°C for upcoming 5 days. Maximum and minimum relative humidity will be 83-89% and 36-44% respectively. There will be clear sky for next 5 days.

Block	Tangi-Chowdwar(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	20/01/2026	21/01/2026	22/01/2026	23/01/2026	24/01/2026
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	31	31	30	30	31
T-Min(°C)	13	14	15	15	15
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	83	89	89	89	87
RH Min (%)	36	39	44	44	41
Wind Speed (Kmph)	8	10	9	8	7
Wind Direction (deg)	328	335	315	322	323

**Agromet Advisory**

**General Advisory:** *As shallow to moderate fog has occurred over isolated places in past 4-5 days and also likely to occur in next 2-3 days, so farmers are advised to monitor potato, tomato and onion crops for incidence of early/late blight disease and mustard for white rust, green and black gram crop for powdery mildew. If symptoms are noticed, apply recommended plant protection measures.*

**Pulses:**

**Farmers should start sowing of pulses by utilization of irrigation water.**

Cloudy and foggy weather may aggravate the infestation of **white fly, aphid and powdery mildew** in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and **Whitefly** and spray 3gm Sulfex or 1ml Propiconazole 10% EC or 1 ml Tebuconazole 25.9%EC per litre water for controlling **Powdery mildew**.

**Rabi-Groundnut:**

**As early sown groundnut crop are at post flowering stage(After 30-40 days of sowing), so farmers are advised to irrigate their field after this stage due to mildly dry condition in Cuttack district in last one month and also in upcoming days.**

Farmers are advised to do intercultivation and apply **Gypsum @200 kg/ acre** in Groundnut which is at flowering stage. To manage **aphids** and **thrips** spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. To manage **tikka** disease in groundnut farmers should spray with 500g Blitox-50 powder in 200 litre water.

**Mustard:** *As dry weather is needed after sowing of mustard, so farmers should complete sowing of Mustard in between upcoming 5 days.*

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of **Aphid** and **stink bug** infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage **Mustard sawfly** and **leafWebber** spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

**Vegetables:**

**Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water. To control the **sucking pests like aphid & white fly**, spray Imidacloprid @ 5ml/10 lit of water.

**Cabbage and Cauliflower:** Current weather is favorable for attack of **Cabbage Webber** and **tobacco caterpillar** in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. **Leaf eating caterpillar and fruit borer** may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectin benzoate 5% SG 80gm or indoxacarb 14.5% SC mixing in 200lit. of water per acre crop. At three-week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by **diamond back moth**. To control this, apply Spinosad 45SC @ 0.5ml/l.

**Tomato:** Transplanted seedlings are at 6-8-week stage in main field. Complete hoeing, weeding and apply 27 kg urea at six-week stage. Irrigate the crops after fertilizer application. To control **tomato leaf curl virus** in field, Spray Imidachloprid 17.8% SL @ 0.5 ml or Dimethoate 30% EC @ 1 ml/litre of 10 water at 15, 25, 45 days after transplanting.

**Potato:** There are chances of Leaf Blight disease incidence in potato crop. To manage **early leaf blight** in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days' interval.

**Onion:** The transplanted seedlings are at seedling establishment to four-week stage in main field. **Irrigate the crop at 10 days' interval.** Go for hoeing, weeding and 1<sup>st</sup> top dressing with 45kg urea per acre crop area at 4-week stage of crop followed by irrigation.

- Animal husbandry:** In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightning alert in your area

**DrRanjan Kumar Mohanta**  
**(Nodal officer)**





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
**(A Unit of ICAR-National Rice Research Institute)**  
**Email: kykcuttack@gmail.com/damukyvkcuttack@gmail.com**



Bulletin No.-613  
Date- 20/01/2026

**AGROMET ADVISORY BULLETIN FOR TIGIRIA**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 26.14°C (-2.57 from normal) and 10.57°C (-4.4 from normal), respectively. The mean relative humidity was 65-77%.

**Forecasted period: Today(8.30IST) to 25<sup>th</sup>January (8.30IST)2026**

**Weather advisory:** *Dry weather is forecasted for upcoming 5 days in the block.* The day temperature will be 30-31°C and night temperature will be 13-14°C for upcoming 5 days. Maximum and minimum relative humidity will be 89-93% and 22-27% respectively. There will be clear sky for next 5 days.

Block	Tigiria(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	20/01/2026	21/01/2026	22/01/2026	23/01/2026	24/01/2026
Rainfall(mm)	00	00	00	00	00
T-Max(°C)	30	30	30	30	31
T-Min(°C)	13	13	14	14	14
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	89	91	92	93	92
RH Min (%)	22	24	26	27	23
Wind Speed (Kmph)	3	4	5	1	1
Wind Direction (deg)	291	325	321	270	315

**Agro met Advisory**

**GeneralAdvisory:** *As shallow to moderate fog has occurred over isolated places in past 4-5 days and also likely to occur in next 2-3 days, so farmers are advised to monitor potato, tomato and onion crops for incidence of early/late blight disease and mustard for white rust, green and black gram crop for powdery mildew. If symptoms are noticed, apply recommended plant protection measures.*

**Pulses:**

**Farmers should start sowing of pulses by utilization of irrigation water.**

Cloudy and foggy weather may aggravate the infestation of white fly, aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and Whitefly and spray 3gm Sulfex or 1ml Propiconazole 10% EC or 1 ml Tebuconazole 25.9%EC per litre water for controlling Powdery mildew.

**Rabi-Groundnut:**

**As early sown groundnut crop are at post flowering stage(After 30-40 days of sowing), so farmers are advised to irrigate their field after this stage due to mildly dry condition in Cuttack district in last one month and also in upcoming days.**

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. To manage aphids and thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. To manage tikka disease in groundnut farmers should spray with 500g Blitox-50 powder in 200 litre water.

**Mustard:** *As dry weather is needed after sowing of mustard, so farmers should complete sowing of Mustard in between upcoming 5 days.*

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leafWebber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyriphos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

**Vegetables:**

**Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water. To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.

**Cabbage and Cauliflower:** Current weather is favorable for attack of **Cabbage Webber** and **tobacco caterpillar** in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. **Leaf eating caterpillar and fruit borer** may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectin benzoate 5% SG 80gm or indoxacarb 14.5% SC mixing in 200lit. of water per acre crop. At three-week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by **diamond back moth**. To control this, apply Spinosad 45SC @ 0.5ml/l.

**Tomato:** Transplanted seedlings are at 6-8-week stage in main field. Complete hoeing, weeding and apply 27 kg urea at six-week stage. Irrigate the crops after fertilizer application. To control **tomato leaf curl virus** in field, Spray Imidachloprid 17.8% SL @ 0.5 ml or Dimethoate 30% EC @ 1 ml/litre of 10 water at 15, 25, 45 days after transplanting.

**Potato:** There are chances of Leaf Blight disease incidence in potato crop. To manage **early leaf blight** in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days' interval.

**Onion:** The transplanted seedlings are at seedling establishment to four-week stage in main field. **Irrigate the crop at 10 days' interval.** Go for hoeing, weeding and 1<sup>st</sup> top dressing with 45kg urea per acre crop area at 4-week stage of crop followed by irrigation.

• **Animal husbandry:** In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta  
(Nodal officer)






**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
**(A Unit of ICAR-National Rice Research Institute)**  
Email: [kvkcuttack@gmail.com](mailto:kvkcuttack@gmail.com)/[damukvkcuttack@gmail.com](mailto:damukvkcuttack@gmail.com)



Bulletin No.-613  
Date- 20/01/2026

				
MEGHDOOT APP	DAMINI APP	FACEBOOK (Damu Cuttack)	BLOCK SPECIFIC WHAT'S APP GROUPS	@DAMUCUTTACK

- Kindly install and use **“Meghdoot mobile App”** for Cuttack district weather forecast, warnings and weather based agromet advisory services.  
<https://play.google.com/store/apps/details?id=com.aas.meghdoot>
- Kindly install and use **“Damini mobile App”** for location specific for lightening forecast and warnings.  
<https://play.google.com/store/apps/details?id=com.lightening.live.damini>
- Follow the **block specific whats app groups** and join in your block wise groups for getting weather based agromet advisory services twice in a week (Tuesday and Friday) and other weather warning based farm management advisories.
- Follow the **Official Website links** to get the weather based agromet advisory service bulletins (**English & Odia**) twice in a week (Tuesday and Friday).

1	<b>ICAR-NRRI, Cuttack, Odisha</b>	<a href="https://icar-nrri.in/agromate/">https://icar-nrri.in/agromate/</a>	
2	<b>Agrimet division, IMD, Pune</b>	<a href="https://imdagrimet.gov.in/imdproject/AGIndex.php">https://imdagrimet.gov.in/imdproject/AGIndex.php</a>	



Follow Meghdoot Mobile APP for Agromet  
advisories in local language and DAMINI  
APP for Lightening alert in your area

DrRanjan Kumar Mohanta  
(Nodal officer)

