




Cuttack District Block Wise Agro-met Advisory Service Bulletin

*Tuesday, 05th May, 2026
(For the period 05th to 09th May, 2026)*



*Issued by
District Agromet Unit,
KrishiVigyan Kendra Cuttack*




Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





STANDARD PRECIPITATION INDEX AND NDVI MAP



LEGEND (Medium range block level rainfall forecast)		
	No Rainfall	00 mm
	Very light rainfall	01-05 mm
	Light rainfall	06-10 mm
	Light to moderate rainfall	11-30 mm
	Moderate to heavy rainfall	31-65 mm
	Heavy to very heavy rainfall	66-115 mm
	Extremely heavy rainfall	>116 mm

LEGEND (Past week temperature)		
Color code	Temperature	Departure
	Below Normal	< 2°C from normal
	Normal	-2°C to +2°C from normal
	Above Normal	>2°C from normal



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)

















GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-630
Date- 05/05/2026

Weather (Past and Forecasted) based Disease and pest in Major Crops and vegetables

Sl No	Crop	Stage and condition of the crop	Anticipated Disease and pest	
1	Maize	Vegetative to tasseling/silking		Maize stalk rot
				Fall army worm
				Red hairy catpillar
2	Brinjal	Fruiting		Phomopsis Blight
				Cercopera leaf spot
				Helicoverpa
3	Tomato	Fruiting		Bacterial wilt
4	Okra	Vegetative to fruiting		Yellow vein mosaiv(YMV)
5	Cucurbits	Flowering to Fruiting stage		Late blight
				Downy Mildew
				Leaf spot
				Stem Rot



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
 (Nodal officer)











GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-630
Date- 05/05/2026

				Fruit rot
6	Banana			Banana leaf spot/sigatoka
				Pseudo stem weevil
				Rhizome rot
7	Mango	Fruiting stage		Fruit borer
8	Papaya	Fruting		phytophthora crown rot or foot rot



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
 (Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-630
Date- 05/05/2026

AGROMET ADVISORY BULLETIN FOR ATHAGAD

District Past week average weather condition and Block past week average Rainfall condition

The block received average 08 mm and district received average 06 mm rainfall during the last week. The mean daily maximum and minimum temperature was 34.57°C (-2.53 from normal) and 21.14°C (-3.41 from normal), respectively. The mean relative humidity was 75-85%.

Forecasted period: Today(8.30IST) to 10th May (8.30IST)2026

Weather advisory: *Light to Moderate Rainfall / thundershower is forecasted in between today & tomorrow and very light Rainfall / thundershower is forecasted for other 3days in the block..* The day temperature will be 33-38°C and night temperature will be 22-25°C for upcoming 5 days. Maximum and minimum relative humidity will be 81-87% and 31-40% respectively. **There will be partly cloudy to overcast sky for next 5days.**

Block	Athagad (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	05/05/2026	06/05/2026	07/05/2026	08/05/2026	09/05/2026
Rainfall (mm)	16	09	05	03	02
T-Max (°C)	33	35	36	37	38
T-Min (°C)	22	23	24	24	25
Cloud Cover (octa)	5	5	4	3	3
RH Max (%)	87	81	84	86	87
RH Min (%)	39	40	35	31	31
Wind Speed (kmph)	8	11	7	7	14
Wind Direction (deg)	124	164	103	158	171

Agromet Advisory

General advisory: As thunderstorm with lightning /gusty winds (speed 40-50 kmph) with light to moderate rainfall is forecasted in upcoming 5 days Farmers should harvest matured green gram, black gram, Maize, Cucurbits and other vegetables and keep the produce at safer places, provide mechanical support to young fruit plants and vegetables to prevent the crops from lodging due to strong winds, avoid standing in the open or working in the fields during thunderstorm/lightning period. Farmers should pause all farm activities in upcoming 5 days.

Pulses:

Harvest matured pulses as light to moderate rainfall or thundershower is forecasted from today on wards at isolated places of Cuttack district.

Summer Maize:

There may be increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of **fall army worm in maize**. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water. If there will be infestation of **Stem rot** at vegetative stage as rainfall is forecasted, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/ltr of water.

Vegetables: After rain there may be chances of increase in infestation of wilting in vegetables. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter.

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/litr of water. **Cercospera leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water. If **helicoverpa infestation** will be found, spray Flunendiamide or indoxacarb 2ml/litr of water.

Cucurbits:

Late Blight may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: If **rhizome rot** will be noticed in the field, drench the solution of Copper oxy chloride at 3g per L of water. For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil @ 2.0 g / litre of water.

Mango: For control of mango **fruit borer** spray Neem oil 3ml + chloripyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The sprayings should be preferably carried out during evening hours.

Papaya: After rain due to water logging there may be increase in the infestation of **phytopthera crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Animal husbandry:

Make necessary arrangement for clean and sufficient drinking water for the animals. keep farm animals indoor during the thunderstorm period.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAM KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-630
Date- 05/05/2026

AGROMET ADVISORY BULLETIN FOR BADAMBA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 08 mm and district received average 06 mm rainfall during the last week. The mean daily maximum and minimum temperature was 34.57°C (-2.53 from normal) and 21.14°C (-3.41 from normal), respectively. The mean relative humidity was 75-85%.

Forecasted period: Today(8.30IST) to 10th May (8.30IST)2026

Weather advisory: Light to Moderate Rainfall / thundershower is forecasted in between today & tomorrow and very light Rainfall / thundershower is forecasted for other 3days in the block. The day temperature will be 34-38°C and night temperature will be 24-25°C for upcoming 5 days. Maximum and minimum relative humidity will be 77-90% and 23-38% respectively. There will be partly cloudy to overcast sky for next 5days.

Block	Badamba (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	05/05/2026	06/05/2026	07/05/2026	08/05/2026	09/05/2026
Rainfall(mm)	14	08	05	02	01
T-Max(°C)	34	36	37	37	38
T-Min(°C)	24	24	25	25	25
Cloud Cover(octa)	5	5	4	3	3
RH Max (%)	90	78	77	82	83
RH Min (%)	38	25	33	26	23
Wind Speed (Kmph)	9	6	7	6	12
Wind Direction (deg)	114	162	141	121	149

Agromet Advisory

General advisory: As thunderstorm with lightning /gusty winds (speed 40-50 kmph) with light to moderate rainfall is forecasted in upcoming 5 days Farmers should harvest matured green gram, black gram, Maize, Cucurbits and other vegetables and keep the produce at safer places, provide mechanical support to young fruit plants and vegetables to prevent the crops from lodging due to strong winds, avoid standing in the open or working in the fields during thunderstorm/lightning period. Farmers should pause all farm activities in upcoming 5 days.

Pulses:

Harvest matured pulses as light to moderate rainfall or thundershower is forecasted from today on wards at isolated places of Cuttack district.

Summer Maize:

There may be increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of **fall army worm in maize**. To control spray **emamectin benzoate (5% S.G.) @ 0.4g per litre of water**. If there will be infestation of **Stem rot** at vegetative stage as rainfall is forecasted, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or vaidamycin 2ml/ltr of water.

Vegetables:

After rain there may be chances of increase in infestation of wilting in vegetables. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply **Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter**.

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/litr of water. **Cercospera leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water. If **helicopterpa infestation** will be found, spray Flunendiamide or indoxacarb 2ml/litr of water.

Cucurbits:

Late Blight may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: If **rhizome rot** will be noticed in the field, drench the solution of Copper oxy chloride at 3g per L of water. For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil @ 2.0 g / litre of water.

Mango: For control of mango **fruit borer** spray Neem oil 3ml + chloriprifos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours.

Papaya: After rain due to water logging there may be increase in the infestation of **phytopthera crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Animal husbandry:

Make necessary arrangement for clean and sufficient drinking water for the animals. keep farm animals indoor during the thunderstorm period.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
 (Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-630
Date- 05/05/2026

AGROMET ADVISORY BULLETIN FOR BANKI

District Past week average weather condition

The district received average 06 mm rainfall during the last week. The mean daily maximum and minimum temperature was 34.57°C (-2.53 from normal) and 21.14°C (-3.41 from normal), respectively. The mean relative humidity was 75-85%.

Forecasted period: Today(8.30IST) to 10th May (8.30IST)2026

Weather advisory: *Light to Moderate Rainfall / thundershower is forecasted in between today & tomorrow and very light Rainfall / thundershower is forecasted for other 3days in the block.* The day temperature will be 34-38°C and night temperature will be 24-25°C for upcoming 5 days. Maximum and minimum relative humidity will be 82-88% and 32-40% respectively. **There will be partly cloudy to overcast sky for next 5days.**

Block	Banki(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	05/05/2026	06/05/2026	07/05/2026	08/05/2026	09/05/2026
Rainfall(mm)	13	08	05	02	01
T-Max(°C)	34	35	36	37	38
T-Min(°C)	24	25	25	25	25
Cloud Cover(octa)	5	5	4	3	2
RH Max (%)	88	82	84	86	87
RH Min (%)	40	40	36	32	32
Wind Speed (Kmph)	8	9	5	6	11
Wind Direction (deg)	114	175	104	140	167

Agromet Advisory

General advisory: As thunderstorm with lightning /gusty winds (speed 40-50 kmph) with light to moderate rainfall is forecasted in upcoming 5 days Farmers should harvest matured green gram, black gram, Maize, Cucurbits and other vegetables and keep the produce at safer places, provide mechanical support to young fruit plants and vegetables to prevent the crops from lodging due to strong winds, avoid standing in the open or working in the fields during thunderstorm/lightning period. Farmers should pause all farm activities in upcoming 5 days.

Pulses:

Harvest matured pulses as light to moderate rainfall or thundershower is forecasted from today on wards at isolated places of Cuttack district.

Summer Maize:

There may be increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of **fall army worm in maize**. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water. If there will be infestation of **Stem rot** at vegetative stage as rainfall is forecasted, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/ltr of water.

Vegetables:

After rain there may be chances of increase in infestation of wilting in vegetables. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter.

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/litr of water. **Cercopspera leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water. If **helicopterpa infestation** will be found, spray Flunendiamide or indoxacarb 2ml/litr of water.

Cucurbits:

Late Blight may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxchloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: If **rhizome rot** will be noticed in the field, drench the solution of Copper oxy chloride at 3g per L of water. For **leaf spot/sigatoka** as a precautionary measure spray carbendizim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalanil @ 2.0 g / litre of water.

Mango: For control of mango **fruit borer** spray Neem oil 3ml + chloripyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The sprayings should be preferably carried out during evening hours.

Papaya: After rain due to water logging there may be increase in the infestation of **phytopthera crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Animal husbandry:

Make necessary arrangement for clean and sufficient drinking water for the animals. keep farm animals indoor during the thunderstorm period.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-630
Date- 05/05/2026

AGROMET ADVISORY BULLETIN FOR BANKI-DAMPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 04 mm and district received average 06 mm rainfall during the last week. The mean daily maximum and minimum temperature was 34.57°C (-2.53 from normal) and 21.14°C (-3.41 from normal), respectively. The mean relative humidity was 75-85%.

Forecasted period: Today(8.30IST) to 10th May (8.30IST)2026

Weather Advisory: Light to Moderate Rainfall / thundershower is forecasted in between today & tomorrow and very light Rainfall / thundershower is forecasted for other 3days in the block. The day temperature will be 34-38°C and night temperature will be 24-25°C for upcoming 5 days. Maximum and minimum relative humidity will be 82-88% and 32-40% respectively. There will be partly cloudy to overcast sky for next 5days.

Block	Banki-Dampada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	05/05/2026	06/05/2026	07/05/2026	08/05/2026	09/05/2026
Rainfall(mm)	12	08	05	03	01
T-Max(°C)	34	35	36	37	38
T-Min(°C)	24	24	25	25	25
Cloud Cover(octa)	5	4	4	3	2
RH Max (%)	88	82	84	86	87
RH Min (%)	40	40	36	32	32
Wind Speed (Kmph)	8	9	5	6	11
Wind Direction (deg)	114	175	104	140	167

Agromet Advisory

General advisory: As thunderstorm with lightning /gusty winds (speed 40-50 kmph) with light to moderate rainfall is forecasted in upcoming 5 days Farmers should harvest matured green gram, black gram, Maize, Cucurbits and other vegetables and keep the produce at safer places, provide mechanical support to young fruit plants and vegetables to prevent the crops from lodging due to strong winds, avoid standing in the open or working in the fields during thunderstorm/lightning period. Farmers should pause all farm activities in upcoming 5 days.

Pulses:

Harvest matured pulses as light to moderate rainfall or thundershower is forecasted from today on wards at isolated places of Cuttack district.

Summer Maize:

There may be increase the infestation of Red hairy caterpillar in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of fall army worm in maize. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water. If there will be infestation of Stem rot at vegetative stage as rainfall is forecasted, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/ltr of water.

Vegetables:

After rain there may be chances of increase in infestation of wilting in vegetables. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter.

Okra: To control Yellow Vain Mosaic Virus (YVMV) in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

If phomopsis blight will occur in brinjal, then spray zineb75%WP 2gram/litr of water. Cercopspera leaf spot may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water. If helioverpa infestation will be found, spray Flunendiamide or indoxacarb 2ml/litr of water.

Cucurbits:

Late Blight may happen due to rising day temperature. There are chances of infestation of Downy Mildew in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: leaf spot, leaf blight, stem rot, fruit rot) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxchloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: If rhizome rot will be noticed in the field, drench the solution of Copper oxy chloride at 3g per L of water. For leaf spot/sigatoka as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil @ 2.0 g / litre of water.

Mango: For control of mango fruit borer spray Neem oil 3ml + chloripyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The sprayings should be preferably carried out during evening hours.

Papaya: After rain due to water logging there may be increase in the infestation of phytopthera crown rot or foot rot in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Animal husbandry:

Make necessary arrangement for clean and sufficient drinking water for the animals. keep farm animals indoor during the thunderstorm period.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-630
Date- 05/05/2026

AGROMET ADVISORY BULLETIN FOR BARANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 09 mm and district received average 06 mm rainfall during the last week. The mean daily maximum and minimum temperature was 34.57°C (-2.53 from normal) and 21.14°C (-3.41 from normal), respectively. The mean relative humidity was 75-85%.

Forecasted period: Today(8.30IST) to 10th May (8.30IST)2026

Weather advisory: Light to Moderate Rainfall / thundershower is forecasted in between today to Thursday and very light Rainfall / thundershower is forecasted for other 2days in the block. The day temperature will be 33-38°C and night temperature will be 23-25°C for upcoming 5 days. Maximum and minimum relative humidity will be 86-95% and 26-39% respectively. There will be partly cloudy to overcast sky for next 5days.

Block	Baranga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	05/05/2026	06/05/2026	07/05/2026	08/05/2026	09/05/2026
Rainfall(mm)	13	08	06	03	01
T-Max(°C)	33	35	36	37	38
T-Min(°C)	23	24	24	24	25
Cloud Cover	5	5	4	3	3
RH Max (%)	94	86	95	95	95
RH Min (%)	39	34	38	29	26
Wind Speed (Kmph)	7	12	11	8	14
Wind Direction (deg)	109	190	174	155	168

Agromet Advisory

General advisory: As thunderstorm with lightning /gusty winds (speed 40-50 kmph) with light to moderate rainfall is forecasted in upcoming 5 days Farmers should harvest matured green gram, black gram, Maize, Cucurbits and other vegetables and keep the produce at safer places, provide mechanical support to young fruit plants and vegetables to prevent the crops from lodging due to strong winds, avoid standing in the open or working in the fields during thunderstorm/lightning period. Farmers should pause all farm activities in upcoming 5 days.

Pulses:

Harvest matured pulses as light to moderate rainfall or thundershower is forecasted from today on wards at isolated places of Cuttack district.

Summer Maize:

There may be increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of **fall army worm in maize**. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water. If there will be infestation of **Stem rot** at vegetative stage as rainfall is forecasted, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/ltr of water.

Vegetables:

After rain there may be chances of increase in infestation of wilting in vegetables. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter.

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/litr of water. **Cercospora leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water. If **helicopterpa infestation** will be found, spray Flunendiamide or indoxacarb 2ml/litr of water.

Cucurbits:

Late Blight may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxchloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: If **rhizome rot** will be noticed in the field, drench the solution of Copper oxy chloride at 3g per L of water. For **leaf spot/sigatoka** as a precautionary measure spray carbendizim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil @ 2.0 g / litre of water.

Mango: For control of mango **fruit borer** spray Neem oil 3ml + chloripyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours.

Papaya: After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Animal husbandry:

Make necessary arrangement for clean and sufficient drinking water for the animals. keep farm animals indoor during the thunderstorm period.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-630
Date- 05/05/2026

AGROMET ADVISORY BULLETIN FOR CUTTACK SADAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 07 mm and district received average 06 mm rainfall during the last week. The mean daily maximum and minimum temperature was 34.57°C (-2.53 from normal) and 21.14°C (-3.41 from normal), respectively. The mean relative humidity was 75-85%.

Forecasted period: Today(8.30IST) to 10th May (8.30IST)2026

Weather advisory: Light to Moderate Rainfall / thundershower is forecasted in between today to Thursday and very light Rainfall / thundershower is forecasted for other 2days in the block. The day temperature will be 33-38°C and night temperature will be 22-24°C for upcoming 5 days. Maximum and minimum relative humidity will be 82-88% and 31-42% respectively. There will be partly cloudy to overcast sky for next 5days.

Block	Cuttack sadar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	05/05/2026	06/05/2026	07/05/2026	08/05/2026	09/05/2026
Rainfall(mm)	14	09	06	03	01
T-Max(°C)	33	35	36	37	38
T-Min(°C)	22	23	24	24	24
Cloud Cover(octa)	5	5	4	3	3
RH Max (%)	86	82	86	86	88
RH Min (%)	39	42	35	32	31
Wind Speed (Kmph)	7	10	7	9	13
Wind Direction (deg)	131	172	139	178	182

Agromet Advisory

General advisory: As thunderstorm with lightning /gusty winds (speed 40-50 kmph) with light to moderate rainfall is forecasted in upcoming 5 days Farmers should harvest matured green gram, black gram, Maize, Cucurbits and other vegetables and keep the produce at safer places, provide mechanical support to young fruit plants and vegetables to prevent the crops from lodging due to strong winds, avoid standing in the open or working in the fields during thunderstorm/lightning period. Farmers should pause all farm activities in upcoming 5 days.

Pulses:

Harvest matured pulses as light to moderate rainfall or thundershower is forecasted from today on wards at isolated places of Cuttack district.

Summer Maize:

There may be increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of **fall army worm in maize**. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water. If there will be infestation of **Stem rot** at vegetative stage as rainfall is forecasted, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/ltr of water.

Vegetables:

After rain there may be chances of increase in infestation of wilting in vegetables. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter.

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/litr of water. **Cercospora leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water. If **helicopterpa infestation** will be found, spray Flunendiamide or indoxacarb 2ml/litr of water.

Cucutbits:

Late Blight may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxchloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: If **rhizome rot** will be noticed in the field, drench the solution of Copper oxy chloride at 3g per L of water. For **leaf spot/sigatoka** as a precautionary measure spray carbendizim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil @ 2.0 g / litre of water.

Mango: For control of mango **fruit borer** spray Neem oil 3ml + chloripyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours.

Papaya: After rain due to water logging there may be increase in the infestation of **phytopthera crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Animal husbandry:

Make necessary arrangement for clean and sufficient drinking water for the animals. keep farm animals indoor during the thunderstorm period.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAM KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-630
Date- 05/05/2026

AGROMET ADVISORY BULLETIN FOR KANTAPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 05 mm and district received average 06 mm rainfall during the last week. The mean daily maximum and minimum temperature was 34.57°C (-2.53 from normal) and 21.14°C (-3.41 from normal), respectively. The mean relative humidity was 75-85%.

Forecasted period: Today(8.30IST) to 10th May (8.30IST)2026

Weather advisory: Light to Moderate Rainfall / thundershower is forecasted in between today to Thursday and very light Rainfall / thundershower is forecasted for other 2days in the block. The day temperature will be 32-37°C and night temperature will be 23-25°C for upcoming 5 days. Maximum and minimum relative humidity will be 84-89% and 37-48% respectively. There will be partly cloudy to overcast sky for next 5days.

Block	Kantapada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	05/05/2026	06/05/2026	07/05/2026	08/05/2026	09/05/2026
Rainfall(mm)	11	08	06	03	02
T-Max(°C)	32	34	35	36	37
T-Min(°C)	23	24	24	24	25
Cloud Cover(octa)	5	5	4	3	3
RH Max (%)	87	84	87	88	89
RH Min (%)	44	48	42	37	37
Wind Speed (Kmph)	6	11	9	11	13
Wind Direction (deg)	135	171	185	186	183

Agromet Advisory

General advisory: As thunderstorm with lightning /gusty winds (speed 40-50 kmph) with light to moderate rainfall is forecasted in upcoming 5 days Farmers should harvest matured green gram, black gram, Maize, Cucurbits and other vegetables and keep the produce at safer places, provide mechanical support to young fruit plants and vegetables to prevent the crops from lodging due to strong winds, avoid standing in the open or working in the fields during thunderstorm/lightning period. Farmers should pause all farm activities in upcoming 5 days.

Pulses:

Harvest matured pulses as light to moderate rainfall or thundershower is forecasted from today on wards at isolated places of Cuttack district.

Summer Maize:

There may be increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of **fall army worm in maize**. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water. If there will be infestation of **Stem rot** at vegetative stage as rainfall is forecasted, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/ltr of water.

Vegetables:

After rain there may be chances of increase in infestation of wilting in vegetables. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter.

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/litr of water. **Cercopspera leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water. If **helicopterpa infestation** will be found, spray Flunendiamide or indoxacarb 2ml/litr of water.

Cucurbits:

Late Blight may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: If **rhizome rot** will be noticed in the field, drench the solution of Copper oxy chloride at 3g per L of water. For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil@ 2.0 g / litre of water.

Mango: For control of mango **fruit borer** spray Neem oil 3ml + chloripyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours.

Papaya: After rain due to water logging there may be increase in the infestation of **phytopthera crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Animal husbandry:

Make necessary arrangement for clean and sufficient drinking water for the animals. keep farm animals indoor during the thunderstorm period.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-630
Date- 05/05/2026

AGROMET ADVISORY BULLETIN FOR MAHANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 07 mm and district received average 06 mm rainfall during the last week. The mean daily maximum and minimum temperature was 34.57°C (-2.53 from normal) and 21.14°C (-3.41 from normal), respectively. The mean relative humidity was 75-85%.

Forecasted period: Today(8.30IST) to 10th May (8.30IST)2026

Weather advisory: Light to Moderate Rainfall / thundershower is forecasted in between today to Thursday and very light Rainfall / thundershower is forecasted for other 2days in the block.. The day temperature will be 32-37°C and night temperature will be 22-25°C for upcoming 5 days. Maximum and minimum relative humidity will be 79-86% and 33-43% respectively. **There will be partly cloudy to overcast sky for next 5days.**

Block	Mahanga(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	05/05/2026	06/05/2026	07/05/2026	08/05/2026	09/05/2026
Rainfall(mm)	16	09	06	04	02
T-Max(°C)	32	34	35	36	37
T-Min(°C)	22	23	24	24	25
Cloud Cover(octal)	5	5	4	3	3
RH Max (%)	84	79	84	84	86
RH Min (%)	41	43	36	34	33
Wind Speed (Kmph)	6	11	9	9	13
Wind Direction (deg)	142	186	164	180	188

Agromet Advisory

General advisory: As thunderstorm with lightning /gusty winds (speed 40-50 kmph) with light to moderate rainfall is forecasted in upcoming 5 days Farmers should harvest matured green gram, black gram, Maize, Cucurbits and other vegetables and keep the produce at safer places, provide mechanical support to young fruit plants and vegetables to prevent the crops from lodging due to strong winds, avoid standing in the open or working in the fields during thunderstorm/lightning period. Farmers should pause all farm activities in upcoming 5 days.

Pulses:

Harvest matured pulses as light to moderate rainfall or thundershower is forecasted from today on wards at isolated places of Cuttack district.

Summer Maize:

There may be increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of **fall army worm in maize**. To control spray **emamectin benzoate (5% S.G.) @ 0.4g per litre of water**. If there will be infestation of **Stem rot** at vegetative stage as rainfall is forecasted, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or vaidamycin 2ml/litr of water.

Vegetables:

After rain there may be chances of increase in infestation of **wilting in vegetables**. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply **Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter**.

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/litr of water. **Cercopspera leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water. If **helicopterpa infestation** will be found, spray Flunendiamide or indoxacarb 2ml/litr of water.

Cucurbits:

Late Blight may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: If **rhizome rot** will be noticed in the field, drench the solution of Copper oxy chloride at 3g per L of water. For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil@ 2.0 g / litre of water.

Mango: For control of mango **fruit borer** spray Neem oil 3ml + chloripyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours.

Papaya: After rain due to water logging there may be increase in the infestation of **phytopthera crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Animal husbandry:

Make necessary arrangement for clean and sufficient drinking water for the animals. keep farm animals indoor during the thunderstorm period.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-630
Date- 05/05/2026

AGROMET ADVISORY BULLETIN FOR NARASINGHPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 09 mm and district received average 06 mm rainfall during the last week. The mean daily maximum and minimum temperature was 34.57°C (-2.53 from normal) and 21.14°C (-3.41 from normal), respectively. The mean relative humidity was 75-85%.

Forecasted period: Today(8.30IST) to 10th May (8.30IST)2026

Weather advisory: Light to Moderate Rainfall / thundershower is forecasted in between today & tomorrow and very light Rainfall / thundershower is forecasted for other 3days in the block. The day temperature will be 35-39°C and night temperature will be 24-25°C for upcoming 5 days. Maximum and minimum relative humidity will be 79-82% and 29-41% respectively. There will be partly cloudy to overcast sky for next 5days.

Block	Narasinghpur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	05/05/2026	06/05/2026	07/05/2026	08/05/2026	09/05/2026
Rainfall(mm)	14	08	05	02	01
T-Max(°C)	35	36	37	38	39
T-Min(°C)	24	24	25	25	25
Cloud Cover(octa)	5	5	4	3	3
RH Max (%)	90	82	80	81	79
RH Min (%)	41	38	37	29	29
Wind Speed (Kmph)	6	2	4	3	6
Wind Direction (deg)	111	194	101	135	159

Agromet Advisory

General advisory: As thunderstorm with lightning /gusty winds (speed 40-50 kmph) with light to moderate rainfall is forecasted in upcoming 5 days Farmers should harvest matured green gram, black gram, Maize, Cucurbits and other vegetables and keep the produce at safer places, provide mechanical support to young fruit plants and vegetables to prevent the crops from lodging due to strong winds, avoid standing in the open or working in the fields during thunderstorm/lightning period. Farmers should pause all farm activities in upcoming 5 days.

Pulses:

Harvest matured pulses as light to moderate rainfall or thundershower is forecasted from today on wards at isolated places of Cuttack district.

Summer Maize:

There may be increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of **fall army worm in maize**. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water. If there will be infestation of **Stem rot** at vegetative stage as rainfall is forecasted, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/ltr of water.

Vegetables:

After rain there may be chances of increase in infestation of wilting in vegetables. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter.

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/litr of water. **Cercospora leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water. If **helicopterpa infestation** will be found, spray Flunendiamide or indoxacarb 2ml/litr of water.

Cucurbits:

Late Blight may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxchloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: If **rhizome rot** will be noticed in the field, drench the solution of Copper oxy chloride at 3g per L of water. For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalanil @ 2.0 g/ litre of water.

Mango: For control of mango **fruit borer** spray Neem oil 3ml + chloripyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours.

Papaya: After rain due to water logging there may be increase in the infestation of **phytopthera crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Animal husbandry:

Make necessary arrangement for clean and sufficient drinking water for the animals. keep farm animals indoor during the thunderstorm period.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-630
Date- 05/05/2026

AGROMET ADVISORY BULLETIN FOR NIALI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 06 mm and district received average 06 mm rainfall during the last week. The mean daily maximum and minimum temperature was 34.57°C (-2.53 from normal) and 21.14°C (-3.41 from normal), respectively. The mean relative humidity was 75-85%.

Forecasted period: Today(8.30IST) to 10th May (8.30IST)2026

Weather advisory: Light to Moderate Rainfall / thundershower is forecasted in between today to Thursday and very light Rainfall / thundershower is forecasted for other 2days in the block. The day temperature will be 31-36°C and night temperature will be 23-25°C for upcoming 5 days. Maximum and minimum relative humidity will be 87-95% and 37-45% respectively. There will be partly cloudy to overcast sky for next 5days.

Block	Niali (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	05/05/2026	06/05/2026	07/05/2026	08/05/2026	09/05/2026
Rainfall(mm)	11	07	07	04	02
T-Max(°C)	31	33	34	35	36
T-Min(°C)	23	24	24	24	25
Cloud Cover(octa)	5	5	5	4	3
RH Max (%)	94	87	93	94	95
RH Min (%)	45	43	45	37	37
Wind Speed(Kmph)	4	12	14	9	12
Wind Direction(deg)	125	187	192	180	182

Agromet Advisory

General advisory: As thunderstorm with lightning /gusty winds (speed 40-50 kmph) with light to moderate rainfall is forecasted in upcoming 5 days Farmers should harvest matured green gram, black gram, Maize, Cucurbits and other vegetables and keep the produce at safer places, provide mechanical support to young fruit plants and vegetables to prevent the crops from lodging due to strong winds, avoid standing in the open or working in the fields during thunderstorm/lightning period. Farmers should pause all farm activities in upcoming 5 days.

Pulses:

Harvest matured pulses as light to moderate rainfall or thundershower is forecasted from today on wards at isolated places of Cuttack district.

Summer Maize:

There may be increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of **fall army worm in maize**. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water. If there will be infestation of **Stem rot** at vegetative stage as rainfall is forecasted, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/ltr of water.

Vegetables:

After rain there may be chances of increase in infestation of wilting in vegetables. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter.

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/litr of water. **Cercospora leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water. If **helicopterpa infestation** will be found, spray Flunendiamide or indoxacarb 2ml/litr of water.

Cucurbits:

Late Blight may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit: Banana: If **rhizome rot** will be noticed in the field, drench the solution of Copper oxy chloride at 3g per L of water. For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalanyl @ 2.0 g/ litre of water.

Mango: For control of mango **fruit borer** spray Neem oil 3ml + chlorpyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours.

Papaya: After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Animal husbandry: Make necessary arrangement for clean and sufficient drinking water for the animals. keep farm animals indoor during the thunderstorm period.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-630
Date- 05/05/2026

AGROMET ADVISORY BULLETIN FOR NISCHINTAKOILI
District Past week average weather condition and Block past week average Rainfall condition

The block received average 04 mm and district received average 06 mm rainfall during the last week. The mean daily maximum and minimum temperature was 34.57°C (-2.53 from normal) and 21.14°C (-3.41 from normal), respectively. The mean relative humidity was 75-85%.

Forecasted period: Today(8.30IST) to 10th May (8.30IST)2026

Weather advisory: *Light to Moderate Rainfall / thundershower is forecasted in between today to Thursday and very light Rainfall / thundershower is forecasted for other 2days in the block.* The day temperature will be 32-36°C and night temperature will be 22-25°C for upcoming 5 days. Maximum and minimum relative humidity will be 80-87% and 34-43% respectively. **There will be partly cloudy to overcast sky for next 5days.**

Block	Nischintakoili (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	05/05/2026	06/05/2026	07/05/2026	08/05/2026	09/05/2026
Rainfall(mm)	13	08	07	04	02
T-Max(°C)	32	33	35	36	36
T-Min(°C)	22	23	24	24	25
Cloud Cover(octa)	5	5	4	3	3
RH Max (%)	85	80	84	85	87
RH Min (%)	41	43	37	34	34
Wind Speed (Kmph)	5	11	9	9	13
Wind Direction (deg)	143	180	164	182	188

Agromet Advisory

General advisory: As thunderstorm with lightning /gusty winds (speed 40-50 kmph) with light to moderate rainfall is forecasted in upcoming 5 days Farmers should harvest matured green gram, black gram, Maize, Cucurbits and other vegetables and keep the produce at safer places, provide mechanical support to young fruit plants and vegetables to prevent the crops from lodging due to strong winds, avoid standing in the open or working in the fields during thunderstorm/lightning period. Farmers should pause all farm activities in upcoming 5 days.

Pulses:

Harvest matured pulses as light to moderate rainfall or thundershower is forecasted from today on wards at isolated places of Cuttack district.

Summer Maize:

There may be increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of **fall army worm in maize**. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water. If there will be infestation of **Stem rot** at vegetative stage as rainfall is forecasted, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/ltr of water.

Vegetables:

After rain there may be chances of increase in infestation of wilting in vegetables. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply **Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter.**

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/litr of water. **Cercospora leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water. If **helicopterpa infestation** will be found, spray Flunendiamide or indoxacarb 2ml/litr of water.

Cucurbits:

Late Blight may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: If **rhizome rot** will be noticed in the field, drench the solution of Copper oxy chloride at 3g per L of water. For **leaf spot/sigatoka** as a precautionary measure spray carbendizim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalanil @ 2.0 g / litre of water.

Mango: For control of mango **fruit borer** spray Neem oil 3ml + chloripyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The sprayings should be preferably carried out during evening hours.

Papaya: After rain due to water logging there may be increase in the infestation of **phytopthera crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Animal husbandry:

Make necessary arrangement for clean and sufficient drinking water for the animals. keep farm animals indoor during the thunderstorm period.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-630
Date- 05/05/2026

AGROMET ADVISORY BULLETIN FOR SALEPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 08 mm and district received average 06 mm rainfall during the last week. The mean daily maximum and minimum temperature was 34.57°C (-2.53 from normal) and 21.14°C (-3.41 from normal), respectively. The mean relative humidity was 75-85%.

Forecasted period: Today(8.30IST) to 10th May (8.30IST)2026

Weather advisory: Light to Moderate Rainfall / thundershower is forecasted in between today to Thursday and very light Rainfall / thundershower is forecasted for other 2days in the block. The day temperature will be 33-37°C and night temperature will be 22-24°C for upcoming 5 days. Maximum and minimum relative humidity will be 80-87% and 34-43% respectively. **There will be partly cloudy to overcast sky for next 5days.**

Block	Salepur(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	05/05/2026	06/05/2026	07/05/2026	08/05/2026	09/05/2026
Rainfall(mm)	15	09	06	04	01
T-Max(°C)	33	34	36	37	37
T-Min(°C)	22	23	24	24	24
Cloud Cover(octa)	5	5	4	3	3
RH Max (%)	85	80	85	85	87
RH Min (%)	41	43	37	34	34
Wind Speed (Kmph)	6	10	9	9	13
Wind Direction (deg)	140	178	159	180	188

Agromet Advisory

General advisory: As thunderstorm with lightning /gusty winds (speed 40-50 kmph) with light to moderate rainfall is forecasted in upcoming 5 days Farmers should harvest matured green gram, black gram, Maize, Cucurbits and other vegetables and keep the produce at safer places, provide mechanical support to young fruit plants and vegetables to prevent the crops from lodging due to strong winds, avoid standing in the open or working in the fields during thunderstorm/lightning period. Farmers should pause all farm activities in upcoming 5 days.

Pulses:

Harvest matured pulses as light to moderate rainfall or thundershower is forecasted from today on wards at isolated places of Cuttack district.

Summer Maize:

There may be increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of **fall army worm in maize**. To control spray **emamectin benzoate (5% S.G.) @ 0.4g per litre of water**. If there will be infestation of **Stem rot** at vegetative stage as rainfall is forecasted, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/ltr of water.

Vegetables:

After rain there may be chances of increase in infestation of wilting in vegetables. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter.

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

If phomopsis blight will occur in brinjal, then spray zineb75%WP 2gram/litr of water.**Cercopspera leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water.If **helicopterpa infestation** will be found, sprayFlunendiamide or indoxacarb 2ml/litr of water.

Cucurbits:

Late Blight may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: If **rhizome rot** will be noticed in the field, drench the solution of Copper oxy chloride at 3g per L of water.For **leaf spot/sigatoka** as a precautionary measure spray carbendizim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil@ 2.0 g / litre of water.

Mango: For control of mango **fruit borer** spray Neem oil 3ml + chloripyrifos 1 ml perlitre of water along with stickers at marble stage of the fruit. The sprayingshould be preferably carried out during evening hours.

Papaya: After rain due to water logging there may be increase in the infestation of **phytopthera crown rot or foot rot** in papya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Animal husbandry:

Make necessary arrangement for clean and sufficient drinking water for the animals. keep farm animals indoor during the thunderstorm period.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-630
Date- 05/05/2026

AGROMET ADVISORY BULLETIN FOR TANGI-CHOUWAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 06 mm and district received average 06 mm rainfall during the last week. The mean daily maximum and minimum temperature was 34.57°C (-2.53 from normal) and 21.14°C (-3.41 from normal), respectively. The mean relative humidity was 75-85%.

Forecasted period: Today(8.30IST) to 10th May (8.30IST)2026

Weather advisory: Light to Moderate Rainfall / thundershower is forecasted in between today to Thursday and very light Rainfall / thundershower is forecasted for other 2days in the block. The day temperature will be 33-38°C and night temperature will be 22-24°C for upcoming 5 days. Maximum and minimum relative humidity will be 85-88% and 52-58% respectively. There will be partly cloudy to overcast sky for next 5days.

Block	Tangi-Choudwar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	05/05/2026	06/05/2026	07/05/2026	08/05/2026	09/05/2026
Rainfall(mm)	16	09	06	03	01
T-Max(°C)	33	34	36	37	38
T-Min(°C)	22	23	24	24	24
Cloud Cover(octa)	5	5	4	3	3
RH Max (%)	87	85	86	87	88
RH Min (%)	58	58	56	52	53
Wind Speed (Kmph)	7	15	12	11	13
Wind Direction (deg)	174	172	191	200	200

Agromet Advisory

General advisory: As thunderstorm with lightning /gusty winds (speed 40-50 kmph) with light to moderate rainfall is forecasted in upcoming 5 days Farmers should harvest matured green gram, black gram, Maize, Cucurbits and other vegetables and keep the produce at safer places, provide mechanical support to young fruit plants and vegetables to prevent the crops from lodging due to strong winds, avoid standing in the open or working in the fields during thunderstorm/lightning period. Farmers should pause all farm activities in upcoming 5 days.

Pulses:

Harvest matured pulses as light to moderate rainfall or thundershower is forecasted from today on wards at isolated places of Cuttack district.

Summer Maize:

There may be increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of **fall army worm in maize**. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water. If there will be infestation of **Stem rot** at vegetative stage as rainfall is forecasted, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/ltr of water.

Vegetables:

After rain there may be chances of increase in infestation of wilting in vegetables. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter.

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/litr of water. **Cercospora leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water. If **helicopterpa infestation** will be found, spray Flunendiamide or indoxacarb 2ml/litr of water.

Cucurbits:

Late Blight may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: If **rhizome rot** will be noticed in the field, drench the solution of Copper oxy chloride at 3g per L of water. For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil @ 2.0 g / litre of water.

Mango: For control of mango **fruit borer** spray Neem oil 3ml + chloripyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours.

Papaya: After rain due to water logging there may be increase in the infestation of **phytopthera crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Animal husbandry:

Make necessary arrangement for clean and sufficient drinking water for the animals. keep farm animals indoor during the thunderstorm period.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-630
Date- 05/05/2026

AGROMET ADVISORY BULLETIN FOR TIGIRIA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 03 mm and district received average 06 mm rainfall during the last week. The mean daily maximum and minimum temperature was 34.57°C (-2.53 from normal) and 21.14°C (-3.41 from normal), respectively. The mean relative humidity was 75-85%.

Forecasted period: Today(8.30IST) to 10th May (8.30IST)2026

Weather advisory: Light to Moderate Rainfall / thundershower is forecasted in between today & tomorrow and very light Rainfall / thundershower is forecasted for other 3days in the block. The day temperature will be 33-38°C and night temperature will be 23-25°C for upcoming 5 days. Maximum and minimum relative humidity will be 81-88 and 31-40% respectively. There will be partly cloudy to overcast sky for next 5days.

Block	Tigiria (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	05/05/2026	06/05/2026	07/05/2026	08/05/2026	09/05/2026
Rainfall(mm)	16	08	05	02	01
T-Max(°C)	33	35	36	37	38
T-Min(°C)	23	24	24	24	25
Cloud Cover(octa)	5	5	4	3	3
RH Max (%)	88	81	83	86	86
RH Min (%)	40	39	35	31	32
Wind Speed (Kmph)	8	9	5	5	12
Wind Direction (deg)	117	175	86	135	166

Agromet Advisory

General advisory: As thunderstorm with lightning /gusty winds (speed 40-50 kmph) with light to moderate rainfall is forecasted in upcoming 5 days Farmers should harvest matured green gram, black gram, Maize, Cucurbits and other vegetables and keep the produce at safer places, provide mechanical support to young fruit plants and vegetables to prevent the crops from lodging due to strong winds, avoid standing in the open or working in the fields during thunderstorm/lightning period. Farmers should pause all farm activities in upcoming 5 days.

Pulses:

Harvest matured pulses as light to moderate rainfall or thundershower is forecasted from today on wards at isolated places of Cuttack district.

Summer Maize:

There may be increase the infestation of Red hairy caterpillar in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of fall army worm in maize. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water. If there will be infestation of Stem rot at vegetative stage as rainfall is forecasted, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/ltr of water.

Vegetables:

After rain there may be chances of increase in infestation of wilting in vegetables. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter.

Okra: To control Yellow Vain Mosaic Virus (YVMV) in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

If phomopsis blight will occur in brinjal, then spray zineb75%WP 2gram/litr of water. Cercopspera leaf spot may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water. If helioverpa infestation will be found, spray Flunendiamide or indoxacarb 2ml/litr of water.

Cucurbits:

Late Blight may happen due to rising day temperature. There are chances of infestation of Downy Mildew in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: leaf spot, leaf blight, stem rot, fruit rot) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: If rhizome rot will be noticed in the field, drench the solution of Copper oxy chloride at 3g per L of water. For leaf spot/sigatoka as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil @ 2.0 g / litre of water.

Mango: For control of mango fruit borer spray Neem oil 3ml + chloripyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The sprayings should be preferably carried out during evening hours.

Papaya: After rain due to water logging there may be increase in the infestation of phytopthera crown rot or foot rot in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Animal husbandry:

Make necessary arrangement for clean and sufficient drinking water for the animals. keep farm animals indoor during the thunderstorm period.



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)








GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-630
Date- 05/05/2026

				
MEGHDOOT APP	DAMINI APP	FACEBOOK (Damu Cuttack)	BLOCK SPECIFIC WHAT'S APP GROUPS	@DAMUCUTTACK

- Kindly install and use **“Meghdoot mobile App”** for Cuttack district weather forecast, warnings and weather based agromet advisory services.
<https://play.google.com/store/apps/details?id=com.aas.meghdoot>
- Kindly install and use **“Damini mobile App”** for location specific for lightening forecast and warnings.
<https://play.google.com/store/apps/details?id=com.lightening.live.damini>
- Follow the **block specific whats app groups** and join in your block wise groups for getting weather based agromet advisory services twice in a week (Tuesday and Friday) and other weather warning based farm management advisories.
- Follow the **Official Website links** to get the weather based agromet advisory service bulletins (**English & Odia**) twice in a week (Tuesday and Friday).

1	ICAR-NRRI, Cuttack, Odisha	https://icar-nrri.in/agromate/	
2	Agrimet division, IMD, Pune	https://imdagrmet.gov.in/imdproject/AGIndex.php	



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

DrRanjan Kumar Mohanta
(Nodal officer)

