




Cuttack District Block Wise Agro-met Advisory Service Bulletin

*Tuesday, 01st September, 2026
(For the period 01st to 05th September, 2026)*



*Issued by
District Agromet Unit,
KrishiVigyan Kendra Cuttack*





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-647
Date- 01/09/2026

STANDARD PRECIPITATION INDEX AND NDVI MAP

Standard Precipitation Index (SPI)	Normalized difference vegetation Index (NDVI)
MINISTRY OF EARTH SCIENCES INDIA METEOROLOGICAL DEPARTMENT HYDROMET SECTION, CAUI, CRS PUNE STANDARDIZED PRECIPITATION INDEX (SPI) RECENT 4 WEEK 23 JULY 2026 TO 19 AUGUST 2026 SPI Values - No Data <= -2 (Extremely Dry) -1.5 to -1.99 (Severely Dry) -1 to -1.49 (Moderately Dry) 0 to -0.99 (Mildly Dry) 0 to 0.99 (Mildly Wet) 1 to 1.49 (Moderately Wet) 1.5 to 1.99 (Severely Wet) >= 2 (Extremely Wet)	NOAA/VIIRS/BLENDED NDVI Composite ending on Week no 32 (06.08.2026 to 12.08.2026) over Agricultural region Odisha Agriculture vigour is moderate in the state. Legend: -9999 - 0 0 - 0.05 0.05 - 0.1 0.1 - 0.15 0.15 - 0.2 0.2 - 0.25 0.25 - 0.3 0.3 - 0.35 0.35 - 0.4 0.4 - 0.45 0.45 - 0.5 0.5 - 0.55 0.55 - 0.6 > 0.6 Bare soil, rocks clouds, rain, snow, etc. Moderate Good Very good
<i>As per SPI value (cumulative 4weeks -23 July to 19 August, 2026), Mildly Wet condition was there in Cuttack district.</i>	<i>According to NDVI map (06.08.2026 to 12.08.2026) the agriculture vigor was Very light to light over most places of Cuttack district.</i>

LEGEND (Medium range block level rainfall forecast)		
	No Rainfall	00 mm
	Very light rainfall	01-05 mm
	Light rainfall	06-10 mm
	Light to moderate rainfall	11-30 mm
	Moderate to heavy rainfall	31-65 mm
	Heavy to very heavy rainfall	66-115 mm
	Extremely heavy rainfall	>116 mm

DrRanjan Kumar Mohanta
 (Nodal officer)

**Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area**






GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT

KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-647
Date- 01/09/2026

Weather (Past and Forecasted) based Disease and pest in Major Crops and vegetables

Sl No	Crop	Stage and condition of the crop	Anticipated Disease and pest	
1	Rice	Tillering		Leaf folder
				Thrips
				Rice leaf streak
2	Sugarcane	Vegetative to Tillering		Red rot
				Early shoot borer
3	Maize	3-5 week stage		Stem fly
				Leaf blight
				Stalk rot
				Aphid and thrips
4	Brinjal, tomato, chilli	Seedling		Seedling rot

DrRanjan Kumar Mohanta
 (Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-647
Date- 01/09/2026

				Stem rot
5	Tomato and brinjal	Fruting		Bacterial wilt
6	Chilli	seedling		Damping off
7	Pointed gourd			Fruit rot
8	Pumpkin(Cucurbits)	Flowering to fruting		Stem and root rot



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-647
Date- 01/09/2026

AGROMET ADVISORY BULLETIN FOR ATHAGAD

District Past week average weather condition and Block past week average Rainfall condition

The block received average 12 mm and district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.28°C (-0.72) and 23.57°C (-0.63), respectively. The mean relative humidity was 88-92%.

Forecasted period: Today(8.30IST) to 06th September (8.30IST)2026

Weather advisory: Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block. The day temperature will be 31-33°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 85-90% and 48-72% respectively. **There will be partly cloudy to overcast sky for next 5days.**

Block	Athagad (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	01/09/2026	02/09/2026	03/09/2026	04/09/2026	05/09/2026
Rainfall (mm)	12	9	8	8	7
T-Max (°C)	32	31	32	33	33
T-Min (°C)	25	25	25	26	26
Cloud Cover (octa)	6	6	6	5	5
RH Max (%)	90	88	88	88	85
RH Min (%)	72	64	63	48	51
Wind Speed (kmph)	11	4	4	4	6
Wind Direction (deg)	289	246	255	281	310

Agromet Advisory

General advisory:

- Resume spraying and fertilization in dry days based upon now-cast warning (before 2-3 hrs).
- By using recent rainfall and soil moisture start field preparation for non-paddy crops like pulse, oilseeds and vegetables in upland condition and take up early sowing/transplanting.
- Before sowing of any crop, farmers are advised for application of well decomposed FYM or compost, it increases the water use efficiency as well as nutrient quantity in the soil. Farmers are also advised to increase the use of potash, it helps in minimizing the losses due to the moisture stress during dry spell

Kharif Rice:

The crop is at tillering to active tillering stage. **Maintain about 2-3 cm standing water during active tillering stage.** At 21 days after transplanting apply 50% N (35 kg Urea/acre) .**Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.**

Weather based Disease and pest

- **In early transplanted/direct sown crop, chance for leaf folder infestation due to the prevalence of cloudy weather. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days.**
- **Due to forecasted increase in temperature there may be increase in infestation of thrips, to control this Spray Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.**
- **Due to high temperature and high humidity there will be chances of infestation of bacterial leaf streak, to control apply Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.**

Maize:

Past week high humidity (>90%) and decrease in temperature (21-33°C) and continuous rain the past week may aggravate **downy mildew disease** in Maize. High humid conditions, temperature between 18-27 Deg. Celsius favors the occurrence of **leaf blight** in Maize. To control both disease spray Azoxystrobin + Mancozeb @ 300 g /ac. Due to water logging condition **stem rot** may be found, then farmers should apply carbendazim @ 2g/ltr water. Due to forecasted hot and humid weather, there may be **Aphids and Thrips** attack, then apply Dimethoate 30EC @ 400ml/Acre.

VEGETABLES:

- Due to past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- **Brinjal, tomato and chilli seedlings** should be planted in the main field. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Past weather may aggravate **fruit rot in pointed gourd**, to control it remove damaged fruit from the field and drain out excess water. Spray alternately Copper Oxychloride 50% WP (4 g/litre of water) and Metalaxyl 8% + Mancozeb 64% (3 g/litre of water).
- Past rainy weather is favorable for **Root and stem rot in cucurbits**. To control farmers may apply Thiophanate Methyl 70% WP @ 2 gm/l or Azoxystrobin 18.2 SC + Difeconazole 11.11% SC 1.5 gm/l to be applied base as well as root zone.
- At present chilli crop at nursery to transplanting stage. Past weather conditions are favourable for **damping off in chilli nurseries**. To control it by drenching of copper oxy chloride @ 3 g per litre of water (or) 1 % Bordeaux mixture, two times at an interval of 7 days. Farmers are advised to dip the seedlings in 5 ml imidachloprid and 10 g carbendazim solution per 10 litres of water for 5 minutes while transplanting.

ANIMAL HUSBANDRY:

Keep animal shed and shelter house free from water logging by keeping it dry and free from weed. Fresh and immature leaves should not to be fed more amount as they may cause diarrhea. If grasses were cut in morning or evening time, it can be sun dried for about 30 minute before feeding.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-647
Date- 01/09/2026

AGROMET ADVISORY BULLETIN FOR BADAMBA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 05 mm and district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.28°C (-0.72) and 23.57°C (-0.63), respectively. The mean relative humidity was 88-92%.

Forecasted period: Today(8.30IST) to 06th September (8.30IST)2026

Weather advisory *Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block.* The day temperature will be 31-33°C and night temperature will be 24-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 82-87% and 46-68% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Badamba (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	01/09/2026	02/09/2026	03/09/2026	04/09/2026	05/09/2026
Rainfall(mm)	12	9	8	8	7
T-Max(°C)	32	31	32	32	33
T-Min(°C)	25	24	25	25	26
Cloud Cover(octa)	6	6	6	5	5
RH Max (%)	87	82	85	82	82
RH Min (%)	68	58	59	53	46
Wind Speed (Kmph)	13	5	5	4	4
Wind Direction (deg)	283	228	238	259	312

Agromet Advisory

General advisory:

- Resume spraying and fertilization in dry days based upon now-cast warning (before 2-3 hrs).
- By using recent rainfall and soil moisture start field preparation for non-paddy crops like pulse, oilseeds and vegetables in upland condition and take up early sowing/transplanting.
- Before sowing of any crop, farmers are advised for application of well decomposed FYM or compost, it increases the water use efficiency as well as nutrient quantity in the soil. Farmers are also advised to increase the use of potash, it helps in minimizing the losses due to the moisture stress during dry spell

Kharif Rice:

The crop is at tillering to active tillering stage. Maintain about 2-3 cm standing water during active tillering stage. At 21 days after transplanting apply 50% N (35 kg Urea/acre). Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.

Weather based Disease and pest

- In early transplanted/direct sown crop, chance for **leaf folder** infestation due to the prevalence of cloudy weather. If observe in field then spray **Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days.**
- Due to forecasted increase in temperature there may be increase in infestation of **thrips**, to control this Spray **Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.**
- Due to high temperature and high humidity there will be chances of infestation of **bacterial leaf streak**, to control apply **Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.**

Maize:

Past week high humidity (>90%) and decrease in temperature (21-33°C) and continuous rain the past week may aggravate **downy mildew disease** in Maize. High humid conditions, temperature between 18-27 Deg. Celsius favors the occurrence of **leaf blight** in Maize. To control both disease spray **Azoxystrobin + Mancozeb @ 300 g /ac.** Due to water logging condition **stem rot** may be found, then farmers should apply **carbendazim @ 2g/ltr water.** Due to forecasted hot and humid weather, there may be **Aphids and Thrips** attack, then apply **Dimethoate 30EC @ 400ml/Acre.**

VEGETABLES:

- Due to past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray **Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water.** If necessary spray second and third round at 15 days interval.
- **Brinjal, tomato and chilli** seedlings should be planted in the main field. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Past weather may aggravate **fruit rot in pointed gourd**, to control it remove damaged fruit from the field and drain out excess water. Spray alternately **Copper Oxychloride 50% WP (4 g/litre of water) and Metalaxyl 8% + Mancozeb 64% (3 g/litre of water).**
- Past rainy weather is favorable for **Root and stem rot in cucurbits.** To control farmers may apply **Thiophanate Methyl 70% WP @ 2 gm/l or Azoxystrobin 18.2 SC + Difeconazole 11.11% SC 1.5 gm/l** to be applied base as well as root zone.
- At present chilli crop at nursery to transplanting stage. Past weather conditions are favourable for **damping off in chilli nurseries.** To control it by drenching of copper oxy chloride @ 3 g per litre of water (or) 1 % Bordeaux mixture, two times at an interval of 7 days. Farmers are advised to dip the seedlings in 5 ml imidachloprid and 10 g carbendazim solution per 10 litres of water for 5 minutes while transplanting.

ANIMAL HUSBANDRY:

Keep animal shed and shelter house free from water logging by keeping it dry and free from weed. Fresh and immature leaves should not to be fed more amount as they may cause diarrhea. If grasses were cut in morning or evening time, it can be sun dried for about 30 minute before feeding.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-647
Date- 01/09/2026

AGROMET ADVISORY BULLETIN FOR BANKI

District Past week average weather condition

The district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.28°C (-0.72) and 23.57°C (-0.63), respectively. The mean relative humidity was 88-92%.

Forecasted period: Today(8.30IST) to 06th September (8.30IST)2026

Weather advisory: *Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block.* The day temperature will be 31-33°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 85-92% and 49-74% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Banki (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	01/09/2026	02/09/2026	03/09/2026	04/09/2026	05/09/2026
Rainfall(mm)	11	9	8	8	7
T-Max(°C)	32	31	32	33	33
T-Min(°C)	25	25	25	26	26
Cloud Cover(octa)	6	6	6	5	5
RH Max (%)	92	86	88	87	85
RH Min (%)	74	63	65	49	51
Wind Speed (Kmph)	10	5	4	4	4
Wind Direction (deg)	282	234	248	270	311

Agromet Advisory

General advisory:

- Resume spraying and fertilization in dry days based upon now-cast warning (before 2-3 hrs).
- By using recent rainfall and soil moisture start field preparation for non-paddy crops like pulse, oilseeds and vegetables in upland condition and take up early sowing/transplanting.
- Before sowing of any crop, farmers are advised for application of well decomposed FYM or compost, it increases the water use efficiency as well as nutrient quantity in the soil. Farmers are also advised to increase the use of potash, it helps in minimizing the losses due to the moisture stress during dry spell

Kharif Rice:

The crop is at tillering to active tillering stage. Maintain about 2-3 cm standing water during active tillering stage. At 21 days after transplanting apply 50% N (35 kg Urea/acre). Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.

Weather based Disease and pest

- In early transplanted/direct sown crop, chance for leaf folder infestation due to the prevalence of cloudy weather. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days.
- Due to forecasted increase in temperature there may be increase in infestation of thrips, to control this Spray Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.
- Due to high temperature and high humidity there will be chances of infestation of bacterial leaf streak, to control apply Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.

Maize:

Past week high humidity (>90%) and decrease in temperature (21-33°C) and continuous rain the past week may aggravate downy mildew disease in Maize. High humid conditions, temperature between 18-27 Deg. Celsius favors the occurrence of leaf blight in Maize. To control both disease spray Azoxystrobin + Mancozeb @ 300 g /ac. Due to water logging condition stem rot may be found, then farmers should apply carbendazim @ 2g/ltr water. Due to forecasted hot and humid weather, there may be Aphids and Thrips attack, then apply Dimethoate 30EC @ 400ml/Acre.

VEGETABLES:

- Due to past rain, Chances of fruit rot infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Past weather may aggravate fruit rot in pointed gourd, to control it remove damaged fruit from the field and drain out excess water. Spray alternately Copper Oxychloride 50% WP (4 g/litre of water) and Metalaxyl 8% + Mancozeb 64% (3 g/litre of water).
- Past rainy weather is favorable for Root and stem rot in cucurbits. To control farmers may apply Thiophanate Methyl 70% WP @ 2 gm/l or Azoxystrobin 18.2 SC + Difeconazole 11.11% SC 1.5 gm/l to be applied base as well as root zone.
- At present chilli crop at nursery to transplanting stage. Past weather conditions are favourable for damping off in chilli nurseries. To control it by drenching of copper oxy chloride @ 3 g per litre of water (or) 1 % Bordeaux mixture, two times at an interval of 7 days. Farmers are advised to dip the seedlings in 5 ml imidachloprid and 10 g carbendazim solution per 10 litres of water for 5 minutes while transplanting.

ANIMAL HUSBANDRY:

Keep animal shed and shelter house free from water logging by keeping it dry and free from weed. Fresh and immature leaves should not to be fed more amount as they may cause diarrhea. If grasses were cut in morning or evening time, it can be sun dried for about 30 minute before feeding.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-647
Date- 01/09/2026

AGROMET ADVISORY BULLETIN FOR BANKI-DAMPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 06 mm and district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.28°C (-0.72) and 23.57°C (-0.63), respectively. The mean relative humidity was 88-92%.

Forecasted period: Today(8.30IST) to 06th September (8.30IST)2026

Weather Advisory:: *Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block.* The day temperature will be 32-33°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 85-92% and 49-74% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Banki-Dampada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	01/09/2026	02/09/2026	03/09/2026	04/09/2026	05/09/2026
Rainfall(mm)	11	9	7	8	7
T-Max(°C)	32	32	32	33	33
T-Min(°C)	25	25	25	26	26
Cloud Cover(octa)	6	6	6	5	5
RH Max (%)	92	86	88	87	85
RH Min (%)	74	63	65	49	51
Wind Speed (Kmph)	10	5	4	4	4
Wind Direction (deg)	282	234	248	270	311

Agromet Advisory

General advisory:

- Resume spraying and fertilization in dry days based upon now-cast warning (before 2-3 hrs).
- By using recent rainfall and soil moisture start field preparation for non-paddy crops like pulse, oilseeds and vegetables in upland condition and take up early sowing/transplanting.
- Before sowing of any crop, farmers are advised for application of well decomposed FYM or compost, it increases the water use efficiency as well as nutrient quantity in the soil. Farmers are also advised to increase the use of potash, it helps in minimizing the losses due to the moisture stress during dry spell

Kharif Rice:

The crop is at tillering to active tillering stage. Maintain about 2-3 cm standing water during active tillering stage. At 21 days after transplanting apply 50% N (35 kg Urea/acre). Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.

Weather based Disease and pest

- In early transplanted/direct sown crop, chance for **leaf folder** infestation due to the prevalence of cloudy weather. If observe in field then spray **Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days.**
- Due to forecasted increase in temperature there may be increase in infestation of **thrips**, to control this Spray **Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.**
- Due to high temperature and high humidity there will be chances of infestation of **bacterial leaf streak**, to control apply **Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.**

Maize:

Past week high humidity (>90%) and decrease in temperature (21-33°C) and continuous rain the past week may aggravate **downy mildew disease** in Maize. High humid conditions, temperature between 18-27 Deg. Celsius favors the occurrence of **leaf blight** in Maize. To control both disease spray **Azoxystrobin + Mancozeb @ 300 g /ac.** Due to water logging condition **stem rot** may be found, then farmers should apply **carbendazim @ 2g/ltr water.** Due to forecasted hot and humid weather, there may be **Aphids and Thrips** attack, then apply **Dimethoate 30EC @ 400ml/Acre.**

VEGETABLES:

- Due to past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray **Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water.** If necessary spray second and third round at 15 days interval.
- **Brinjal, tomato and chilli** seedlings should be planted in the main field. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Past weather may aggravate **fruit rot in pointed gourd**, to control it remove damaged fruit from the field and drain out excess water. Spray alternately **Copper Oxychloride 50% WP (4 g/litre of water) and Metalaxyl 8% + Mancozeb 64% (3 g/litre of water).**
- Past rainy weather is favorable for **Root and stem rot in cucurbits.** To control farmers may apply **Thiophanate Methyl 70% WP @ 2 gm/l or Azoxystrobin 18.2 SC + Difeconazole 11.11% SC 1.5 gm/l** to be applied base as well as root zone.
- At present chilli crop at nursery to transplanting stage. Past weather conditions are favourable for **damping off in chilli nurseries.** To control it by drenching of copper oxy chloride @ 3 g per litre of water (or) 1 % Bordeaux mixture, two times at an interval of 7 days. Farmers are advised to dip the seedlings in 5 ml imidachloprid and 10 g carbendazim solution per 10 litres of water for 5 minutes while transplanting.

ANIMAL HUSBANDRY:

Keep animal shed and shelter house free from water logging by keeping it dry and free from weed. Fresh and immature leaves should not to be fed more amount as they may cause diarrhea. If grasses were cut in morning or evening time, it can be sun dried for about 30 minute before feeding.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-647
Date- 01/09/2026

AGROMET ADVISORY BULLETIN FOR BARANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 09 mm and district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.28°C (-0.72) and 23.57°C (-0.63), respectively. The mean relative humidity was 88-92%.

Forecasted period: Today(8.30IST) to 06th September (8.30IST)2026

Weatheradvisory: *Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block.* The day temperature will be 31-33°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 81-88% and 42-65% respectively. There will be partly cloudy to overcast sky for next 5days.

Block	Baranga(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	01/09/2026	02/09/2026	03/09/2026	04/09/2026	05/09/2026
Rainfall(mm)	10	8	8	8	7
T-Max(°C)	32	31	32	32	33
T-Min(°C)	25	25	25	26	26
Cloud Cover	6	6	6	5	5
RH Max (%)	88	81	85	82	81
RH Min (%)	65	60	53	45	42
Wind Speed (Kmph)	12	7	5	5	4
Wind Direction (deg)	280	227	241	295	275

Agromet Advisory

General advisory:

- Resume spraying and fertilization in dry days based upon now-cast warning (before 2-3 hrs).
- By using recent rainfall and soil moisture start field preparation for non-paddy crops like pulse, oilseeds and vegetables in upland condition and take up early sowing/transplanting.
- Before sowing of any crop, farmers are advised for application of well decomposed FYM or compost, it increases the water use efficiency as well as nutrient quantity in the soil. Farmers are also advised to increase the use of potash, it helps in minimizing the losses due to the moisture stress during dry spell

Kharif Rice:

The crop is at tillering to active tillering stage. Maintain about 2-3 cm standing water during active tillering stage. At 21 days after transplanting apply 50% N (35 kg Urea/acre) .Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.

Weather based Disease and pest

- *In early transplanted/direct sown crop, chance for leaf folder infestation due to the prevalence of cloudy weather. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days.*
- Due to forecasted increase in temperature there may be increase in infestation of thrips, to control this Spray Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.
- Due to high temperature and high humidity there will be chances of infestation of bacterial leaf streak, to control apply Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.

Maize:

Past week high humidity (>90%) and decrease in temperature (21-33°C) and continuous rain the past week may aggravate downy mildew disease in Maize. High humid conditions, temperature between 18-27 Deg. Celsius favors the occurrence of leaf blight in Maize. To control both disease spray Azoxystrobin + Mancozeb @ 300 g /ac.Due to water logging condition stem rot may be found, then farmers should apply carbendazim @ 2g/ltr water.Due to forecasted hot and humid weather, there may be Aphids and Thrips attack,then applyDimethoate 30EC @ 400ml/Acre.

VEGETABLES:

- Due to past rain, Chances of fruit rot infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilliseedlings should be planted in the main field. To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Past weather may aggravate fruit rot in pointed gourd, to control it remove damaged fruit from the field and drain out excess water. Spray alternately Copper Oxychloride 50% WP (4 g/litre of water) and Metalaxyl 8% + Mancozeb 64% (3 g/litre of water).
- Past rainy weather is favorable for Root and stem rot in cucurbits. To control farmers may apply Thiophanate Methyl 70% WP @ 2 gm/l or Azoxystrobin 18.2 SC+ Difeconazole 11.11% SC 1.5 gm/l to be applied base as well as root zone.
- At present chilli crop at nursery to transplanting stage. Past weather conditions are favourable for damping off in chilli nurseries. To control it by drenching of copper oxy chloride @ 3 g per litre of water (or) 1 % Bordeaux mixture, two times at an interval of 7 days. Farmers are advised to dip the seedlings in 5 ml imidachloprid and 10 g carbendazim solution per 10 litres of water for 5 minutes while transplanting.

ANIMAL HUSBANDRY:

Keep animal shed and shelter house free from water logging by keeping it dry and free from weed. Fresh and immature leaves should not to be fed more amount as they may cause diarrhea. If grasses were cut in morning or evening time, it can be sun dried for about 30 minute before feeding.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-647
Date- 01/09/2026

AGROMET ADVISORY BULLETIN FOR CUTTACK SADAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 06 mm and district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.28°C (-0.72) and 23.57°C (-0.63), respectively. The mean relative humidity was 88-92%.

Forecasted period: Today(8.30IST) to 06th September (8.30IST)2026

Weather advisory: *Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block.* The day temperature will be 31-32°C and night temperature will be 24-25°C for upcoming 5 days. Maximum and minimum relative humidity will be 83-89% and 45-69% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Cuttack sadar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	01/09/2026	02/09/2026	03/09/2026	04/09/2026	05/09/2026
Rainfall(mm)	10	8	9	7	7
T-Max(°C)	32	31	31	32	32
T-Min(°C)	25	24	24	25	25
Cloud Cover(octa)	6	6	6	5	5
RH Max (%)	89	86	87	85	83
RH Min (%)	69	63	58	45	49
Wind Speed (Kmph)	14	6	6	5	6
Wind Direction (deg)	282	243	259	278	284

Agromet Advisory

General advisory:

- Resume spraying and fertilization in dry days based upon now-cast warning (before 2-3 hrs).
- By using recent rainfall and soil moisture start field preparation for non-paddy crops like pulse, oilseeds and vegetables in upland condition and take up early sowing/transplanting.
- Before sowing of any crop, farmers are advised for application of well decomposed FYM or compost, it increases the water use efficiency as well as nutrient quantity in the soil. Farmers are also advised to increase the use of potash, it helps in minimizing the losses due to the moisture stress during dry spell

Kharif Rice:

The crop is at tillering to active tillering stage. Maintain about 2-3 cm standing water during active tillering stage. At 21 days after transplanting apply 50% N (35 kg Urea/acre). Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.

Weather based Disease and pest

- In early transplanted/direct sown crop, chance for **leaf folder** infestation due to the prevalence of cloudy weather. If observe in field then spray **Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days.**
- Due to forecasted increase in temperature there may be increase in infestation of **thrips**, to control this Spray **Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.**
- Due to high temperature and high humidity there will be chances of infestation of **bacterial leaf streak**, to control apply **Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.**

Sugarcane:

Wrapping and propping should be done when crop are 4-5 month old in medium and late duration varieties. For **early shoot borer** Spray Chloropyrifos @ 400 ml/ac in 200 lts of water after cessation of rain. For **Red rot**, avoid rationing in red rot affected area; spray the crop with Mancozeb 0.3% or carbendazim 0.1% at 15 days interval. Complete hoeing, weeding and apply fertilizer urea 100kg with 34 kg MOP in lines followed by earthing up during non rainy period.

VEGETABLES:

- Due to past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- **Brinjal, tomato and chilli** seedlings should be planted in the main field. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Past weather may aggravate **fruit rot in pointed gourd**, to control it remove damaged fruit from the field and drain out excess water. Spray alternately Copper Oxychloride 50% WP (4 g/litre of water) and Metalaxyl 8% + Mancozeb 64% (3 g/litre of water).
- Past rainy weather is favorable for **Root and stem rot in cucurbits**. To control farmers may apply Thiophanate Methyl 70% WP @ 2 gm/l or Azoxistrobin 18.2 SC + Difeconazole 11.11% SC 1.5 gm/l to be applied base as well as root zone.
- At present chilli crop at nursery to transplanting stage. Past weather conditions are favourable for **damping off in chilli nurseries**. To control it by drenching of copper oxy chloride @ 3 g per litre of water (or) 1 % Bordeaux mixture, two times at an interval of 7 days. Farmers are advised to dip the seedlings in 5 ml imidachloprid and 10 g carbendazim solution per 10 litres of water for 5 minutes while transplanting.

ANIMAL HUSBANDRY:

Keep animal shed and shelter house free from water logging by keeping it dry and free from weed. Fresh and immature leaves should not to be fed more amount as they may cause diarrhea. If grasses were cut in morning or evening time, it can be sun dried for about 30 minute before feeding.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-647
Date- 01/09/2026

AGROMET ADVISORY BULLETIN FOR KANTAPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 09 mm and district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.28°C (-0.72) and 23.57°C (-0.63), respectively. The mean relative humidity was 88-92%.

Forecasted period: Today(8.30IST) to 06th September (8.30IST)2026

Weather advisory: *Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block.* The day temperature will be 32-33°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 81-89% and 45-68% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Kantapada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	01/09/2026	02/09/2026	03/09/2026	04/09/2026	05/09/2026
Rainfall(mm)	10	8	8	7	7
T-Max(°C)	32	32	32	33	33
T-Min(°C)	25	25	25	26	26
Cloud Cover(octa)	6	6	6	5	5
RH Max (%)	89	85	86	85	81
RH Min (%)	68	62	56	45	50
Wind Speed (Kmph)	11	9	6	6	5
Wind Direction (deg)	266	242	247	297	291

Agromet Advisory

General advisory:

- Resume spraying and fertilization in dry days based upon now-cast warning (before 2-3 hrs).
- By using recent rainfall and soil moisture start field preparation for non-paddy crops like pulse, oilseeds and vegetables in upland condition and take up early sowing/transplanting.
- Before sowing of any crop, farmers are advised for application of well decomposed FYM or compost, it increases the water use efficiency as well as nutrient quantity in the soil. Farmers are also advised to increase the use of potash, it helps in minimizing the losses due to the moisture stress during dry spell

Kharif Rice:

The crop is at tillering to active tillering stage. Maintain about 2-3 cm standing water during active tillering stage. At 21 days after transplanting apply 50% N (35 kg Urea/acre). Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.

Weather based Disease and pest

- In early transplanted/direct sown crop, chance for leaf folder infestation due to the prevalence of cloudy weather. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days.
- Due to forecasted increase in temperature there may be increase in infestation of thrips, to control this Spray Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.
- Due to high temperature and high humidity there will be chances of infestation of bacterial leaf streak, to control apply Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.

Sugarcane:

Wrapping and propping should be done when crop are 4-5 month old in medium and late duration varieties. For early shoot borer Spray Chloropyrifos @ 400 ml/ac in 200 lts of water after cessation of rain. For Red rot, avoid rationing in red rot affected area; spray the crop with Mancozeb 0.3% or carbendazim 0.1% at 15 days interval. Complete hoeing, weeding and apply fertilizer urea 100kg with 34 kg MOP in lines followed by earthing up during non rainy period.

VEGETABLES:

- Due to past rain, Chances of fruit rot infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Past weather may aggravate fruit rot in pointed gourd, to control it remove damaged fruit from the field and drain out excess water. Spray alternately Copper Oxychloride 50% WP (4 g/litre of water) and Metalaxyl 8% + Mancozeb 64% (3 g/litre of water).
- Past rainy weather is favorable for Root and stem rot in cucurbits. To control farmers may apply Thiophanate Methyl 70% WP @ 2 gm/l or Azoxistrobin 18.2 SC + Difeconazole 11.11% SC 1.5 gm/l to be applied base as well as root zone.
- At present chilli crop at nursery to transplanting stage. Past weather conditions are favourable for damping off in chilli nurseries. To control it by drenching of copper oxy chloride @ 3 g per litre of water (or) 1 % Bordeaux mixture, two times at an interval of 7 days. Farmers are advised to dip the seedlings in 5 ml imidachloprid and 10 g carbendazim solution per 10 litres of water for 5 minutes while transplanting.

ANIMAL HUSBANDRY:

Keep animal shed and shelter house free from water logging by keeping it dry and free from weed. Fresh and immature leaves should not to be fed more amount as they may cause diarrhea. If grasses were cut in morning or evening time, it can be sun dried for about 30 minute before feeding.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-647
Date- 01/09/2026

AGROMET ADVISORY BULLETIN FOR MAHANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 07 mm and district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.28°C (-0.72) and 23.57°C (-0.63), respectively. The mean relative humidity was 88-92%.

Forecasted period: Today(8.30IST) to 06th September (8.30IST)2026

Weather advisory: *Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block.* The day temperature will be and it will be 31-32°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 83-88% and 44-68 respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Mahanga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	01/09/2026	02/09/2026	03/09/2026	04/09/2026	05/09/2026
Rainfall(mm)	12	9	8	8	7
T-Max(°C)	32	31	32	32	32
T-Min(°C)	25	25	25	26	26
Cloud Cover(octal)	6	6	6	6	6
RH Max (%)	88	87	88	85	83
RH Min (%)	68	64	55	44	51
Wind Speed (Kmph)	17	9	6	6	6
Wind Direction (deg)	277	241	246	277	266

Agromet Advisory

General advisory:

- Resume spraying and fertilization in dry days based upon now-cast warning (before 2-3 hrs).
- By using recent rainfall and soil moisture start field preparation for non-paddy crops like pulse, oilseeds and vegetables in upland condition and take up early sowing/transplanting.
- Before sowing of any crop, farmers are advised for application of well decomposed FYM or compost, it increases the water use efficiency as well as nutrient quantity in the soil. Farmers are also advised to increase the use of potash, it helps in minimizing the losses due to the moisture stress during dry spell

Kharif Rice:

The crop is at tillering to active tillering stage. Maintain about 2-3 cm standing water during active tillering stage. At 21 days after transplanting apply 50% N (35 kg Urea/acre). Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.

Weather based Disease and pest

- In early transplanted/direct sown crop, chance for leaf folder infestation due to the prevalence of cloudy weather. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days.
- Due to forecasted increase in temperature there may be increase in infestation of thrips, to control this Spray Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.
- Due to high temperature and high humidity there will be chances of infestation of bacterial leaf streak, to control apply Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.

Sugarcane:

Wrapping and propping should be done when crop are 4-5 month old in medium and late duration varieties. For early shoot borer Spray Chloropyrifos @ 400 ml/ac in 200 lts of water after cessation of rain. For Red rot, avoid rationing in red rot affected area; spray the crop with Mancozeb 0.3% or carbendazim 0.1% at 15 days interval. Complete hoeing, weeding and apply fertilizer urea 100kg with 34 kg MOP in lines followed by earthing up during non rainy period.

VEGETABLES:

- Due to past rain, Chances of fruit rot infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Past weather may aggravate fruit rot in pointed gourd, to control it remove damaged fruit from the field and drain out excess water. Spray alternately Copper Oxychloride 50% WP (4 g/litre of water) and Metalaxyl 8% + Mancozeb 64% (3 g/litre of water).
- Past rainy weather is favorable for Root and stem rot in cucurbits. To control farmers may apply Thiophanate Methyl 70% WP @ 2 gm/l or Azoxistrobin 18.2 SC + Difeconazole 11.11% SC 1.5 gm/l to be applied base as well as root zone.
- At present chilli crop at nursery to transplanting stage. Past weather conditions are favourable for damping off in chilli nurseries. To control it by drenching of copper oxy chloride @ 3 g per litre of water (or) 1 % Bordeaux mixture, two times at an interval of 7 days. Farmers are advised to dip the seedlings in 5 ml imidachloprid and 10 g carbendazim solution per 10 litres of water for 5 minutes while transplanting.

ANIMAL HUSBANDRY:

Keep animal shed and shelter house free from water logging by keeping it dry and free from weed. Fresh and immature leaves should not to be fed more amount as they may cause diarrhea. If grasses were cut in morning or evening time, it can be sun dried for about 30 minute before feeding.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-647
Date- 01/09/2026

AGROMET ADVISORY BULLETIN FOR NARASINGHPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 04 mm and district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.28°C (-0.72) and 23.57°C (-0.63), respectively. The mean relative humidity was 88-92%.

Forecasted period: Today(8.30IST) to 06th September (8.30IST)2026

Weather advisory: *Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block.* The day temperature will be 31-32°C and night temperature will be 24-25°C for upcoming 5 days. Maximum and minimum relative humidity will be 90-95% and 57-77% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Narasinghpur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	01/09/2026	02/09/2026	03/09/2026	04/09/2026	05/09/2026
Rainfall(mm)	14	9	8	8	8
T-Max(°C)	31	31	31	32	32
T-Min(°C)	25	24	25	25	25
Cloud Cover(/octa)	6	6	6	5	5
RH Max (%)	95	92	92	90	91
RH Min (%)	77	67	70	57	57
Wind Speed (Kmph)	8	3	4	3	4
Wind Direction (deg)	290	243	233	243	297

Agromet Advisory

General advisory:

- Resume spraying and fertilization in dry days based upon now-cast warning (before 2-3 hrs).
- By using recent rainfall and soil moisture start field preparation for non-paddy crops like pulse, oilseeds and vegetables in upland condition and take up early sowing/transplanting.
- Before sowing of any crop, farmers are advised for application of well decomposed FYM or compost, it increases the water use efficiency as well as nutrient quantity in the soil. Farmers are also advised to increase the use of potash, it helps in minimizing the losses due to the moisture stress during dry spell

Kharif Rice:

The crop is at tillering to active tillering stage. Maintain about 2-3 cm standing water during active tillering stage. At 21 days after transplanting apply 50% N (35 kg Urea/acre). Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.

Weather based Disease and pest

- In early transplanted/direct sown crop, chance for **leaf folder** infestation due to the prevalence of cloudy weather. If observe in field then spray **Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days.**
- Due to forecasted increase in temperature there may be increase in infestation of **thrips**, to control this Spray **Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.**
- Due to high temperature and high humidity there will be chances of infestation of **bacterial leaf streak**, to control apply **Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.**

Maize:

Past week high humidity (>90%) and decrease in temperature (21-33°C) and continuous rain the past week may aggravate **downy mildew disease** in Maize. High humid conditions, temperature between 18-27 Deg. Celsius favors the occurrence of **leaf blight** in Maize. To control both disease spray **Azoxystrobin + Mancozeb @ 300 g /ac.** Due to water logging condition **stem rot** may be found, then farmers should apply **carbendazim @ 2g/ltr water.** Due to forecasted hot and humid weather, there may be **Aphids and Thrips** attack, then apply **Dimethoate 30EC @ 400ml/Acre.**

VEGETABLES:

- Due to past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray **Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water.** If necessary spray second and third round at 15 days interval.
- **Brinjal, tomato and chilli** seedlings should be planted in the main field. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Past weather may aggravate **fruit rot in pointed gourd**, to control it remove damaged fruit from the field and drain out excess water. Spray alternately **Copper Oxychloride 50% WP (4 g/litre of water) and Metalaxyl 8% + Mancozeb 64% (3 g/litre of water).**
- Past rainy weather is favorable for **Root and stem rot in cucurbits.** To control farmers may apply **Thiophanate Methyl 70% WP @ 2 gm/l or Azoxystrobin 18.2 SC + Difeconazole 11.11% SC 1.5 gm/l** to be applied base as well as root zone.
- At present chilli crop at nursery to transplanting stage. Past weather conditions are favourable for **damping off in chilli nurseries.** To control it by drenching of copper oxy chloride @ 3 g per litre of water (or) 1 % Bordeaux mixture, two times at an interval of 7 days. Farmers are advised to dip the seedlings in 5 ml imidachloprid and 10 g carbendazim solution per 10 litres of water for 5 minutes while transplanting.

ANIMAL HUSBANDRY:

Keep animal shed and shelter house free from water logging by keeping it dry and free from weed. Fresh and immature leaves should not to be fed more amount as they may cause diarrhea. If grasses were cut in morning or evening time, it can be sun dried for about 30 minute before feeding.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-647
Date- 01/09/2026

AGROMET ADVISORY BULLETIN FOR NIALI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 09 mm and district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.28°C (-0.72) and 23.57°C (-0.63), respectively. The mean relative humidity was 88-92%.

Forecasted period: Today(8.30IST) to 06th September (8.30IST)2026

Weather advisory: *Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block.* The day temperature will be 32-33°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 78-88% and 42-62% respectively.

There will be partly cloudy to overcast sky for next 5 days.

Block	Niali (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	01/09/2026	02/09/2026	03/09/2026	04/09/2026	05/09/2026
Rainfall(mm)	10	8	7	7	7
T-Max(°C)	33	32	32	33	33
T-Min(°C)	26	25	25	26	26
Cloud Cover(octa)	6	5	5	6	5
RH Max (%)	88	78	82	81	81
RH Min (%)	62	57	47	42	46
Wind Speed(Kmph)	11	10	5	6	7
Wind Direction(deg)	264	232	249	305	238

Agromet Advisory

General advisory:

- Resume spraying and fertilization in dry days based upon now-cast warning (before 2-3 hrs).
- By using recent rainfall and soil moisture start field preparation for non-paddy crops like pulse, oilseeds and vegetables in upland condition and take up early sowing/transplanting.
- Before sowing of any crop, farmers are advised for application of well decomposed FYM or compost, it increases the water use efficiency as well as nutrient quantity in the soil. Farmers are also advised to increase the use of potash, it helps in minimizing the losses due to the moisture stress during dry spell

Kharif Rice:

The crop is at tillering to active tillering stage. Maintain about 2-3 cm standing water during active tillering stage. At 21 days after transplanting apply 50% N (35 kg Urea/acre). Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.

Weather based Disease and pest

- In early transplanted/direct sown crop, chance for leaf folder infestation due to the prevalence of cloudy weather. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days.
- Due to forecasted increase in temperature there may be increase in infestation of thrips, to control this Spray Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.
- Due to high temperature and high humidity there will be chances of infestation of bacterial leaf streak, to control apply Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.

Sugarcane:

Wrapping and propping should be done when crop are 4-5 month old in medium and late duration varieties. For early shoot borer Spray Chloropyrifos @ 400 ml/ac in 200 lts of water after cessation of rain. For Red rot, avoid rationing in red rot affected area; spray the crop with Mancozeb 0.3% or carbendazim 0.1% at 15 days interval. Complete hoeing, weeding and apply fertilizer urea 100kg with 34 kg MOP in lines followed by earthing up during non rainy period.

VEGETABLES:

- Due to past rain, Chances of fruit rot infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Past weather may aggravate fruit rot in pointed gourd, to control it remove damaged fruit from the field and drain out excess water. Spray alternately Copper Oxychloride 50% WP (4 g/litre of water) and Metalaxyl 8% + Mancozeb 64% (3 g/litre of water).
- Past rainy weather is favorable for Root and stem rot in cucurbits. To control farmers may apply Thiophanate Methyl 70% WP @ 2 gm/l or Azoxistrobin 18.2 SC + Difeconazole 11.11% SC 1.5 gm/l to be applied base as well as root zone.
- At present chilli crop at nursery to transplanting stage. Past weather conditions are favourable for damping off in chilli nurseries. To control it by drenching of copper oxy chloride @ 3 g per litre of water (or) 1 % Bordeaux mixture, two times at an interval of 7 days. Farmers are advised to dip the seedlings in 5 ml imidachloprid and 10 g carbendazim solution per 10 litres of water for 5 minutes while transplanting.

ANIMAL HUSBANDRY:

Keep animal shed and shelter house free from water logging by keeping it dry and free from weed. Fresh and immature leaves should not to be fed more amount as they may cause diarrhea. If grasses were cut in morning or evening time, it can be sun dried for about 30 minute before feeding.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-647
Date- 01/09/2026

AGROMET ADVISORY BULLETIN FOR NISCHINTAKOILI
District Past week average weather condition and Block past week average Rainfall condition

The block received average 12 mm and district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.28°C (-0.72) and 23.57°C (-0.63), respectively. The mean relative humidity was 88-92%.

Forecasted period: Today(8.30IST) to 06th September (8.30IST)2026

Weatheradvisory: Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block. The day temperature will be 31-32°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 83-88% and 44-68% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Nischintakoili (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	01/09/2026	02/09/2026	03/09/2026	04/09/2026	05/09/2026
Rainfall(mm)	11	9	8	7	7
T-Max(°C)	32	31	32	32	32
T-Min(°C)	25	25	25	26	26
Cloud Cover(octa)	6	6	6	6	5
RH Max (%)	88	87	88	85	83
RH Min (%)	68	64	55	44	51
Wind Speed (Kmph)	15	9	6	6	6
Wind Direction (deg)	275	241	249	277	263

Agromet Advisory

General advisory:

- Resume spraying and fertilization in dry days based upon now-cast warning (before 2-3 hrs).
- By using recent rainfall and soil moisture start field preparation for non-paddy crops like pulse, oilseeds and vegetables in upland condition and take up early sowing/transplanting.
- Before sowing of any crop, farmers are advised for application of well decomposed FYM or compost, it increases the water use efficiency as well as nutrient quantity in the soil. Farmers are also advised to increase the use of potash, it helps in minimizing the losses due to the moisture stress during dry spell

Kharif Rice:

The crop is at tillering to active tillering stage. Maintain about 2-3 cm standing water during active tillering stage. At 21 days after transplanting apply 50% N (35 kg Urea/acre). Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.

Weather based Disease and pest

- In early transplanted/direct sown crop, chance for leaf folder infestation due to the prevalence of cloudy weather. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days.
- Due to forecasted increase in temperature there may be increase in infestation of thrips, to control this Spray Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.
- Due to high temperature and high humidity there will be chances of infestation of bacterial leaf streak, to control apply Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.

Sugarcane:

Wrapping and propping should be done when crop are 4-5 month old in medium and late duration varieties. For early shoot borer Spray Chloropyrifos @ 400 ml/ac in 200 lts of water after cessation of rain. For Red rot, avoid rationing in red rot affected area; spray the crop with Mancozeb 0.3% or carbendazim 0.1% at 15 days interval. Complete hoeing, weeding and apply fertilizer urea 100kg with 34 kg MOP in lines followed by earthing up during non rainy period.

VEGETABLES:

- Due to past rain, Chances of fruit rot infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Past weather may aggravate fruit rot in pointed gourd, to control it remove damaged fruit from the field and drain out excess water. Spray alternately Copper Oxychloride 50% WP (4 g/litre of water) and Metalaxyl 8% + Mancozeb 64% (3 g/litre of water).
- Past rainy weather is favorable for Root and stem rot in cucurbits. To control farmers may apply Thiophanate Methyl 70% WP @ 2 gm/l or Azoxistrobin 18.2 SC + Difeconazole 11.11% SC 1.5 gm/l to be applied base as well as root zone.
- At present chilli crop at nursery to transplanting stage. Past weather conditions are favourable for damping off in chilli nurseries. To control it by drenching of copper oxy chloride @ 3 g per litre of water (or) 1 % Bordeaux mixture, two times at an interval of 7 days. Farmers are advised to dip the seedlings in 5 ml imidachloprid and 10 g carbendazim solution per 10 litres of water for 5 minutes while transplanting.

ANIMAL HUSBANDRY:

Keep animal shed and shelter house free from water logging by keeping it dry and free from weed. Fresh and immature leaves should not to be fed more amount as they may cause diarrhea. If grasses were cut in morning or evening time, it can be sun dried for about 30 minute before feeding.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-647
Date- 01/09/2026

AGROMET ADVISORY BULLETIN FOR SALEPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 06 mm and district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.28°C (-0.72) and 23.57°C (-0.63), respectively. The mean relative humidity was 88-92%.

Forecasted period: Today(8.30IST) to 06th September (8.30IST)2026

Weather advisory: *Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block.* The day temperature will be 31-32°C and night temperature will be 24-25°C for upcoming 5 days. Maximum and minimum relative humidity will be 83-88% and 44-68% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Salepur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	01/09/2026	02/09/2026	03/09/2026	04/09/2026	05/09/2026
Rainfall(mm)	11	8	9	7	7
T-Max(°C)	32	31	31	32	32
T-Min(°C)	25	24	24	25	25
Cloud Cover(octa)	6	6	6	5	5
RH Max (%)	88	87	88	85	83
RH Min (%)	68	64	55	44	51
Wind Speed (Kmph)	15	9	6	6	6
Wind Direction (deg)	277	241	249	281	270

Agromet Advisory

General advisory:

- Resume spraying and fertilization in dry days based upon now-cast warning (before 2-3 hrs).
- By using recent rainfall and soil moisture start field preparation for non-paddy crops like pulse, oilseeds and vegetables in upland condition and take up early sowing/transplanting.
- Before sowing of any crop, farmers are advised for application of well decomposed FYM or compost, it increases the water use efficiency as well as nutrient quantity in the soil. Farmers are also advised to increase the use of potash, it helps in minimizing the losses due to the moisture stress during dry spell

Kharif Rice:

The crop is at tillering to active tillering stage. Maintain about 2-3 cm standing water during active tillering stage. At 21 days after transplanting apply 50% N (35 kg Urea/acre). Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.

Weather based Disease and pest

- In early transplanted/direct sown crop, chance for **leaf folder** infestation due to the prevalence of cloudy weather. If observe in field then spray **Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days.**
- Due to forecasted increase in temperature there may be increase in infestation of **thrips**, to control this Spray **Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.**
- Due to high temperature and high humidity there will be chances of infestation of **bacterial leaf streak**, to control apply **Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.**

Maize:

Past week high humidity (>90%) and decrease in temperature (21-33°C) and continuous rain the past week may aggravate **downy mildew disease** in Maize. High humid conditions, temperature between 18-27 Deg. Celsius favors the occurrence of **leaf blight** in Maize. To control both disease spray **Azoxystrobin + Mancozeb @ 300 g /ac.** Due to water logging condition **stem rot** may be found, then farmers should apply **carbendazim @ 2g/ltr water.** Due to forecasted hot and humid weather, there may be **Aphids and Thrips** attack, then apply **Dimethoate 30EC @ 400ml/Acre.**

VEGETABLES:

- Due to past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray **Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water.** If necessary spray second and third round at 15 days interval.
- **Brinjal, tomato and chilli** seedlings should be planted in the main field. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Past weather may aggravate **fruit rot in pointed gourd**, to control it remove damaged fruit from the field and drain out excess water. Spray alternately **Copper Oxychloride 50% WP (4 g/litre of water) and Metalaxyl 8% + Mancozeb 64% (3 g/litre of water).**
- Past rainy weather is favorable for **Root and stem rot in cucurbits.** To control farmers may apply **Thiophanate Methyl 70% WP @ 2 gm/l or Azoxystrobin 18.2 SC + Difeconazole 11.11% SC 1.5 gm/l** to be applied base as well as root zone.
- At present chilli crop at nursery to transplanting stage. Past weather conditions are favourable for **damping off in chilli nurseries.** To control it by drenching of copper oxy chloride @ 3 g per litre of water (or) 1 % Bordeaux mixture, two times at an interval of 7 days. Farmers are advised to dip the seedlings in 5 ml imidachloprid and 10 g carbendazim solution per 10 litres of water for 5 minutes while transplanting.

ANIMAL HUSBANDRY:

Keep animal shed and shelter house free from water logging by keeping it dry and free from weed. Fresh and immature leaves should not to be fed more amount as they may cause diarrhea. If grasses were cut in morning or evening time, it can be sun dried for about 30 minute before feeding.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-647
Date- 01/09/2026

AGROMET ADVISORY BULLETIN FOR TANGI-CHOUDWAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 09 mm and district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.28°C (-0.72) and 23.57°C (-0.63), respectively. The mean relative humidity was 88-92%.

Forecasted period: Today(8.30IST) to 06th September (8.30IST)2026

Weather advisory: *Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block.* The day temperature will be 31-32°C and night temperature will be 24-25°C for upcoming 5 days. Maximum and minimum relative humidity will be 82-87% and 55-72% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Tangi-Choudwar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	01/09/2026	02/09/2026	03/09/2026	04/09/2026	05/09/2026
Rainfall(mm)	11	9	9	8	7
T-Max(°C)	32	31	31	32	32
T-Min(°C)	25	24	24	25	25
Cloud Cover(octa)	6	6	6	5	5
RH Max (%)	87	84	84	84	82
RH Min (%)	70	64	58	55	59
Wind Speed (Kmph)	17	13	9	9	6
Wind Direction (deg)	265	249	261	287	260

Agromet Advisory

General advisory:

- Resume spraying and fertilization in dry days based upon now-cast warning (before 2-3 hrs).
- By using recent rainfall and soil moisture start field preparation for non-paddy crops like pulse, oilseeds and vegetables in upland condition and take up early sowing/transplanting.
- Before sowing of any crop, farmers are advised for application of well decomposed FYM or compost, it increases the water use efficiency as well as nutrient quantity in the soil. Farmers are also advised to increase the use of potash, it helps in minimizing the losses due to the moisture stress during dry spell

Kharif Rice:

The crop is at tillering to active tillering stage. Maintain about 2-3 cm standing water during active tillering stage. At 21 days after transplanting apply 50% N (35 kg Urea/acre). Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.

Weather based Disease and pest

- In early transplanted/direct sown crop, chance for leaf folder infestation due to the prevalence of cloudy weather. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days.
- Due to forecasted increase in temperature there may be increase in infestation of thrips, to control this Spray Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.
- Due to high temperature and high humidity there will be chances of infestation of bacterial leaf streak, to control apply Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.

Maize:

Past week high humidity (>90%) and decrease in temperature (21-33°C) and continuous rain the past week may aggravate downy mildew disease in Maize. High humid conditions, temperature between 18-27 Deg. Celsius favors the occurrence of leaf blight in Maize. To control both disease spray Azoxystrobin + Mancozeb @ 300 g /ac. Due to water logging condition stem rot may be found, then farmers should apply carbendazim @ 2g/ltr water. Due to forecasted hot and humid weather, there may be Aphids and Thrips attack, then apply Dimethoate 30EC @ 400ml/Acre.

VEGETABLES:

- Due to past rain, Chances of fruit rot infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Past weather may aggravate fruit rot in pointed gourd, to control it remove damaged fruit from the field and drain out excess water. Spray alternately Copper Oxychloride 50% WP (4 g/litre of water) and Metalaxyl 8% + Mancozeb 64% (3 g/litre of water).
- Past rainy weather is favorable for Root and stem rot in cucurbits. To control farmers may apply Thiophanate Methyl 70% WP @ 2 gm/l or Azoxystrobin 18.2 SC + Difeconazole 11.11% SC 1.5 gm/l to be applied base as well as root zone.
- At present chilli crop at nursery to transplanting stage. Past weather conditions are favourable for damping off in chilli nurseries. To control it by drenching of copper oxy chloride @ 3 g per litre of water (or) 1 % Bordeaux mixture, two times at an interval of 7 days. Farmers are advised to dip the seedlings in 5 ml imidachloprid and 10 g carbendazim solution per 10 litres of water for 5 minutes while transplanting.

ANIMAL HUSBANDRY:

Keep animal shed and shelter house free from water logging by keeping it dry and free from weed. Fresh and immature leaves should not to be fed more amount as they may cause diarrhea. If grasses were cut in morning or evening time, it can be sun dried for about 30 minute before feeding.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-647
Date- 01/09/2026

AGROMET ADVISORY BULLETIN FOR TIGIRIA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 06 mm and district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.28°C (-0.72) and 23.57°C (-0.63), respectively. The mean relative humidity was 88-92%.

Forecasted period: Today(8.30IST) to 06th September (8.30IST)2026

Weatheradvisory: *Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block.* The day temperature will be 31-33°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 85-92% and 50-75% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Tigiria (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	01/09/2026	02/09/2026	03/09/2026	04/09/2026	05/09/2026
Rainfall(mm)	12	9	8	8	7
T-Max(°C)	32	31	32	33	33
T-Min(°C)	25	25	25	26	26
Cloud Cover(octa)	6	6	6	5	5
RH Max (%)	92	87	89	87	85
RH Min (%)	75	64	66	50	52
Wind Speed (Kmph)	10	4	4	4	4
Wind Direction (deg)	287	243	248	264	315

Agromet Advisory

General advisory:

- Resume spraying and fertilization in dry days based upon now-cast warning (before 2-3 hrs).
- By using recent rainfall and soil moisture start field preparation for non-paddy crops like pulse, oilseeds and vegetables in upland condition and take up early sowing/transplanting.
- Before sowing of any crop, farmers are advised for application of well decomposed FYM or compost, it increases the water use efficiency as well as nutrient quantity in the soil. Farmers are also advised to increase the use of potash, it helps in minimizing the losses due to the moisture stress during dry spell

Kharif Rice:

The crop is at tillering to active tillering stage. Maintain about 2-3 cm standing water during active tillering stage. At 21 days after transplanting apply 50% N (35 kg Urea/acre). Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.

Weather based Disease and pest

- *In early transplanted/direct sown crop, chance for leaf folder infestation due to the prevalence of cloudy weather. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days.*
- Due to forecasted increase in temperature there may be increase in infestation of thrips, to control this Spray Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.
- Due to high temperature and high humidity there will be chances of infestation of bacterial leaf streak, to control apply Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.

Maize:

Past week high humidity (>90%) and decrease in temperature (21-33°C) and continuous rain the past week may aggravate downy mildew disease in Maize. High humid conditions, temperature between 18-27 Deg. Celsius favors the occurrence of leaf blight in Maize. To control both disease spray Azoxystrobin + Mancozeb @ 300 g /ac. Due to water logging condition stem rot may be found, then farmers should apply carbendazim @ 2g/ltr water. Due to forecasted hot and humid weather, there may be Aphids and Thrips attack, then apply Dimethoate 30EC @ 400ml/Acre.

VEGETABLES:

- Due to past rain, Chances of fruit rot infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Past weather may aggravate fruit rot in pointed gourd, to control it remove damaged fruit from the field and drain out excess water. Spray alternately Copper Oxychloride 50% WP (4 g/litre of water) and Metalaxyl 8% + Mancozeb 64% (3 g/litre of water).
- Past rainy weather is favorable for Root and stem rot in cucurbits. To control farmers may apply Thiophanate Methyl 70% WP @ 2 gm/l or Azoxystrobin 18.2 SC + Difeconazole 11.11% SC 1.5 gm/l to be applied base as well as root zone.
- At present chilli crop at nursery to transplanting stage. Past weather conditions are favourable for damping off in chilli nurseries. To control it by drenching of copper oxy chloride @ 3 g per litre of water (or) 1 % Bordeaux mixture, two times at an interval of 7 days. Farmers are advised to dip the seedlings in 5 ml imidachloprid and 10 g carbendazim solution per 10 litres of water for 5 minutes while transplanting.

ANIMAL HUSBANDRY:

Keep animal shed and shelter house free from water logging by keeping it dry and free from weed. Fresh and immature leaves should not to be fed more amount as they may cause diarrhea. If grasses were cut in morning or evening time, it can be sun dried for about 30 minute before feeding.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area







GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-647
Date- 01/09/2026

				
MEGHDOOT APP	DAMINI APP	FACEBOOK (Damu Cuttack)	BLOCK SPECIFIC WHAT'S APP GROUPS	@DAMUCUTTACK

- Kindly install and use **“Meghdoot mobile App”** for Cuttack district weather forecast, warnings and weather based agromet advisory services.
<https://play.google.com/store/apps/details?id=com.aas.meghdoot>
- Kindly install and use **“Damini mobile App”** for location specific for lightening forecast and warnings.
<https://play.google.com/store/apps/details?id=com.lightening.live.damini>
- Follow the **block specific whats app groups** and join in your block wise groups for getting weather based agromet advisory services twice in a week (Tuesday and Friday) and other weather warning based farm management advisories.
- Follow the **Official Website links** to get the weather based agromet advisory service bulletins (**English & Odia**) twice in a week (Tuesday and Friday).

1	ICAR-NRRI, Cuttack, Odisha	https://icar-nrri.in/agromate/	
2	Agrimet division, IMD, Pune	https://imdagrmet.gov.in/imdproject/AGIndex.php	



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

