




Cuttack District Block Wise Agro-met Advisory Service Bulletin

*Tuesday, 29th September, 2026
(For the period 29th September 03rd
October, 2026)*



*Issued by
District Agromet Unit,
KrishiVigyan Kendra Cuttack*







GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-651
Date- 29/09/2026

STANDARD PRECIPITATION INDEX AND NDVI MAP

Standard Precipitation Index (SPI)	Normalized difference vegetation Index (NDVI)
MINISTRY OF EARTH SCIENCES INDIA METEOROLOGICAL DEPARTMENT HYDROMET SECTION, CAUI, CRS PUNE STANDARDIZED PRECIPITATION INDEX (SPI) RECENT 4 WEEK 27 AUGUST 2026 to 23 SEPTEMBER 2026  As per SPI value (cumulative 4weeks -27 August to 23 September, 2026), Mildly Dry condition was there in Cuttack district.	NOAA/VIIRS/BLENDED NDVI Composite ending on Week no 38 (17.09.2026 to 23.09.2026) over Agricultural region Odisha  Agriculture vigour is moderate at many places in the state. According to NDVI map (17.09.2026 to 23.09.2026) the agriculture vigor was light to moderate over most places of Cuttack district.

LEGEND (Medium range block level rainfall forecast)		
	No Rainfall	00 mm
	Very light rainfall	01-05 mm
	Light rainfall	06-10 mm
	Light to moderate rainfall	11-30 mm
	Moderate to heavy rainfall	31-65 mm
	Heavy to very heavy rainfall	66-115 mm
	Extremely heavy rainfall	>116 mm



Dr Ranjan Kumar Mohanta
(Nodal officer)

**Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area**





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-651
Date- 29/09/2026

Weather (Past and Forecasted) based Disease and pest in Major Crops and vegetables

Sl No	Crop	Stage and condition of the crop	Anticipated Disease and pest	
1	Rice	Panicle initiation		Leaf Streak
				Brown plant hopper
				Gall mildge
				Mealy bug
				Hispa
				Caseworm
2	Sugarcane	6-7 month old		Mealy bug
				Scales
				White fly



Dr Ranjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-651
Date- 29/09/2026

3	Maize	Cob initiation		Aphid/thrips
				Stem borer
				Tobacco caterpillar
4	Vegetables			Aphid
				Leaf hopper
				White fly
5	Banana			Sigatoka
				Panama wilt



Dr Ranjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-651
Date- 29/09/2026

AGROMET ADVISORY BULLETIN FOR ATHAGAD

District Past week average weather condition and Block past week average Rainfall condition

The block received average 25 mm and district received average 33 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.57°C (-2.54) and 23°C (-0.82), respectively. The mean relative humidity was 88-95%.

Forecasted period: Today(8.30IST) to 04th October (8.30IST)2026

Weather advisory: Very light to light rainfall/thundershower is forecasted for today to Saturday in the block. The day temperature will be 35-36°C and night temperature will be 24-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 85-90% and 46-53% respectively. **There will be partly cloudy to overcast sky for next 5days.**

Block	Athagad (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	29/09/2026	30/09/2026	01/10/2026	02/10/2026	03/10/2026
Rainfall (mm)	4	5	5	4	3
T-Max (°C)	35	35	36	36	36
T-Min (°C)	24	25	25	25	26
Cloud Cover (octa)	3	4	4	3	4
RH Max (%)	88	85	88	88	90
RH Min (%)	50	48	51	53	46
Wind Speed (kmph)	7	6	4	3	4
Wind Direction (deg)	321	303	301	264	232

Agromet Advisory

General advisory:

Drain out excess rain water from crop fields. Farmers should continue pesticide and fertilizer application activities in between upcoming 5 days based upon rainfall forecast (Now cast). Start sowing of non-paddy crops and vegetables after rain to utilize the rain water.

Kharif Rice

Maintain about 5-10cm standing water during panicle initiation stage. In normal transplanted rice, apply second top dressing of fertilizer at Panicle Initiation stage with 17kg of urea and 13kg MOP per acre in upcoming dry days based upon rainfall forecast.

Weather based Disease and pest of rice

- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Make a gape after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest.
- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5%EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 ltr water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one-acre area.

Maize:

- Maize is at cob initiation stage. There may be **Aphids and Thrips** infestation, then Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre. To control **stem borer** apply fipronil 5SC 400ml/acre in 200 litre water. There may be chances of **tobacco caterpillar** in maize in between temperature 15-35 °C. To control the pest apply Thiodicarb 75% WP 400gram/acre in 200 liter water.

VEGETABLES:

- **Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.** Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field. Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.. Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm. Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.

Fruits:

- Banana Present weather conditions are favourable for the incidence and spread of sigatoka leaf spot disease in banana. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2 -3 times in 25 days interval. Go for Drenching of Carbendazim 0.2% or Propiconazole 0.5 ml/ litre once in a month for panama wilt of Banana disease.

Animal husbandry:

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



Dr Ranjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-651
Date- 29/09/2026

AGROMET ADVISORY BULLETIN FOR BADAMBA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 72 mm and district received average 33 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.57°C (-2.54) and 23°C (-0.82), respectively. The mean relative humidity was 88-95%.

Forecasted period: Today(8.30IST) to 04th October (8.30IST)2026

Weather advisory *Very light to light rainfall/thundershower is forecasted for today to Saturday in the block.* The day temperature will be 35-36°C and night temperature will be 24-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 87-93% and 43-53% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Badamba (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	29/09/2026	30/09/2026	01/10/2026	02/10/2026	03/10/2026
Rainfall(mm)	3	4	4	3	3
T-Max(°C)	35	35	35	36	36
T-Min(°C)	24	25	25	25	26
Cloud Cover(octa)	2	4	4	3	4
RH Max (%)	89	87	90	93	93
RH Min (%)	53	48	46	46	43
Wind Speed (Kmph)	9	6	3	3	3
Wind Direction (deg)	323	306	283	256	230

Agromet Advisory

General advisory:

Drain out excess rain water from crop fields. Farmers should continue pesticide and fertilizer application activities in between upcoming 5 days based upon rainfall forecast (Now cast). Start sowing of non-paddy crops and vegetables after rain to utilize the rain water.

Kharif Rice

Maintain about 5-10cm standing water during panicle initiation stage. In normal transplanted rice, apply second top dressing of fertilizer at Panicle Initiation stage with 17kg of urea and 13kg MOP per acre in upcoming dry days based upon rainfall forecast.

Weather based Disease and pest of rice

- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Make a gape after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest.
- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200ltr water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5%EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 ltr water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one-acre area.

Maize:

- Maize is at cob initiation stage. There may be **Aphids and Thrips** infestation, then Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200ltr water for one acre. To control **stem borer** apply fipronil 5SC 400ml/acre in 200 ltr water. There may be chances of **tobacco caterpillar** in maize in between temperature 15-35 °C. To control the pest apply Thiodicarb 75% WP 400gram/acre in 200 liter water.

VEGETABLES:

- **Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.** Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field. Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.. Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm. Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.

Fruits:

- Banana Present weather conditions are favourable for the incidence and spread of sigatoka leaf spot disease in banana. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2 -3 times in 25 days interval. Go for Drenching of Carbendizim 0.2% or Propiconazole 0.5 ml/ litre once in a month for panama wilt of Banana disease.

Animal husbandry:

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



Dr Ranjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-651
Date- 29/09/2026

AGROMET ADVISORY BULLETIN FOR BANKI

District Past week average weather condition

The district received average 33 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.57°C (-2.54) and 23°C (-0.82), respectively. The mean relative humidity was 88-95%.

Forecasted period: Today(8.30IST) to 04th October (8.30IST)2026

Weather advisory: Very light to light rainfall/thundershower is forecasted for today to Saturday in the block. The day temperature will be 35-36°C and night temperature will be 24-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 85-90% and 45-51% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Banki (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	29/09/2026	30/09/2026	01/10/2026	02/10/2026	03/10/2026
Rainfall(mm)	4	4	4	4	3
T-Max(°C)	35	35	35	36	36
T-Min(°C)	24	25	25	26	26
Cloud Cover(octa)	2	4	4	3	4
RH Max (%)	87	85	87	88	90
RH Min (%)	51	47	49	51	45
Wind Speed (Kmph)	8	6	4	3	4
Wind Direction (deg)	323	305	294	270	236

Agromet Advisory

General advisory:

Drain out excess rain water from crop fields. Farmers should continue pesticide and fertilizer application activities in between upcoming 5 days based upon rainfall forecast (Now cast). Start sowing of non-paddy crops and vegetables after rain to utilize the rain water.

Kharif Rice

Maintain about 5-10cm standing water during panicle initiation stage. In normal transplanted rice, apply second top dressing of fertilizer at Panicle Initiation stage with 17kg of urea and 13kg MOP per acre in upcoming dry days based upon rainfall forecast.

Weather based Disease and pest of rice

- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Make a gape after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest.
- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5%EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 litre water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one-acre area.

Maize:

- Maize is at cob initiation stage. There may be **Aphids and Thrips** infestation, then Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre. To control **stem borer** apply fipronil 5SC 400ml/acre in 200 litre water. There may be chances of **tobacco caterpillar** in maize in between temperature 15-35 °C. To control the pest apply Thiodicarb 75% WP 400gram/acre in 200 litre water.

VEGETABLES:

- **Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.** Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field. Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.. Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm. Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.

Fruits:

- Banana Present weather conditions are favourable for the incidence and spread of sigatoka leaf spot disease in banana. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2 -3 times in 25 days interval. Go for Drenching of Carbendazim 0.2% or Propiconazole 0.5 ml/ litre once in a month for panama wilt of Banana disease.

Animal husbandry:

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



Dr Ranjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNITKRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.comBulletin No.-651
Date- 29/09/2026

AGROMET ADVISORY BULLETIN FOR BANKI-DAMPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 42 mm and district received average 33 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.57°C (-2.54) and 23°C (-0.82), respectively. The mean relative humidity was 88-95%.

Forecasted period: Today(8.30IST) to 04th October (8.30IST)2026

Weather Advisory: Very light to light rainfall/thundershower is forecasted for today to Saturday in the block. The day temperature will be 35-36°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 85-90% and 45-51% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Banki-Dampada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	29/09/2026	30/09/2026	01/10/2026	02/10/2026	03/10/2026
Rainfall(mm)	4	5	4	4	4
T-Max(°C)	35	35	35	36	36
T-Min(°C)	25	25	25	26	26
Cloud Cover(octa)	3	4	5	3	4
RH Max (%)	87	85	87	88	90
RH Min (%)	51	47	49	51	45
Wind Speed (Kmph)	8	6	4	3	4
Wind Direction (deg)	323	305	294	270	236

Agromet Advisory

General advisory:

Drain out excess rain water from crop fields. Farmers should continue pesticide and fertilizer application activities in between upcoming 5 days based upon rainfall forecast (Now cast). Start sowing of non-paddy crops and vegetables after rain to utilize the rain water.

Kharif Rice

Maintain about 5-10cm standing water during panicle initiation stage. In normal transplanted rice, apply second top dressing of fertilizer at Panicle Initiation stage with 17kg of urea and 13kg MOP per acre in upcoming dry days based upon rainfall forecast.

Weather based Disease and pest of rice

- If **Brown Plant Hopper (BPH)** infestation will observe in rice, go for alternate wetting and drying. Make a gape after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest.
- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5%EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 ltr water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one-acre area.

Maize:

- Maize is at cob initiation stage. There may be **Aphids and Thrips** infestation, then Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre. To control **stem borer** apply fipronil 5SC 400ml/acre in 200 litre water. There may be chances of **tobacco caterpillar** in maize in between temperature 15-35 °C. To control the pest apply Thiodicarb 75% WP 400gram/acre in 200 liter water.

VEGETABLES:

- Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.** Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field. Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.. Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm. Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.

Fruits:

- Banana Present weather conditions are favourable for the incidence and spread of sigatoka leaf spot disease in banana. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2 -3 times in 25 days interval. Go for Drenching of Carbendazim 0.2% or Propiconazole 0.5 ml/ litre once in a month for panama wilt of Banana disease.

Animal husbandry:

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.

Dr Ranjan Kumar Mohanta
(Nodal officer)Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-651
Date- 29/09/2026

AGROMET ADVISORY BULLETIN FOR BARANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 25 mm and district received average 33 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.57°C (-2.54) and 23°C (-0.82), respectively. The mean relative humidity was 88-95%.

Forecasted period: Today(8.30IST) to 04th October (8.30IST)2026

Weatheradvisory: Very light to light rainfall/thundershower is forecasted for today to Saturday in the block. The day temperature will be 35-36°C and night temperature will be 24-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 85-94% and 49-52% respectively. **There will be partly cloudy to overcast sky for next 5days.**

Block	Baranga(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	29/09/2026	30/09/2026	01/10/2026	02/10/2026	03/10/2026
Rainfall(mm)	5	6	5	4	4
T-Max(°C)	35	35	36	36	36
T-Min(°C)	24	25	25	26	26
Cloud Cover	3	4	5	3	4
RH Max (%)	89	85	89	93	94
RH Min (%)	52	49	51	50	49
Wind Speed (Kmph)	7	6	5	4	7
Wind Direction (deg)	319	281	247	236	219

Agromet Advisory

General advisory:

Drain out excess rain water from crop fields. Farmers should continue pesticide and fertilizer application activities in between upcoming 5 days based upon rainfall forecast (Now cast). Start sowing of non-paddy crops and vegetables after rain to utilize the rain water.

Kharif Rice

Maintain about 5-10cm standing water during panicle initiation stage. In normal transplanted rice, apply second top dressing of fertilizer at Panicle Initiation stage with 17kg of urea and 13kg MOP per acre in upcoming dry days based upon rainfall forecast.

Weather based Disease and pest of rice

- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Make a gape after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @500ml/acre to manage the pest.
- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5%EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 ltr water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one-acre area.

Maize:

- Maize is at cob initiation stage. There may be **Aphids and Thrips** infestation, then Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre. To control **stem borer** apply fipronil 5SC 400ml/acre in 200 litr water. There may be chances of **tobacco caterpillar** in maize in between temperature 15t-35 °C. To control the pest apply Thiodicarb 75% WP 400gram/acre in 200 liter water.

VEGETABLES:

- **Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.** Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field. Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.. Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm. Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.

Fruits:

- Banana Present weather conditions are favourable for the incidence and spread of sigatoka leaf spot disease in banana. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2 -3 times in 25 days interval. Go for Drenching of Carbendizim 0.2% or Propiconazole 0.5 ml/ litre once in a month for panama wilt of Banana disease.

Animal husbandry:

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



Dr Ranjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-651
Date- 29/09/2026

AGROMET ADVISORY BULLETIN FOR CUTTACK SADAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 22 mm and district received average 33 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.57°C (-2.54) and 23°C (-0.82), respectively. The mean relative humidity was 88-95%.

Forecasted period: Today(8.30IST) to 04th October (8.30IST)2026

Weather advisory: Very light to light rainfall/thundershower is forecasted for today to Saturday in the block. The day temperature will be 35-37°C and night temperature will be 24-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 83-89% and 47-52% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Cuttack sadar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	29/09/2026	30/09/2026	01/10/2026	02/10/2026	03/10/2026
Rainfall(mm)	6	7	6	4	3
T-Max(°C)	35	35	36	36	37
T-Min(°C)	24	25	25	25	26
Cloud Cover(octa)	3	4	5	3	4
RH Max (%)	85	83	84	88	89
RH Min (%)	49	47	49	52	47
Wind Speed (Kmph)	6	6	4	5	6
Wind Direction (deg)	317	284	275	241	227

Agromet Advisory

General advisory:

Drain out excess rain water from crop fields. Farmers should continue pesticide and fertilizer application activities in between upcoming 5 days based upon rainfall forecast (Now cast). Start sowing of non-paddy crops and vegetables after rain to utilize the rain water.

Kharif Rice

Maintain about 5-10cm standing water during panicle initiation stage. In normal transplanted rice, apply second top dressing of fertilizer at Panicle Initiation stage with 17kg of urea and 13kg MOP per acre in upcoming dry days based upon rainfall forecast.

Weather based Disease and pest of rice

- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Make a gape after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest.
- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5%EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 ltr water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one-acre area.

Sugarcane:

- Prevailing hot weather favours the infestation of Mites, Mealy bug, Whitefly and scale insect in sugarcane. If mites will be found the spray spiromesifen 22.9%SC 1ml/litre of water. To control mealy bug spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre. To control scale insects apply Dimethoate 2ml/litre water or methyl demeton 2ml/litre water. To control white fly go for spraying of Thiamethoxam 40 gram in 200litre water for one acre.

VEGETABLES:

- **Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.** Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field. Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.. Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm. Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.

Fruits:

- Banana Present weather conditions are favourable for the incidence and spread of sigatoka leaf spot disease in banana. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2 -3 times in 25 days interval. Go for Drenching of Carbendazim 0.2% or Propiconazole 0.5 ml/ litre once in a month for panama wilt of Banana disease.

Animal husbandry:

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



Dr Ranjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-651
Date- 29/09/2026

AGROMET ADVISORY BULLETIN FOR KANTAPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 22 mm and district received average 33 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.57°C (-2.54) and 23°C (-0.82), respectively. The mean relative humidity was 88-95%.

Forecasted period: Today(8.30IST) to 04th October (8.30IST)2026

Weather advisory: Very light to light rainfall/thundershower is forecasted for today to Saturday in the block. The day temperature will be 35-36°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 84-90% and 49-57% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Kantapada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	29/09/2026	30/09/2026	01/10/2026	02/10/2026	03/10/2026
Rainfall(mm)	5	6	5	4	4
T-Max(°C)	35	35	36	36	36
T-Min(°C)	25	25	26	26	26
Cloud Cover(octa)	3	5	5	3	4
RH Max (%)	86	84	86	89	90
RH Min (%)	52	49	53	57	52
Wind Speed (Kmph)	7	7	5	6	8
Wind Direction (deg)	317	273	249	235	223

Agromet Advisory

General advisory:

Drain out excess rain water from crop fields. Farmers should continue pesticide and fertilizer application activities in between upcoming 5 days based upon rainfall forecast (Now cast). Start sowing of non-paddy crops and vegetables after rain to utilize the rain water.

Kharif Rice

Maintain about 5-10cm standing water during panicle initiation stage. In normal transplanted rice, apply second top dressing of fertilizer at Panicle Initiation stage with 17kg of urea and 13kg MOP per acre in upcoming dry days based upon rainfall forecast.

Weather based Disease and pest of rice

- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Make a gape after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest.
- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5%EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 ltr water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one-acre area.

Sugarcane:

- Prevailing hot weather favours the infestation of Mites, Mealy bug, Whitefly and scale insect in sugarcane. If mites will be found the spray spiromesifen 22.9%SC 1ml/litre of water. To control mealy bug spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre. To control scale insects apply Dimethoate 2ml/litre water or methyl demeton 2ml/litre water. To control white fly go for spraying of Thiamethoxam 40 gram in 200litre water for one acre.

VEGETABLES:

- Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.** Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field. Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.. Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm. Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.

Fruits:

- Banana Present weather conditions are favourable for the incidence and spread of sigatoka leaf spot disease in banana. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2 -3 times in 25 days interval. Go for Drenching of Carbendazim 0.2% or Propiconazole 0.5 ml/ litre once in a month for panama wilt of Banana disease.

Animal husbandry:

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



Dr Ranjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-651
Date- 29/09/2026

AGROMET ADVISORY BULLETIN FOR MAHANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 40 mm and district received average 33 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.57°C (-2.54) and 23°C (-0.82), respectively. The mean relative humidity was 88-95%.

Forecasted period: Today(8.30IST) to 04th October (8.30IST)2026

Weather advisory: Very light to light rainfall/thundershower is forecasted for today to Saturday in the block. The day temperature will be and it will be 34-36°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 83-89% and 48-56 respectively. **There will be partly cloudy to overcast sky for next 5days.**

Block	Mahanga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	29/09/2026	30/09/2026	01/10/2026	02/10/2026	03/10/2026
Rainfall(mm)	5	5	6	4	4
T-Max(°C)	34	35	35	35	36
T-Min(°C)	25	25	26	26	26
Cloud Cover(octal)	3	4	4	3	4
RH Max (%)	87	83	85	88	89
RH Min (%)	48	48	51	56	48
Wind Speed (Kmph)	6	7	6	5	8
Wind Direction (deg)	303	270	243	222	225

Agromet Advisory

General advisory:

Drain out excess rain water from crop fields. Farmers should continue pesticide and fertilizer application activities in between upcoming 5 days based upon rainfall forecast (Now cast). Start sowing of non-paddy crops and vegetables after rain to utilize the rain water.

Kharif Rice

Maintain about 5-10cm standing water during panicle initiation stage. In normal transplanted rice, apply second top dressing of fertilizer at Panicle Initiation stage with 17kg of urea and 13kg MOP per acre in upcoming dry days based upon rainfall forecast.

Weather based Disease and pest of rice

- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Make a gape after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest.
- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5%EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 ltr water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one-acre area.

Sugarcane:

- Prevailing hot weather favours the infestation of Mites, Mealy bug, Whitefly and scale insect in sugarcane. If mites will be found the spray spiromesifen 22.9%SC 1ml/litre of water. To control mealy bug spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre. To control scale insects apply Dimethoate 2ml/litre water or methyl demeton 2ml/litre water. To control white fly go for spraying of Thiamethoxam 40 gram in 200litre water for one acre.

VEGETABLES:

- **Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.** Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field. Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.. Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm. Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.

Fruits:

- Banana Present weather conditions are favourable for the incidence and spread of sigatoka leaf spot disease in banana. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2 -3 times in 25 days interval. Go for Drenching of Carbendazim 0.2% or Propiconazole 0.5 ml/ litre once in a month for panama wilt of Banana disease.

Animal husbandry:

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



Dr Ranjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-651
Date- 29/09/2026

AGROMET ADVISORY BULLETIN FOR NARASINGHPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 24 mm and district received average 33 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.57°C (-2.54) and 23°C (-0.82), respectively. The mean relative humidity was 88-95%.

Forecasted period: Today(8.30IST) to 04th October (8.30IST)2026

Weatheradvisory: Very light to light rainfall/thundershower is forecasted for today to Saturday in the block. The day temperature will be 34-36°C and night temperature will be 24-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 89-94% and 46-56% respectively. **There will be partly cloudy to overcast sky for next 5days.**

Block	Narasinghpur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	29/09/2026	30/09/2026	01/10/2026	02/10/2026	03/10/2026
Rainfall(mm)	3	3	3	3	2
T-Max(°C)	34	35	35	36	36
T-Min(°C)	24	25	25	25	26
Cloud Cover(octa)	2	3	4	3	3
RH Max (%)	94	89	90	93	94
RH Min (%)	56	50	49	51	46
Wind Speed (Kmph)	8	6	4	3	3
Wind Direction (deg)	315	301	304	286	278

Agromet Advisory

General advisory:

Drain out excess rain water from crop fields. Farmers should continue pesticide and fertilizer application activities in between upcoming 5 days based upon rainfall forecast (Now cast). Start sowing of non-paddy crops and vegetables after rain to utilize the rain water.

Kharif Rice

Maintain about 5-10cm standing water during panicle initiation stage. In normal transplanted rice, apply second top dressing of fertilizer at Panicle Initiation stage with 17kg of urea and 13kg MOP per acre in upcoming dry days based upon rainfall forecast.

Weather based Disease and pest of rice

- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Make a gape after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest.
- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litr water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5%EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 ltr water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one-acre area.

Maize:

- Maize is at cob initiation stage. There may be **Aphids and Thrips** infestation, then Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litr water for one acre. To control **stem borer** apply fipronil 5SC 400ml/acre in 200 ltr water. There may be chances of **tobacco caterpillar** in maize in between temperature 15t-35 °C. To control the pest apply Thiodicarb 75% WP 400gram/acre in 200 liter water.

VEGETABLES:

- **Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.** Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field. Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.. Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm. Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.

Fruits:

- Banana Present weather conditions are favourable for the incidence and spread of sigatoka leaf spot disease in banana. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2 -3 times in 25 days interval. Go for Drenching of Carbendizim 0.2% or Propiconazole 0.5 ml/ litre once in a month for panama wilt of Banana disease.

Animal husbandry:

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



Dr Ranjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-651
Date- 29/09/2026

AGROMET ADVISORY BULLETIN FOR NIALI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 29 mm and district received average 33 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.57°C (-2.54) and 23°C (-0.82), respectively. The mean relative humidity was 88-95%.

Forecasted period: Today(8.30IST) to 04th October (8.30IST)2026

Weather advisory: Very light to light rainfall/thundershower is forecasted for today to Saturday in the block. The day temperature will be 35-36°C and night temperature will be 25-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 89-94% and 51-55% respectively.

There will be partly cloudy to overcast sky for next 5 days.

Block	Niali (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	29/09/2026	30/09/2026	01/10/2026	02/10/2026	03/10/2026
Rainfall(mm)	5	6	6	4	5
T-Max(°C)	35	35	36	36	36
T-Min(°C)	25	26	26	26	27
Cloud Cover(octa)	3	5	5	3	4
RH Max (%)	90	89	93	94	94
RH Min (%)	55	51	55	55	54
Wind Speed(Kmph)	7	7	6	5	9
Wind Direction(deg)	311	259	243	242	225

Agromet Advisory

General advisory:

Drain out excess rain water from crop fields. Farmers should continue pesticide and fertilizer application activities in between upcoming 5 days based upon rainfall forecast (Now cast). Start sowing of non-paddy crops and vegetables after rain to utilize the rain water.

Kharif Rice

Maintain about 5-10cm standing water during panicle initiation stage. In normal transplanted rice, apply second top dressing of fertilizer at Panicle Initiation stage with 17kg of urea and 13kg MOP per acre in upcoming dry days based upon rainfall forecast.

Weather based Disease and pest of rice

- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Make a gape after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest.
- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5%EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 ltr water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one-acre area.

Sugarcane:

- Prevailing hot weather favours the infestation of Mites, Mealy bug, Whitefly and scale insect in sugarcane. If mites will be found the spray spiromesifen 22.9%SC 1ml/litre of water. To control mealy bug spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre. To control scale insects apply Dimethoate 2ml/litre water or methyl demeton 2ml/litre water. To control white fly go for spraying of Thiamethoxam 40 gram in 200litre water for one acre.

VEGETABLES:

- **Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.** Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field. Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.. Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm. Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.

Fruits:

- Banana Present weather conditions are favourable for the incidence and spread of sigatoka leaf spot disease in banana. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2 -3 times in 25 days interval. Go for Drenching of Carbendizim 0.2% or Propiconazole 0.5 ml/ litre once in a month for panama wilt of Banana disease.

Animal husbandry:

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



Dr Ranjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-651
Date- 29/09/2026

AGROMET ADVISORY BULLETIN FOR NISCHINTAKOILI
District Past week average weather condition and Block past week average Rainfall condition

The block received average 40 mm and district received average 33 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.57°C (-2.54) and 23°C (-0.82), respectively. The mean relative humidity was 88-95%.

Forecasted period: Today(8.30IST) to 04th October (8.30IST)2026

Weatheradvisory: *Very light to light rainfall/thundershower is forecasted for today to Saturday in the block.* The day temperature will be 35-36°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 84-89% and 48-56% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Nischintakoili (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	29/09/2026	30/09/2026	01/10/2026	02/10/2026	03/10/2026
Rainfall(mm)	5	6	6	4	4
T-Max(°C)	35	35	35	36	36
T-Min(°C)	25	25	26	26	26
Cloud Cover(octa)	3	5	5	3	4
RH Max (%)	87	84	86	89	89
RH Min (%)	48	48	51	56	49
Wind Speed (Kmph)	6	7	6	5	9
Wind Direction (deg)	303	267	240	222	225

Agromet Advisory

General advisory:

Drain out excess rain water from crop fields. Farmers should continue pesticide and fertilizer application activities in between upcoming 5 days based upon rainfall forecast (Now cast). Start sowing of non-paddy crops and vegetables after rain to utilize the rain water.

Kharif Rice

Maintain about 5-10cm standing water during panicle initiation stage. In normal transplanted rice, apply second top dressing of fertilizer at Panicle Initiation stage with 17kg of urea and 13kg MOP per acre in upcoming dry days based upon rainfall forecast.

Weather based Disease and pest of rice

- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Make a gape after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest.
- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5%EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 ltr water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one-acre area.

Sugarcane:

- Prevailing hot weather favours the infestation of Mites, Mealy bug, Whitefly and scale insect in sugarcane. If mites will be found the spray spiromesifen 22.9%SC 1ml/litre of water. To control mealy bug spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre. To control scale insects apply Dimethoate 2ml/litre water or methyl demeton 2ml/litre water. To control white fly go for spraying of Thiamethoxam 40 gram in 200litre water for one acre.

VEGETABLES:

- **Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.** Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field. Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.. Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm. Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.

Fruits:

- Banana Present weather conditions are favourable for the incidence and spread of sigatoka leaf spot disease in banana. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2 -3 times in 25 days interval. Go for Drenching of Carbendazim 0.2% or Propiconazole 0.5 ml/ litre once in a month for panama wilt of Banana disease.

Animal husbandry:

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



Dr Ranjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-651
Date- 29/09/2026

AGROMET ADVISORY BULLETIN FOR SALEPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 30 mm and district received average 33 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.57°C (-2.54) and 23°C (-0.82), respectively. The mean relative humidity was 88-95%.

Forecasted period: Today(8.30IST) to 04th October (8.30IST)2026

Weather advisory: Very light to light rainfall/thundershower is forecasted for today to Saturday in the block. The day temperature will be 35-36°C and night temperature will be 24-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 83-89% and 48-56% respectively. **There will be partly cloudy to overcast sky for next 5days.**

Block	Salepur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	29/09/2026	30/09/2026	01/10/2026	02/10/2026	03/10/2026
Rainfall(mm)	6	7	6	4	4
T-Max(°C)	35	35	36	36	36
T-Min(°C)	24	25	25	26	26
Cloud Cover(octa)	3	4	5	3	4
RH Max (%)	87	83	85	88	89
RH Min (%)	48	48	51	56	49
Wind Speed (Kmph)	6	7	5	5	8
Wind Direction (deg)	306	273	242	225	225

Agromet Advisory

General advisory:

Drain out excess rain water from crop fields. Farmers should continue pesticide and fertilizer application activities in between upcoming 5 days based upon rainfall forecast (Now cast). Start sowing of non-paddy crops and vegetables after rain to utilize the rain water.

Kharif Rice

Maintain about 5-10cm standing water during panicle initiation stage. In normal transplanted rice, apply second top dressing of fertilizer at Panicle Initiation stage with 17kg of urea and 13kg MOP per acre in upcoming dry days based upon rainfall forecast.

Weather based Disease and pest of rice

- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Make a gape after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest.
- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5%EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 litre water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one-acre area.

Maize:

- Maize is at cob initiation stage. There may be **Aphids and Thrips** infestation, then Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre. To control **stem borer** apply fipronil 5SC 400ml/acre in 200 litre water. There may be chances of **tobacco caterpillar** in maize in between temperature 15-35 °C. To control the pest apply Thiodicarb 75% WP 400gram/acre in 200 liter water.

VEGETABLES:

- **Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.** Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field. Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.. Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm. Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.

Fruits:

- Banana Present weather conditions are favourable for the incidence and spread of sigatoka leaf spot disease in banana. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2 -3 times in 25 days interval. Go for Drenching of Carbendazim 0.2% or Propiconazole 0.5 ml/ litre once in a month for panama wilt of Banana disease.

Animal husbandry:

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



Dr Ranjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-651
Date- 29/09/2026

AGROMET ADVISORY BULLETIN FOR TANGI-CHOUDWAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 33 mm and district received average 33 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.57°C (-2.54) and 23°C (-0.82), respectively. The mean relative humidity was 88-95%.

Forecasted period: Today(8.30IST) to 04th October (8.30IST)2026

Weather advisory: Very light to light rainfall/thundershower is forecasted for today to Saturday in the block. The day temperature will be 35-37°C and night temperature will be 24-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 83-86% and 56-63% respectively. **There will be partly cloudy to overcast sky for next 5days.**

Block	Tangi-Choudwar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	29/09/2026	30/09/2026	01/10/2026	02/10/2026	03/10/2026
Rainfall(mm)	6	6	6	4	3
T-Max(°C)	35	35	36	36	37
T-Min(°C)	24	25	25	25	26
Cloud Cover(octa)	3	4	5	3	4
RH Max (%)	84	83	84	86	85
RH Min (%)	56	56	62	63	60
Wind Speed (Kmph)	8	11	9	9	13
Wind Direction (deg)	300	257	243	235	227

Agromet Advisory

General advisory:

Drain out excess rain water from crop fields. Farmers should continue pesticide and fertilizer application activities in between upcoming 5 days based upon rainfall forecast (Now cast). Start sowing of non-paddy crops and vegetables after rain to utilize the rain water.

Kharif Rice

Maintain about 5-10cm standing water during panicle initiation stage. In normal transplanted rice, apply second top dressing of fertilizer at Panicle Initiation stage with 17kg of urea and 13kg MOP per acre in upcoming dry days based upon rainfall forecast.

Weather based Disease and pest of rice

- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Make a gape after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest.
- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200ltr water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5%EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 ltr water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one-acre area.

Maize:

- Maize is at cob initiation stage. There may be **Aphids and Thrips** infestation, then Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200ltr water for one acre. To control **stem borer** apply fipronil 5SC 400ml/acre in 200 ltr water. There may be chances of **tobacco caterpillar** in maize in between temperature 15t-35 °C. To control the pest apply Thiodicarb 75% WP 400gram/acre in 200 liter water.

VEGETABLES:

- **Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.** Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field. Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.. Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm. Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.

Fruits:

- Banana Present weather conditions are favourable for the incidence and spread of sigatoka leaf spot disease in banana. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2 -3 times in 25 days interval. Go for Drenching of Carbendizim 0.2% or Propiconazole 0.5 ml/ litre once in a month for panama wilt of Banana disease.

Animal husbandry:

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



Dr Ranjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-651
Date- 29/09/2026

AGROMET ADVISORY BULLETIN FOR TIGIRIA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 27 mm and district received average 33 mm rainfall during the last week. The mean daily maximum and minimum temperature was 30.57°C (-2.54) and 23°C (-0.82), respectively. The mean relative humidity was 88-95%.

Forecasted period: Today(8.30IST) to 04th October (8.30IST)2026

Weather advisory: Very light to light rainfall/thundershower is forecasted for today to Saturday in the block. The day temperature will be 35-36°C and night temperature will be 24-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 86-91% and 44-51% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Tigiria (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	29/09/2026	30/09/2026	01/10/2026	02/10/2026	03/10/2026
Rainfall(mm)	3	4	4	3	3
T-Max(°C)	35	35	35	36	36
T-Min(°C)	24	25	25	25	26
Cloud Cover(octa)	2	4	4	3	4
RH Max (%)	88	86	88	89	91
RH Min (%)	51	47	49	51	44
Wind Speed (Kmph)	8	6	4	3	3
Wind Direction (deg)	321	308	304	277	243

Agromet Advisory

General advisory:

Drain out excess rain water from crop fields. Farmers should continue pesticide and fertilizer application activities in between upcoming 5 days based upon rainfall forecast (Now cast). Start sowing of non-paddy crops and vegetables after rain to utilize the rain water.

Kharif Rice

Maintain about 5-10cm standing water during panicle initiation stage. In normal transplanted rice, apply second top dressing of fertilizer at Panicle Initiation stage with 17kg of urea and 13kg MOP per acre in upcoming dry days based upon rainfall forecast.

Weather based Disease and pest of rice

- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Make a gap after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest.
- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200ltr water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5%EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 ltr water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one-acre area.

Maize:

- Maize is at cob initiation stage. There may be **Aphids and Thrips** infestation, then Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200ltr water for one acre. To control **stem borer** apply fipronil 5SC 400ml/acre in 200 ltr water. There may be chances of **tobacco caterpillar** in maize in between temperature 15-35 °C. To control the pest apply Thiodicarb 75% WP 400gram/acre in 200 liter water.

VEGETABLES:

- **Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.** Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field. Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.. Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm. Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.

Fruits:

- Banana Present weather conditions are favourable for the incidence and spread of sigatoka leaf spot disease in banana. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2 -3 times in 25 days interval. Go for Drenching of Carbendazim 0.2% or Propiconazole 0.5 ml/ litre once in a month for panama wilt of Banana disease.

Animal husbandry:

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



Dr Ranjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area






GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-651
Date- 29/09/2026

				
MEGHDOOT APP	DAMINI APP	FACEBOOK (Damu Cuttack)	BLOCK SPECIFIC WHAT'S APP GROUPS	@DAMUCUTTACK

- Kindly install and use **“Meghdoot mobile App”** for Cuttack district weather forecast, warnings and weather based agromet advisory services.
<https://play.google.com/store/apps/details?id=com.aas.meghdoot>
- Kindly install and use **“Damini mobile App”** for location specific for lightening forecast and warnings.
<https://play.google.com/store/apps/details?id=com.lightening.live.damini>
- Follow the **block specific whats app groups** and join in your block wise groups for getting weather based agromet advisory services twice in a week (Tuesday and Friday) and other weather warning based farm management advisories.
- Follow the **Official Website links** to get the weather based agromet advisory service bulletins (English & Odia) twice in a week (Tuesday and Friday).

1	ICAR-NRRI, Cuttack, Odisha	https://icar-nrri.in/agromate/	
2	Agrimet division, IMD, Pune	https://imdagrimet.gov.in/imdproject/AGIndex.php	



Dr Ranjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

